

In[]:= **f[x_] := x^3 - I s x^2 + x + I s**

In[]:= **PolynomialRemainder[f[x], f'[x], x]**

$$\text{Out[]} = \frac{10 i s}{9} + \left(\frac{2}{3} + \frac{2 s^2}{9} \right) x$$

- which is not a zero function for all s

In[]:= **r1 = PolynomialRemainder[f[x], f'[x], x];**

PolynomialRemainder[f'[x], r1, x]

Solve[PolynomialRemainder[f'[x], r1, x] == 0, s]

$$\text{Out[]} = 1 - \frac{100 s^2}{27 \left(\frac{2}{3} + \frac{2 s^2}{9} \right)^2} - \frac{20 s^2}{9 \left(\frac{2}{3} + \frac{2 s^2}{9} \right)}$$

$$\text{Out[]} = \left\{ \left\{ s \rightarrow -\sqrt{-\frac{11}{2} + \frac{5\sqrt{5}}{2}} \right\}, \left\{ s \rightarrow \sqrt{-\frac{11}{2} + \frac{5\sqrt{5}}{2}} \right\}, \right. \\ \left. \left\{ s \rightarrow -i\sqrt{\frac{1}{2} (11 + 5\sqrt{5})} \right\}, \left\{ s \rightarrow i\sqrt{\frac{1}{2} (11 + 5\sqrt{5})} \right\} \right\}$$

- Hence, when s is large enough, f[x] has no multiple zeros.
- From the discussion before (2.10), set $\alpha=2\pi/L$ there exists integer $k>l>0$, such that

In[6]:= **$\xi_2 = \xi_1 + k \alpha$;**

$\xi_3 = \xi_1 + l \alpha$;

Collect[($\xi - \xi_1$) ($\xi - \xi_2$) ($\xi - \xi_3$), ξ]

$$\text{Out[8]} = \xi^3 + \xi^2 (-k \alpha - l \alpha - 3 \xi_1) - k l \alpha^2 \xi_1 - k \alpha \xi_1^2 - l \alpha \xi_1^2 - \xi_1^3 + \xi (k l \alpha^2 + 2 k \alpha \xi_1 + 2 l \alpha \xi_1 + 3 \xi_1^2)$$

- **($\xi - \xi_1$) ($\xi - \xi_2$) ($\xi - \xi_3$) = $\xi^3 + I s \xi^2 - \xi + I s$, where in my draft, I use "s" denote the eigenvalue λ . We aim to eliminate s and ξ_1 .**

- Now we know $e_1 = e_2 = e_3 = 0$, which is the same as equation (), where e_1, e_2, e_3 defined as follows

In[12]:= **$e_1 = (-k \alpha - l \alpha - 3 \xi_1) - I s$;**

$e_2 = (k l \alpha^2 + 2 k \alpha \xi_1 + 2 l \alpha \xi_1 + 3 \xi_1^2) + 1$;

$e_3 = -k l \alpha^2 \xi_1 - k \alpha \xi_1^2 - l \alpha \xi_1^2 - \xi_1^3 - I s$;

- Eliminate ξ_1^3 term.

In[17]:= **$e_4 = \text{Collect}[\xi_1 e_2 + 3 e_3, \xi_1]$**

$$\text{Out[17]} = -3 i s + (1 - 2 k l \alpha^2) \xi_1 + (-k \alpha - l \alpha) \xi_1^2$$

- Notice that $e_1=e_2=e_3=0$ is equivalent to $e_1=e_2=e_4=0$, since $(e_4-\xi_1 e_2)/3=e_3$.
- Eliminate ξ_1^2 term in e_3 .

In[23]:= **$e_5 = \text{Collect}[\text{Simplify}[(k+1) \alpha e_2 + 3 e_4], \xi_1]$**

$$\text{Out[23]} = -9 i s + k \alpha + l \alpha + k^2 l \alpha^3 + k l^2 \alpha^3 + (3 + 2 k^2 \alpha^2 - 2 k l \alpha^2 + 2 l^2 \alpha^2) \xi_1$$

- Notice that $e_1=e_2=e_4=0$ is equivalent to $e_1=e_2=e_5$, since $(e_5-(k+l)\alpha e_2)/3=e_4$.

■ Now we solve s from e_1 , and then plug-in e_5 to obtain e_6 .

```
In[31]:= e6 = Collect[Simplify[e5 /. Solve[e1 == 0, s][[1]]], ξ1]
```

```
Out[31]= 10 k α + 10 l α + k^2 l α^3 + k l^2 α^3 + (30 + 2 k^2 α^2 - 2 k l α^2 + 2 l^2 α^2) ξ1
```

■ Now we have $e_2=e_6=0$ is equivalent to $e_1=e_2=e_5=0$.

```
In[37]:= e7 = e2 /. Solve[e6 == 0, ξ1][[1]]
```

```
Out[37]= 1 + k l α^2 +  $\frac{k \alpha (-10 k \alpha - 10 l \alpha - k^2 l \alpha^3 - k l^2 \alpha^3)}{15 + k^2 \alpha^2 - k l \alpha^2 + l^2 \alpha^2}$  +  $\frac{l \alpha (-10 k \alpha - 10 l \alpha - k^2 l \alpha^3 - k l^2 \alpha^3)}{15 + k^2 \alpha^2 - k l \alpha^2 + l^2 \alpha^2}$  +  $\frac{3 (-10 k \alpha - 10 l \alpha - k^2 l \alpha^3 - k l^2 \alpha^3)^2}{4 (15 + k^2 \alpha^2 - k l \alpha^2 + l^2 \alpha^2)^2}$ 
```

```
In[48]:= FullSimplify[-e7 * 4 (15 + k^2 α^2 - k l α^2 + l^2 α^2)^2 / 9]
```

```
Out[48]= -100 + 20 (k^2 - k l + l^2) α^2 + 4 (k^2 - k l + l^2)^2 α^4 + k^2 (k - l)^2 l^2 α^6
```