

MAROTO MEDICAL CENTER · CARDIOLOGY DEPARTMENT

Maroto Medical Center 112 Harbor Bay
Road San Francisco, CA 94107 +1 415 555
0118 · records@maroto.example

Anamnesis & Initial Assessment Report

Document ref: ANAM-2026-04812 · Encounter: 2026-05-20 09:14 · 7 of 9 pages

PATIENT DEMOGRAPHICS				
PATIENT NAME	MRN	DATE OF BIRTH & AGE	INSURANCE	REFERRING CLINICIAN
Reyes, Marcus J.	MR-04812-26	1972-08-14 · 53 y	BlueCare Premier	Dr. A. Patel, GP
Preferred: "Marc"	Maroto Health ID	Sex: Male	Policy #BC-77821-3	Bayview Family Clinic

BLOOD PRESSURE	HEART RATE	BMI	SPO ₂	TEMPERATURE
138 / 88	82 bpm	28.7	97%	36.8°C
. stage 1 hypertension	within normal range	overweight	normal	afebrile

CHIEF COMPLAINT & HISTORY OF PRESENT ILLNESS

Chief complaint: Episodic central chest pressure with mild shortness of breath, worse on exertion. Onset approximately 6 weeks ago, increasing in frequency over the past 10 days.

Patient describes a substernal pressure-like sensation, non-radiating, lasting 5–10 minutes, reproducibly triggered by climbing two flights of stairs or carrying groceries. Pain resolves with rest within 10 minutes. No associated diaphoresis, nausea, or palpitations. No syncope. Sleep is undisturbed. No recent infections or fevers.

RISK FACTOR SCREENING (MODIFIED WHO ROSE QUESTIONNAIRE)	
Q1 Q2 Q3 Q4 Q5 Q6 Q7 Q8 Q1 Q2 Q3 Q4 CORONARY SYMPTOMS Have you ever had pain or discomfort in your chest? YES Patient confirms pressure-like episodes on exertion, see HPI above. Does it ever make you stop walking? YES "I have to slow down and lean on something for a minute." Does the pain go away when you stand still? YES Resolves within 5–10 minutes of rest. Where do you feel the discomfort? Substernal · non-radiating · no jaw / arm involvement.	Q5 Q6 Q7 Q8 Q5 Q6 Q7 Q8 SMOKING Do you currently smoke tobacco? NO Quit 8 years ago. Prior 15 pack-year history (15 cigs/day × 22 years). Q6 Q7 Q8 DIABETES Have you been told you have diabetes? PRE-DIABETIC HbA1c 6.1% in 2025. Not on metformin. Q7 Q8 HYPERTENSION Are you being treated for high blood pressure? YES Lisinopril 10 mg daily since 2023. Q8 FAMILY HISTORY First-degree relative with early cardiac disease? YES Father MI at age 54. Brother CABG at age 58.

COMPOSITE RISK PROFILE

Family history CAD

Hypertension

Pre-diabetic

Former smoker

BMI 28.7

No CKD

No prior MI

No arrhythmia

Active lifestyle (3x/wk walk)

No EtOH abuse

Framingham 10-year CVD risk estimate: 17.4% (high). ASCVD pooled cohort risk: 15.8%. Recommend statin initiation per ACC/AHA 2018 guidelines.

CURRENT MEDICATIONS & ALLERGIES

Active medications

Drug	Dose	Freq	Since
Lisinopril	10 mg	QD	2023-02
Atorvastatin	20 mg	QHS	2024-09
Aspirin (low-dose)	81 mg	QD	2024-12
Vitamin D.	1000 IU	QD	2022-01


Allergies & intolerances

Agent	Reaction	Severity
Penicillin	Urticaria	Severe
Shellfish	Lip swelling	Mild
Latex	Contact rash	Mild

IMAGING & DIAGNOSTIC STUDIES

PA CHEST X-RAY ACQUIRED 2026-05-20 09:38

Cardiothoracic ratio 0.51 — **borderline**



Mild cardiomegaly, no consolidation, lung fields clear. No pleural effusion. Mediastinum not widened.

12-LEAD ECG RESTING 125 MM/S


NSR 78 bpm · **borderline LVH**



Normal sinus rhythm. Mild voltage criteria for left ventricular hypertrophy. No ST elevation, no Q waves.

HOME VITALS TREND LAST 7 DAYS

SOURCE: PATIENT-RECORDED VALIDATED CUFF (OMRON HEM-7156T)



. Systolic BP (mmHg, scale 110–145) . Heart rate (bpm, scale 60–95)

LABORATORY RESULTS (DRAWN 2026-05-20)

Analyte	Result	Unit	Reference range	Flag
Total cholesterol	224	mg/dL	< 200	HIGH
LDL cholesterol	148	mg/dL	< 100	HIGH
HDL cholesterol	38	mg/dL	> 40	LOW
Triglycerides	192	mg/dL	< 150	HIGH
HbA1c	6.1	%	< 5.7	HIGH
Fasting glucose	108	mg/dL	70–99	HIGH
Creatinine	0.94	mg/dL	0.7–1.3	OK
eGFR	92	mL/min	> 60	OK
hs-Troponin I	8	ng/L	< 19	OK
BNP	42	pg/mL	< 100	OK

ASSESSMENT & IMPRESSION

Primary: Probable stable angina pectoris (exertional class II). Multifactorial cardiovascular risk profile dominated by family history, dyslipidemia, and uncontrolled hypertension.

Secondary considerations: Pre-diabetes mellitus (HbA1c 6.1%); overweight (BMI 28.7); borderline LVH on ECG; mild cardiomegaly on CXR — likely chronic hypertensive remodelling.

Reassuring findings: Negative high-sensitivity troponin and BNP — acute coronary syndrome not currently active. Renal function preserved.

RECOMMENDED PLAN

- Confirm ischemia.** Schedule exercise stress test with myocardial perfusion imaging within 14 days. If positive or equivocal, proceed to coronary CT angiography.
- Optimize antihypertensive therapy.** Increase lisinopril to 20 mg QD; recheck BP in 4 weeks. Add amlodipine 5 mg if SBP > 135 at follow-up.
- Intensify lipid therapy.** Atorvastatin 40 mg QHS. Repeat full lipid panel in 8 weeks. Discuss PCSK9 inhibitor if LDL > 100 at recheck.
- Lifestyle counselling.** Mediterranean diet referral to nutrition. Goal: 5% body weight loss over 6 months. Continue 30-min walks 5x/week.
- Pre-diabetes management.** Recheck HbA1c in 3 months. Refer to diabetes prevention program.
- Patient safety net.** Provide chest-pain action card with symptom thresholds for ER presentation. Sublingual nitroglycerin 0.4 mg SL PRN, max x3 with 5-min interval.
- Follow-up cardiology visit.** In 6 weeks (post stress test & lipid recheck).

Clinician statement

I have personally reviewed the history, examined the patient, and discussed the assessment and recommended plan. Patient demonstrated understanding (teach-back verified) and consented to the proposed investigations.

Report prepared automatically from the Maroto Anamnesis questionnaire and finalised by the attending clinician at 2026-05-20 11:02 PT. Document checksum: 6f9e83...b4c1.

ATTENDING PHYSICIAN

Dr. Helena Costa, MD Board-certified
Cardiologist NPI 1235647890 License
#MD-CA-082114

Confidential medical record. This document contains protected health information subject to applicable privacy laws (HIPAA, GDPR). Unauthorised disclosure, copying or distribution is prohibited. If received in error, please destroy and notify records@maroto.example. Findings reflect a single point-in-time clinical encounter; treatment decisions remain the responsibility of the patient's clinical team.