Testing the Effects of Transfection on Mammalian Cytokine RNA Expression

Austin Mitchell Vecchio

Abstract—Abstract outline

This experiment aims to study the effect of genetic expression on polyplex treated cells against β -actin and INF α .

I. INTRODUCTION

This section introduces the topic and leads the reader on to the main part.

Real Time PCR Generate large quantity of DNA from cDNA templates. Can view amount of DNA in each cycle. Can be seen if immunological response was indicated.

There are multiple controls used throughout this experiment. For controls that compare directly against the immunological transcripts β -actin and RPL13A will be used. INF β will not be used as it will only appear after cycle 40 in the experiment producing undesirable results. The No Template Control will consist of DEPC water.

Outline

polyplex polycationic treatments samples form an immune response The mRNA sampl

Test the effects on

Each student was given a sample as follows

IL6 and IFN α are used as the immunilogical transcripts DEPC water is the no template control

Poly24 Av Poly24 Ts Just Cells No Wash

It is virtually impossible to completely eliminate all genomic DNA from RNA preparations. Therefore, if the assay is not cDNA-specific, it is important to include a minus-reverse transcriptase ("-RT") control in real-time RT-PCR experiments. Typically, the "-RT" control is a mock reverse transcription containing all the RT-PCR reagents, except the reverse transcriptase. The presence of an amplification product in the "-RT" control is indicative of contaminating DNA in the sample.

II. METHODS

A. Purification mRNA

B. Reverse Transcription

C. Qt PCR

iTaq universal SYBR Green supermix $20\mu LR eaction 10 Permuation of Green supermix with four <> ina2: 1 ratio.5 microliters and 10 microliters.$

Each well will get 6μ L of $4ng/\mu$ L of dna as per. The concentrations of cDNA for Ts, Av and Mi were all around

Advisor: Dipl.–Ing. Michelle Ammerman, Lehrstuhl für Nachrichtentechnik, TUM, WS 2050/2051.

TABLE I CDNA CONCENTRATIONS

1

| +RT | -RT | | |
|------------------------|---------------------------|--|--|
| 1μL 10x dsDNase Buffer | 10x 1L 10x dsDNase Buffer | | |
| 1μL dsDNase | 1L dsDNase | | |
| 1μL Template RNA | 1L Template RNA | | |
| 2μL Total RNA (Poly24) | 2L Total RNA (Poly24) | | |

TABLE II CDNA CONCENTRATIONS

| | +RT | -RT |
|---|---|--|
| ĺ | 4μL 5x Reaction Mix | 4μ L 5x Reaction Mix height 2μ L |
| | 2μ L DEPC H20 height 4μ L Nuclease Free Water | 4μL Nuclease Free Wa |

TABLE III CDNA CONCENTRATIONS

| INFA | (Ts, AV, Mi) | 1.32 |
|----------------|--------------|-------|
| INFA | (Ni) | 11.79 |
| β -Actin | (Ts, AV, Mi) | 2.00 |
| β -Actin | (Ni) | 2.9 |

100ng/ μ L. Therefore one set of calculations can be used for all three experiments. The concentrations of Ni, however, were reported ot be 11.2ng/ μ L and thus, had to have a different set of calculations.

Adding supermix, following charts from above and water makex mix.

D. Analysis

Used $\Delta\Delta$ ct method was used in calculating fold inductions.

III. RESULTS

A. Qubit

TABLE IV OUBIT RESULTS

| SD1 | 54.27 |
|-----|---------|
| SD2 | 1057.52 |
| Tk | 8.75 |
| DA | 68.0 |
| AV | 487.0 |
| Tk | 28.0 |

IV. DISCUSSION

For a future experiment, it would be suggested that the polyplex cells are treated in 3 hour increments from 3 hours to 24 hours.

V. CONCLUSION

The mRNA was purified and converted to cDNA. THe resulting concentration was relatively high compared to peers. This eludes that the treatment for these cells of polycationic DNA for 24 hours could result in higher transcription rates.

VI. FIGURES

TABLE V
WELLS FOR IMMUNOLOGICAL RESPONSES

| | 1 | 2 | 3 | 4 | 5 | 6 |
|---|--------|--------|--------|--------|--------|--------|
| Α | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL |
| В | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL |
| С | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL |
| D | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL |
| Е | Ts | Ni | AV | Ts | Ni | AV |
| Е | Ts | Ni | AV | Ts | Ni | AV |
| G | Mi -RT | Mi NTC | | Mi -RT | Mi NTC | |
| Н | | | | | | |

Tasha (Ts) Micah (Mi) Nick (Ni) Austin (AV) -rt controls NTC

Columns 1-3 IL6 Columns 4-6 INF $\alpha Columns7 - 9\beta - actingColumns10 - 12RPL13A$

TABLE VI WELLS FOR CONTROLS

| | 7 | 8 | 9 | 10 | 11 | 12 |
|---|--------|--------|--------|--------|--------|--------|
| A | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL |
| В | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL |
| С | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL |
| D | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL | 4ng/μL |
| Е | Ts | Ni | AV | Ts | Ni | AV |
| Е | Ts | Ni | AV | Ts | Ni | AV |
| G | Mi -RT | Mi NTC | | Mi -RT | Mi NTC | |
| Н | | | | | | |

REFERENCES

- [1] J. Hagenauer, E. Offer, and L. Papke. Iterative decoding of binary block and convolutional codes. *IEEE Trans. Inform. Theory*, vol. 42, no. 2, pp. 429-445, Mar. 1996.
- [2] T. Mayer, H. Jenkac, and J. Hagenauer. Turbo base-station cooperation for intercell interference cancellation. *IEEE Int. Conf. Commun. (ICC)*, Istanbul, Turkey, pp. 356–361, June 2006.
- [3] J. G. Proakis. *Digital Communications*. McGraw-Hill Book Co., New York, USA, 3rd edition, 1995.
- [4] F. R. Kschischang. Giving a talk: Guidelines for the Preparation and Presentation of Technical Seminars. http://www.comm.toronto.edu/frank/ guide/guide.pdf.
- [5] IEEE Transactions LaTeX and Microsoft Word Style Files. http://www.ieee. org/web/publications/authors/transjnl/index.html