

Cyclotomic Polynomials

A Complete Treatment

December 2025

1 Preliminaries

Definition 1.1 (Primitive Root of Unity). Let $n \geq 1$ be a positive integer. A complex number ζ is a *primitive n -th root of unity* if $\zeta^n = 1$ and $\zeta^k \neq 1$ for all $1 \leq k < n$. Equivalently, ζ is primitive if and only if $\text{ord}(\zeta) = n$ in the multiplicative group \mathbb{C}^\times .

The n -th roots of unity are precisely $\zeta_n^k := e^{2\pi i k/n}$ for $k = 0, 1, \dots, n-1$. Among these, ζ_n^k is primitive if and only if $\gcd(k, n) = 1$. The number of primitive n -th roots of unity is therefore $\varphi(n)$, Euler's totient function.

Definition 1.2 (Cyclotomic Polynomial). The *n -th cyclotomic polynomial* is defined as:

$$\Phi_n(x) := \prod_{\substack{1 \leq k \leq n \\ \gcd(k, n) = 1}} (x - \zeta_n^k)$$

where $\zeta_n = e^{2\pi i/n}$. Equivalently, $\Phi_n(x)$ is the minimal polynomial over \mathbb{Q} of any primitive n -th root of unity.

By construction, $\Phi_n(x)$ is a monic polynomial of degree $\varphi(n)$ with roots precisely the primitive n -th roots of unity.

Index convention. For a divisor $d \mid n$, each n -th root of unity ζ_n^k is a primitive d -th root of unity for exactly one d , namely $d = n/\gcd(k, n)$. This partitions the roots of $x^n - 1$ by their primitive order.

2 The Fundamental Factorisation

Theorem 2.1 (Divisor Product Identity). For all positive integers n :

$$x^n - 1 = \prod_{d \mid n} \Phi_d(x)$$

Proof. Both sides are monic polynomials of degree n . The left side has roots $\{\zeta_n^k : 0 \leq k \leq n-1\}$. Each such root ζ_n^k has multiplicative order $d := n/\gcd(k, n)$, which divides n . Thus ζ_n^k is a primitive d -th root of unity and hence a root of $\Phi_d(x)$.

Conversely, every primitive d -th root of unity (for $d \mid n$) satisfies $\zeta^d = 1$, hence $\zeta^n = (\zeta^d)^{n/d} = 1$, so it is an n -th root of unity.

This establishes a bijection between the roots of $x^n - 1$ and the union $\bigcup_{d \mid n} \{\text{roots of } \Phi_d(x)\}$. Since the $\Phi_d(x)$ have pairwise disjoint root sets (a primitive d -th root has order exactly d), and since

$$\sum_{d|n} \deg \Phi_d(x) = \sum_{d|n} \phi(d) = n,$$

the factorisation follows from unique factorisation in $\mathbb{K}[x]$. \square

Corollary 2.2. The cyclotomic polynomial admits the recursive formula:

$$\Phi_n(x) = \frac{x^n - 1}{\prod_{\substack{d|n \\ d < n}} \Phi_d(x)}$$

3 Integrality of Coefficients

Lemma 3.1 (Monic Division in $\mathbb{K}[x]$). Let $f(x), g(x) \in \mathbb{K}[x]$ with $g(x)$ monic. If $g(x) \mid f(x)$ in $\mathbb{K}[x]$, then $g(x) \mid f(x)$ in $\mathbb{K}[x]$; that is, $f(x)/g(x) \in \mathbb{K}[x]$.

Proof. By the division algorithm in $\mathbb{K}[x]$, write $f(x) = g(x)q(x) + r(x)$ where $q(x), r(x) \in \mathbb{K}[x]$ and $\deg r < \deg g$. Since $g \mid f$ in $\mathbb{K}[x]$, we have $r(x) = 0$, so $f(x) = g(x)q(x)$.

We claim $q(x) \in \mathbb{K}[x]$. Suppose not; let $q(x) = \sum_{i=0}^m q_i x^i$ with some $q_i \notin \mathbb{K}$. Write $q_i = a_i/b_i$ in lowest terms. Let p be a prime dividing some denominator b_i , and let j be maximal such that $p \mid b_i$.

Consider the coefficient of $x^{j + \deg g}$ in $f(x) = g(x)q(x)$. Since g is monic of degree $d := \deg g$, this coefficient is:

$$q_j + \sum_{i>j} g_{d-(i-j)} q_i$$

where we set $g_k = 0$ for $k < 0$. By maximality of j , each q_i with $i > j$ has denominator coprime to p , and $g_{d-(i-j)} \in \mathbb{K}$. Thus the sum $\sum_{i>j} g_{d-(i-j)} q_i$ has denominator coprime to p . But q_j has p in its denominator, so the total cannot be an integer, contradicting $f \in \mathbb{K}[x]$. \square

Theorem 3.2 (Integrality). For all $n \geq 1$, $\Phi_n(x) \in \mathbb{K}[x]$.

Proof. We proceed by strong induction on n .

Base case. $\Phi_1(x) = x - 1 \in \mathbb{K}[x]$.

Inductive step. Assume $\Phi_d(x) \in \mathbb{K}[x]$ for all $d < n$. Define:

$$D_n(x) := \prod_{\substack{d|n \\ d < n}} \Phi_d(x)$$

By the inductive hypothesis, each factor lies in $\mathbb{K}[x]$, hence $D\Phi(x) \in \mathbb{K}[x]$. Moreover, $D\Phi(x)$ is monic (being a product of monic polynomials).

By Corollary 2.2:

$$\Phi_n(x) = \frac{x^n - 1}{D_n(x)}$$

Both $x^n - 1 \in \mathbb{K}[x]$ and $D\Phi(x) \in \mathbb{K}[x]$ is monic. By Theorem 2.1, this division is exact in $\mathbb{K}[x]$. Since $\mathbb{K}[x] \cong \mathbb{K}[x]$, polynomial division of $x^n - 1$ by $D\Phi(x)$ in $\mathbb{K}[x]$ yields zero remainder, so the quotient lies in $\mathbb{K}[x]$. By Lemma 3.1, $\Phi\Phi(x) \in \mathbb{K}[x]$. \square

4 The Möbius Inversion Formula

The divisor product identity admits an explicit inversion via the Möbius function.

Definition 4.1 (Möbius Function). The Möbius function $\mu: \mathbb{N} \rightarrow \{-1, 0, 1\}$ is defined by:

$$\mu(n) = \begin{cases} 1 & \text{if } n = 1 \\ (-1)^k & \text{if } n = p_1 p_2 \cdots p_k \text{ for distinct primes } p_i \\ 0 & \text{if } p^2 \mid n \text{ for some prime } p \end{cases}$$

The fundamental property is:

$$\sum_{d \mid n} \mu(d) = \begin{cases} 1 & \text{if } n = 1 \\ 0 & \text{if } n > 1 \end{cases}$$

Theorem 4.2 (Möbius Inversion for Cyclotomic Polynomials). For all $n \geq 1$:

$$\Phi_n(x) = \prod_{d \mid n} (x^{n/d} - 1)^{\mu(d)} = \prod_{d \mid n} (x^d - 1)^{\mu(n/d)}$$

Proof. Taking formal logarithms of the divisor product identity (Theorem 2.1):

$$\log(x^n - 1) = \sum_{d \mid n} \log \Phi_d(x)$$

Define $f(n) := \log \Phi_n(x)$ and $g(n) := \log(x^n - 1)$. The identity states $g(n) = \sum_{d \mid n} f(d)$. By Möbius inversion on the divisor poset:

$$f(n) = \sum_{d \mid n} \mu(n/d) g(d) = \sum_{d \mid n} \mu(n/d) \log(x^d - 1)$$

Exponentiating:

$$\Phi_n(x) = \exp\left(\sum_{d|n} \mu(n/d) \log(x^d - 1)\right) = \prod_{d|n} (x^d - 1)^{\mu(n/d)}$$

The substitution $d \mapsto n/d$ yields the equivalent form $\prod_{d|n} (x^{n/d} - 1)^{\mu(d)}$.

Although the right-hand side is a priori in $\mathbb{R}(x)$, it equals $\Phi_{\mathbb{R}}(x)$ by inversion of Theorem 2.1, hence lies in $\mathbb{R}[x]$ by Theorem 3.2. \square

Remark 4.3 (Validity of the formal argument). The logarithmic manipulation is justified in the ring of formal power series $\mathbb{R}[[x^{\pm 1}]]$. Writing $x^n - 1 = x^n(1 - x^{n/n})$, the expression $\log(1 - x^{n/n}) = -\sum_{k \geq 1} x^{n/n}^k/k$ is a well-defined element of $\mathbb{R}[[x^{\pm 1}]]$. The identity holds in this ring, and the resulting polynomial identity can be verified by observing that both sides are polynomials (by Theorem 3.2) agreeing as formal Laurent series.

5 Explicit Formulae for Special Cases

Proposition 5.1 (Prime Index). For prime p :

$$\Phi_p(x) = \frac{x^p - 1}{x - 1} = x^{p-1} + x^{p-2} + \cdots + x + 1 = \sum_{k=0}^{p-1} x^k$$

Proof. The divisors of p are 1 and p . Thus $x^p - 1 = \Phi_1(x)\Phi_p(x) = (x-1)\Phi_p(x)$. \square

Proposition 5.2 (Prime Power Index). For prime p and $k \geq 1$:

$$\Phi_{p^k}(x) = \Phi_p(x^{p^{k-1}}) = \sum_{j=0}^{p-1} x^{j \cdot p^{k-1}}$$

In particular, $\deg \Phi_{p^k} = \varphi(p^k) = p^{k-1}(p-1)$.

Proof. The divisors of p^k are $1, p, p^2, \dots, p^k$. Among these, $\mu(p^j) \neq 0$ only for $j \in \{0, 1\}$, with $\mu(1) = 1$ and $\mu(p) = -1$. By Theorem 4.2:

$$\Phi_{p^k}(x) = \frac{(x^{p^k} - 1)^{\mu(1)}}{(x^{p^{k-1}} - 1)^{-\mu(p)}} = \frac{x^{p^k} - 1}{x^{p^{k-1}} - 1}$$

The substitution $y = x^{p^{k-1}}$ gives:

$$\Phi_{p^k}(x) = \frac{y^p - 1}{y - 1} = \Phi_p(y) = \Phi_p(x^{p^{k-1}}) \quad \blacksquare$$

Proposition 5.3 (Product of Two Distinct Primes). For distinct primes $p < q$:

$$\Phi_{pq}(x) = \frac{(x^{pq} - 1)(x - 1)}{(x^p - 1)(x^q - 1)}$$

Proof. The divisors of pq are $\{1, p, q, pq\}$ with Möbius values $\mu(1) = 1, \mu(p) = \mu(q) = -1, \mu(pq) = 1$. The formula follows from Theorem 4.2. \square

Proposition 5.4 (The Case $2p$ for Odd Prime p). For odd prime p :

$$\Phi_{2p}(x) = \Phi_p(-x) = x^{p-1} - x^{p-2} + x^{p-3} - \dots - x + 1 = \sum_{k=0}^{p-1} (-1)^{p-1-k} x^k$$

Proof. By Proposition 5.3 with the pair $(2, p)$:

$$\Phi_{2p}(x) = \frac{(x^{2p} - 1)(x - 1)}{(x^2 - 1)(x^p - 1)} = \frac{(x^{2p} - 1)(x - 1)}{(x - 1)(x + 1)(x^p - 1)} = \frac{x^{2p} - 1}{(x + 1)(x^p - 1)}$$

Observe that $x^{2p} - 1 = (x^p)^2 - 1 = (x^p - 1)(x^p + 1)$. Thus:

$$\Phi_{2p}(x) = \frac{x^p + 1}{x + 1}$$

Now $\Phi_p(-x) = ((-x)^p - 1)/((-x) - 1) = (-x^p - 1)/(-x - 1) = (x^p + 1)/(x + 1)$, using that p is odd. \square

6 Reduction Formulae

Theorem 6.1 (Reduction Formulae). Let $n > 1$ and let p be a prime dividing n . Write $n = p^a m$ where $\gcd(p, m) = 1$.

(i) If $a = 1$ (so $n = pm$):

$$\Phi_{pm}(x) = \frac{\Phi_m(x^p)}{\Phi_m(x)}$$

(ii) If $a \geq 2$ (so $p^2 \mid n$):

$$\Phi_n(x) = \Phi_{n/p}(x^p)$$

Proof of (i). The divisors of pm partition as $\{d : d \mid m\} \sqcup \{pd : d \mid m\}$. By Theorem 4.2:

$$\Phi_{pm}(x) = \prod_{d|pm} (x^d - 1)^{\mu(pm/d)}$$

Splitting by whether p divides the divisor:

$$= \prod_{d|m} (x^d - 1)^{\mu(pm/d)} \cdot \prod_{d|m} (x^{pd} - 1)^{\mu(m/d)}$$

Since $\gcd(p, m) = 1$, for $d \mid m$ we have $\mu(pm/d) = \mu(p)\mu(m/d) = -\mu(m/d)$. Thus:

$$\Phi_{pm}(x) = \prod_{d|m} (x^d - 1)^{-\mu(m/d)} \cdot \prod_{d|m} (x^{pd} - 1)^{\mu(m/d)} = \frac{\Phi_m(x^p)}{\Phi_m(x)}$$

where the final identification uses Theorem 4.2 applied to m . \square

Proof of (ii). Write $n = p^a m$ with $a \geq 2$ and $\gcd(p, m) = 1$. By Theorem 4.2:

$$\Phi_n(x) = \prod_{d|n} (x^d - 1)^{\mu(n/d)}$$

For $\mu(n/d) \neq 0$, we require n/d to be squarefree. If $p \nmid d$, then $p^2 \mid (n/d)$, so n/d is not squarefree, hence $\mu(n/d) = 0$. Thus only divisors d with $p \mid d$ contribute.

Writing $d = pe$ for $e \mid (n/p)$, and noting that $n/d = n/(pe) = (n/p)/e$:

$$\Phi_n(x) = \prod_{e|(n/p)} (x^{pe} - 1)^{\mu((n/p)/e)} = \prod_{e|(n/p)} ((x^p)^e - 1)^{\mu((n/p)/e)}$$

The final product is precisely $\Phi_{(n/p)}(x^p)$ by Theorem 4.2 applied to n/p . \square

7 The General Closed Form

Theorem 7.1 (Canonical Form). Let $n = p_1^{a_1} p_2^{a_2} \cdots p_r^{a_r}$ be the prime factorisation of $n > 1$. Define the *radical* $\text{rad}(n) := p_1 p_2 \cdots p_r$. Then:

$$\Phi_n(x) = \Phi_{\text{rad}(n)}(x^{n/\text{rad}(n)})$$

For the squarefree case $n = p_1 p_2 \cdots p_r$:

$$\Phi_n(x) = \prod_{S \subseteq \{1, \dots, r\}} \left(x^{n/\prod_{i \in S} p_i} - 1 \right)^{(-1)^{|S|}}$$

Equivalently:

$$\Phi_n(x) = \prod_{d|n} (x^d - 1)^{\mu(n/d)}$$

Proof. For the first identity, apply Theorem 6.1(ii) repeatedly for each prime p with exponent $a \geq 2$ in n , reducing the exponent by one at each step until all exponents equal one, yielding $\text{rad}(n)$.

For the squarefree expansion, note that $\mu(d) \neq 0$ precisely when d is squarefree. For n squarefree with r prime factors, the divisors $d \mid n$ biject with subsets $S \subseteq \{1, \dots, r\}$ via $d = \prod_{i \in S} p_i$. Then $\mu(d) = (-1)^{|S|}$ and $n/d = \prod_{i \notin S} p_i$. \square

8 Application: Explicit Computation of $\Phi_{12}(x)$

Setting $n = 12 = 2^2 \cdot 3$, we have $\text{rad}(12) = 6$ and $12/\text{rad}(12) = 2$.

Step 1. Compute $\Phi_6(x)$. Since $6 = 2 \cdot 3$, by Proposition 5.3:

$$\Phi_6(x) = \frac{(x^6 - 1)(x - 1)}{(x^2 - 1)(x^3 - 1)}$$

Factoring: $x^6 - 1 = (x^3 - 1)(x^3 + 1)$ and $x^2 - 1 = (x - 1)(x + 1)$. Thus:

$$\Phi_6(x) = \frac{(x^3 - 1)(x^3 + 1)(x - 1)}{(x - 1)(x + 1)(x^3 - 1)} = \frac{x^3 + 1}{x + 1} = x^2 - x + 1$$

Step 2. Apply Theorem 7.1:

$$\Phi_{12}(x) = \Phi_6(x^2) = (x^2)^2 - (x^2) + 1 = x^4 - x^2 + 1$$

Verification. The degree is $\varphi(12) = \varphi(4)\varphi(3) = 2 \cdot 2 = 4$. \checkmark

The roots are $e^{2\pi i k/12}$ for $\gcd(k, 12) = 1$, i.e., $k \in \{1, 5, 7, 11\}$. \checkmark

9 Table of Cyclotomic Polynomials

n	$\Phi_n(x)$	$\deg \Phi_n = \varphi(n)$
1	$x - 1$	1
2	$x + 1$	1

n	$\Phi_n(x)$	$\deg \Phi_n = \varphi(n)$
3	$x^2 + x + 1$	2
4	$x^2 + 1$	2
5	$x^4 + x^3 + x^2 + x + 1$	4
6	$x^2 - x + 1$	2
7	$x^6 + x^5 + x^4 + x^3 + x^2 + x + 1$	6
8	$x^4 + 1$	4
9	$x^6 + x^3 + 1$	6
10	$x^4 - x^3 + x^2 - x + 1$	4
12	$x^4 - x^2 + 1$	4
15	$x^8 - x^7 + x^5 - x^4 + x^3 - x + 1$	8

10 Values at Special Points

Lemma 10.0. For all odd $m > 1$:

$$\Phi_{2m}(x) = \Phi_m(-x)$$

Proof. By Theorem 6.1(i) with $p = 2$ and $\gcd(2, m) = 1$:

$$\Phi_{2m}(x) = \frac{\Phi_m(x^2)}{\Phi_m(x)}$$

We show this equals $\Phi_m(-x)$. By Theorem 4.2:

$$\frac{\Phi_m(x^2)}{\Phi_m(x)} = \prod_{d|m} \frac{(x^{2d} - 1)^{\mu(m/d)}}{(x^d - 1)^{\mu(m/d)}} = \prod_{d|m} \left(\frac{x^{2d} - 1}{x^d - 1} \right)^{\mu(m/d)} = \prod_{d|m} (x^d + 1)^{\mu(m/d)}$$

For the right-hand side, since m is odd, every divisor $d \mid m$ is odd, so $(-x)^d = -x^d$. Thus:

$$\Phi_m(-x) = \prod_{d|m} ((-x)^d - 1)^{\mu(m/d)} = \prod_{d|m} (-x^d - 1)^{\mu(m/d)} = \prod_{d|m} (-(x^d + 1))^{\mu(m/d)}$$

Since $\mu(m/d) \in \{-1, 0, 1\}$, each factor contributes either 1 or -1 accordingly. The total sign is $(-1)^{\sum \{d \mid m\} \mu(m/d)} = (-1)^0 = 1$, since $\sum \{d \mid m\} \mu(m/d) = 0$ for $m > 1$. Hence $\Phi_{2m}(x) = \Phi_m(-x)$. \square

Proposition 10.1.

(i) For $n > 1$:

$$\Phi_n(1) = \begin{cases} p & \text{if } n = p^k \text{ for some prime } p \\ 1 & \text{otherwise} \end{cases}$$

(ii) For $n \geq 1$:

$$\Phi_n(-1) = \begin{cases} -2 & \text{if } n = 1 \\ 0 & \text{if } n = 2 \\ 2 & \text{if } n = 2^k \text{ for } k \geq 2 \\ p & \text{if } n = 2p^k \text{ for an odd prime } p, k \geq 1 \\ 1 & \text{otherwise} \end{cases}$$

Proof of (i). From Theorem 2.1, $x^n - 1 = \prod_{d|n} \Phi_d(x)$. Differentiating and evaluating at $x = 1$:

$$n = \left. \frac{d}{dx}(x^n - 1) \right|_{x=1} = \sum_{d|n} \Phi'_d(1) \prod_{\substack{e|n \\ e \neq d}} \Phi_e(1)$$

Since $\Phi_1(1) = 0$, only the term $d = 1$ survives, yielding:

$$n = \Phi'_1(1) \cdot \prod_{\substack{d|n \\ d > 1}} \Phi_d(1) = 1 \cdot \prod_{\substack{d|n \\ d > 1}} \Phi_d(1)$$

Thus $\prod_{d|n, d > 1} \Phi_d(1) = n$. We prove the closed form by strong induction on n .

For $n = p$ prime, the only divisor greater than 1 is p itself, so $\Phi_p(1) = p$.

For $n = p^k$ with $k \geq 2$, the divisors greater than 1 are p, p^2, \dots, p^k . By induction, $\Phi_{p^j}(1) = p$ for $j < k$. Thus:

$$\prod_{j=1}^k \Phi_{p^j}(1) = p^k \implies p^{k-1} \cdot \Phi_{p^k}(1) = p^k \implies \Phi_{p^k}(1) = p$$

For n with at least two distinct prime factors, write $n = p^a m$ with $\gcd(p, m) = 1$ and $m > 1$. The divisors of n greater than 1 include all divisors of m greater than 1, all divisors of p^a greater than 1, and mixed divisors. By the multiplicative structure:

$$\prod_{\substack{d|n \\ d > 1}} \Phi_d(1) = n = p^a \cdot m$$

The divisors p^j for $1 \leq j \leq a$ contribute p^j (by induction). The divisors of m greater than 1 contribute m (by induction on m). The remaining divisors (those involving both p and primes of m) must therefore contribute 1. Each such $\Phi_d(1)$ is a positive integer: it lies in \mathbb{R} by Theorem 3.2, and $\Phi_d(1) > 0$ because pairing conjugate roots gives $\Phi_d(1) = \prod |1 - \zeta|^2 > 0$. Since their product is 1 and each factor is a positive integer, each equals 1. In particular, $\Phi_n(1) = 1$.

Proof of (ii). Write $n = 2^a m$ with m odd.

Case 1: $a = 0$ (n odd). For $n = 1$, direct computation gives $\Phi_1(-1) = -2$. For odd $n > 1$, Lemma 10.0 gives $\Phi_{2n}(x) = \Phi_n(-x)$, so $\Phi_n(-1) = \Phi_{2n}(1)$. Because $n > 1$ is odd, it has an odd prime divisor p , so $2n$ is divisible by both 2 and p . Hence $2n$ is not a prime power, and $\Phi_{2n}(1) = 1$ by part (i).

Case 2: $a = 1$ ($n = 2m$ with m odd). For $m = 1$, direct computation gives $\Phi_2(-1) = 0$. For $m > 1$, Lemma 10.0 gives $\Phi_{2m}(x) = \Phi_m(-x)$, hence $\Phi_{2m}(-1) = \Phi_m(1)$. By part (i), $\Phi_m(1) = p$ if $m = p^j$ for some odd prime p , and $\Phi_m(1) = 1$ otherwise. This yields $\Phi_{2p^j}(-1) = p$ for odd primes p , and $\Phi_{2m}(-1) = 1$ for other odd $m > 1$.

Case 3: $a \geq 2$ ($4 \mid n$). Apply Theorem 6.1(ii) repeatedly with $p = 2$:

$$\Phi_{2^a m}(x) = \Phi_{2m}(x^{2^{a-1}})$$

Evaluating at $x = -1$:

$$\Phi_{2^a m}(-1) = \Phi_{2m}\left((-1)^{2^{a-1}}\right) = \Phi_{2m}(1)$$

since $2^{a-1} \geq 2$ implies $(-1)^{2^{a-1}} = 1$. If $m = 1$, this gives $\Phi_{2^a}(-1) = \Phi_2(1) = 2$. If $m > 1$, then $2m$ is not a prime power (having both 2 and an odd prime as factors), so $\Phi_{2m}(1) = 1$ by part (i), yielding $\Phi_{2^a m}(-1) = 1$.

11 Generalisations

11.1 Cyclotomic Polynomials over Finite Fields

For a finite field \mathbb{F}_q with $\gcd(q, n) = 1$, the polynomial $\Phi_n(x) \in \mathbb{F}_q[x]$ reduces to $\Phi_n(x) \in \mathbb{F}_q[x]$. This reduced polynomial factors into irreducible factors of equal degree d , where d is the multiplicative order of q modulo n . The number of irreducible factors is $\varphi(n)/d$.

11.2 Generalised Cyclotomic Polynomials

For integers a, b with $\gcd(a, b) = 1$, define:

$$\Phi_n(a, b) := \prod_{\substack{1 \leq k \leq n \\ \gcd(k, n) = 1}} (a - \zeta_n^k b)$$

This is a homogeneous polynomial in a, b of degree $\varphi(n)$ with integer coefficients. The substitution $a = x, b = 1$ recovers $\Phi_{\mathbb{Q}}(x)$.

12 Concluding Remarks

The cyclotomic polynomial $\Phi_{\mathbb{Q}}(x)$ admits a closed form via Möbius inversion:

$$\Phi_n(x) = \prod_{d|n} (x^d - 1)^{\mu(n/d)}$$

The integrality $\Phi_{\mathbb{Q}}(x) \in \mathbb{Z}[x]$ follows from strong induction using monic polynomial division. The derivation rests on three principles:

1. **Divisor partition.** The n -th roots of unity partition by primitive order into disjoint sets indexed by $d \mid n$.
2. **Möbius inversion.** The divisor sum $g(n) = \sum_{d \mid n} f(d)$ *inverts to* $f(n) = \sum_{d \mid n} \mu(n/d) g(d)$.
3. **Monic divisibility.** Division by monic polynomials preserves integrality in $\mathbb{Z}[x]$.

The reduction $\Phi_{\mathbb{Q}}(x) = \Phi_{\{\text{rad}(n)\}}(x^{n/\text{rad}(n)})$ shows that computation of $\Phi_{\mathbb{Q}}(x)$ reduces to the squarefree case, where the Möbius formula involves 2^r terms for r distinct prime factors.

13 Appendix: Algorithmic Summary

To compute $\Phi_{\mathbb{K}}(x)$:

1. Compute the prime factorisation $n = p_1^{a_1} \cdots p_r^{a_r}$.
2. Compute $m := \text{rad}(n) = p_1 \cdots p_r$ and $e := n/m = p_1^{a_1-1} \cdots p_r^{a_r-1}$.
3. Compute $\Phi_{\mathbb{K}}(x)$ via:

$$\Phi_m(x) = \prod_{S \subseteq \{1, \dots, r\}} \left(x^{m/\prod_{i \in S} p_i} - 1 \right)^{(-1)^{|S|}}$$

4. Return $\Phi_{\mathbb{K}}(x) = \Phi_{\mathbb{K}}(x^e)$.

For implementation, the product in step 3 is computed iteratively: initialise $P(x) := 1$ (empty product), then for each subset S , multiply or divide by $x^{m/d_S} - 1$ according to the parity of $|S|$.