

Notes on fixed-point procedure

Contents

1	Notations and assumptions	1
2	Estimates	2
2.1	Useful elementary lemmas	2
2.2	Boundedness estimates	4
2.2.1	Electronic density n_e	4
2.2.2	Ion density n_i	4
2.3	Continuity estimates	9
2.3.1	Electron density n_e	9
2.3.2	Ion density n_i	10
3	Results	11

1 Notations and assumptions

We suppose that

1. $f_{e,b}$ is nonnegative.
2. $f_{e,b}(v) = f_{e,b}(-v)$ for all $v \in \mathbb{R}$.
3. $f_{e,b}$ is compactly supported, i.e. there exists $v^* \geq 0$ such that $f_{e,b}(v) = 0$ for all $|v| \geq v^*$.
4. There exists $\bar{c} \leq 0$ such that $f_{e,b}(v) \leq \bar{c}$ for all $v \in \mathbb{R}$.
5. There exists $\underline{c} > 0$ and $0 \leq \underline{v} < \bar{v}$ such that $f_{e,b}(v) \geq \underline{c}$ for all $v \in [-\bar{v}, -\underline{v}]$.
6. The function φ satisfies $\varphi(0) = \varphi'(0) = 0$.
7. The function φ is strongly concave, i.e. there exists $\alpha > 0$ such that $\varphi((1-\tau)x + \tau y) \geq (1-\tau)\varphi(x) + \tau\varphi(y) + \frac{\alpha}{2}(1-\tau)\tau|x-y|^2$ for all $(x, y, \tau) \in [0, 1]^3$.
8. There exists $\beta > \alpha$ such that φ is $(-\beta)$ -convex, i.e. $\varphi((1-\tau)x + \tau y) \leq (1-\tau)\varphi(x) + \tau\varphi(y) + \frac{\beta}{2}(1-\tau)\tau|x-y|^2$ for all $(x, y, \tau) \in [0, 1]^3$. Notice that in this case, $\varphi(x) \geq -\beta\frac{x^2}{2}$, and in particular, $\varphi(1) \geq -\beta$.

For the reader: notations \underline{v}, \bar{v} indicates a domain on which $f_{e,b}$ is *lower bounded*, while v_*, v^* denote a domain on which $f_{e,b}$ is *upper bounded by 0* (with a global bound by \bar{c} anyway).

We use the following notations: the Liapunov functions of the electrons and the ions are

$$\mathcal{L}_e(x, v) := \frac{v^2}{2} - \frac{1}{\mu}\varphi(x). \quad (1.1)$$

$$\mathcal{L}_i(x, v) := \frac{v^2}{2} + \varphi(x). \quad (1.2)$$

For a given point $(x, v) \in [0, 1] \times \mathbb{R}^-$, we denote by (x_b, v_b) the intersection of the boundary $\{x \geq 0, v = 0\} \cup \{x = 0, v \leq 0\}$ with the ion characteristic issued from (x, v) . The values are given by

$$\begin{pmatrix} x_b(x, v) \\ v_b(x, v) \end{pmatrix} := \begin{pmatrix} \varphi^{-1} \left(\min \left(0, \frac{v^2}{2} + \varphi(x) \right) \right) \\ -\sqrt{\max(0, v^2 + 2\varphi(x))} \end{pmatrix} \quad (1.3)$$

The satisfaction of the non-emitting boundary condition in $(x = 1, v \in \mathbb{R}_-)$ may interfere with the value on the line $(x = 0, v \in \mathbb{R})$. **In the following notes, we consider that the non-emitting boundary condition has the precedence**, i.e. $f_e(x, v) = 0$ if $\mathcal{L}_e(x, v) \geq \mathcal{L}_e(1, 0)$, regardless of the prescribed value of $f_{e,b}$ along this line.

2 Estimates

2.1 Useful elementary lemmas

Lemma 2.1. *Let $\alpha > 0$, $\bar{x} \geq 0$ and $\bar{y} \geq 0$. Then*

$$\mathcal{I} := \int_{x=0}^{\bar{x}} \int_{y=0}^{\bar{y}} \frac{1}{(x^2 + \alpha y^2)^{1/2}} dy dx \leq 2 \frac{\sqrt{2\bar{x}\bar{y}}}{\alpha^{1/4}}.$$

Demonstration By the change of variable $z = \sqrt{\alpha}y$, with $\bar{z} := \sqrt{\alpha}\bar{y}$, we have

$$\mathcal{I} = \frac{1}{\sqrt{\alpha}} \int_{x=0}^{\bar{x}} \int_{z=0}^{\bar{z}} \frac{1}{(x^2 + z^2)^{1/2}} dz dx.$$

Notice that $x + z \leq \sqrt{2}(x^2 + y^2)^{1/2}$. Then,

$$\mathcal{I} \leq \frac{1}{\sqrt{\alpha}} \int_{x=0}^{\bar{x}} \int_{z=0}^{\bar{z}} \frac{\sqrt{2}}{x+z} dz dx = \sqrt{\frac{2}{\alpha}} \int_{x=0}^{\bar{x}} \ln \left(\frac{x+\bar{z}}{x} \right) dx = \sqrt{\frac{2}{\alpha}} \int_{x=0}^{\bar{x}} \ln \left(1 + \frac{\bar{z}}{x} \right) dx.$$

Using $\ln(1+a) \leq \sqrt{a}$, we get

$$\mathcal{I} \leq \sqrt{\frac{2}{\alpha}} \int_{x=0}^{\bar{x}} \sqrt{\frac{\bar{z}}{x}} dx = \sqrt{\frac{2}{\alpha}} 2\sqrt{\bar{x}\bar{z}} = 2 \frac{\sqrt{2\bar{x}\bar{y}}}{\alpha^{1/4}}.$$

□

Remark 2.1 (Exact value). *Let $\bar{r} := \sqrt{\bar{x}^2 + \bar{z}^2}$. Then*

$$\mathcal{I} = \bar{x} \ln \left(\frac{1 + \bar{z}/\bar{r}}{\bar{x}/\bar{r}} \right) + \bar{z} \ln \left(\frac{1 + \bar{x}/\bar{r}}{\bar{z}/\bar{r}} \right).$$

Lemma 2.2. *If $a \geq 0$ and $b \geq 0$, then $|a - b| \leq \sqrt{|a^2 - b^2|}$.*

Demonstration If $a \geq b$, then $|a - b| = \sqrt{(a - b)(a - b)} \leq \sqrt{(a - b)(a + b)} = \sqrt{a^2 - b^2}$, else $|a - b| = |b - a|$. □

Lemma 2.3. *Let $\alpha > 0$, $L \in \mathbb{R}$ and $0 \leq a \leq 1$. Suppose that $L + \alpha a^2 \geq 0$, and $a > 0$ if $L = 0$. Then*

$$\mathcal{I} := \int_{z=a}^1 \frac{1}{(L + \alpha z^2)^{1/2}} dz \leq \min \left(|L|^{-1/2}, \frac{2}{\sqrt{\alpha a}} \right).$$

Demonstration If $L > 0$, we have

$$\mathcal{I} = \frac{1}{\sqrt{\alpha}} \int_{z=a}^1 \frac{1}{\left(1 + \left(\sqrt{\frac{\alpha}{L}} z\right)^2\right)^{1/2}} \frac{\sqrt{\alpha} dz}{\sqrt{L}} = \frac{1}{\alpha} \int_{w=a\sqrt{\frac{\alpha}{L}}}^{\sqrt{\frac{\alpha}{L}}} \frac{1}{(1+w^2)^{1/2}} dw = \frac{1}{\sqrt{\alpha}} \left(\sinh^{-1} \left(\sqrt{\frac{\alpha}{L}} \right) - \sinh^{-1} \left(a \sqrt{\frac{\alpha}{L}} \right) \right).$$

Using the positivity of $\sinh^{-1} \left(a \sqrt{\frac{\alpha}{L}} \right)$, and the coarse estimate $\sinh^{-1}(x) \leq x$, we get $\mathcal{I} \leq L^{-1/2}$. Moreover, using that $\sinh^{-1}(x) = \log(x + \sqrt{x^2 + 1})$, we get

$$\mathcal{I} = \frac{1}{\sqrt{\alpha}} \log \left(\frac{\sqrt{\frac{\alpha}{L}} + \sqrt{\frac{\alpha}{L} + 1}}{a \sqrt{\frac{\alpha}{L}} + \sqrt{a^2 \frac{\alpha}{L} + 1}} \right) = \frac{1}{\sqrt{\alpha}} \log \left(\frac{1 + \sqrt{1 + \frac{L}{\alpha}}}{a + \sqrt{a^2 + \frac{L}{\alpha}}} \right) \leq \frac{1}{\sqrt{\alpha}} \log \left(\frac{2 + \sqrt{\frac{L}{\alpha}}}{2a} \right) \leq \frac{1 + \left(\frac{L}{\alpha}\right)^{1/4}}{\sqrt{\alpha a}}.$$

I shamelessly used $\frac{1}{2} \leq 1$. Then $\mathcal{I} \leq \min \left(L^{-1/2}, \frac{1 + \left(\frac{L}{\alpha}\right)^{1/4}}{\sqrt{\alpha a}} \right)$ on $L > 0$. But whenever $L \geq \alpha$, the min is attained in its first argument: indeed, $L^{-1/2} \leq \frac{1}{\sqrt{\alpha a}} \leq \frac{1 + \left(\frac{L}{\alpha}\right)^{1/4}}{\sqrt{\alpha a}}$. Then we obtained $\mathcal{I} \leq \min(L^{-1/2}, \frac{2}{\sqrt{\alpha a}})$ on $L > 0$. If $L = 0$, we have

$$\int_{z=a}^1 \frac{1}{\sqrt{\alpha} z} dz = \frac{\log(1/a)}{\sqrt{\alpha}} \frac{2}{\sqrt{\alpha a}}.$$

Finally, if $L < 0$, the integral becomes

$$\mathcal{I} = \frac{1}{\sqrt{\alpha}} \int_a^1 \frac{1}{\left(\left(\sqrt{\frac{\alpha}{|L|}} z \right)^2 - 1 \right)^{1/2}} \frac{\sqrt{\alpha} dz}{\sqrt{|L|}} = \frac{1}{\alpha} \int_{a\sqrt{\frac{\alpha}{|L|}}}^{\sqrt{\frac{\alpha}{|L|}}} \frac{1}{(w^2 - 1)^{1/2}} dw = \frac{\cosh^{-1} \left(\sqrt{\frac{\alpha}{|L|}} \right) - \cosh^{-1} \left(a \sqrt{\frac{\alpha}{|L|}} \right)}{\sqrt{\alpha}}.$$

The expression is well-defined, since $L + \alpha a^2 \geq 0$ implies $a \sqrt{\frac{\alpha}{-L}} \geq 1$ when $-L > 0$. Since $\cosh^{-1}(x) \leq x$, we obtain the coarse estimate $\mathcal{I} \leq |L|^{-1/2}$. Using that $\cosh^{-1}(x) = \log(x + \sqrt{x^2 - 1})$, we get

$$\mathcal{I} = \frac{1}{\sqrt{\alpha}} \log \left(\frac{1 + \sqrt{1 - \frac{|L|}{\alpha}}}{a + \sqrt{a^2 - \frac{|L|}{\alpha}}} \right) \leq \frac{1}{\sqrt{\alpha}} \log \left(\frac{2}{a} \right) \leq \sqrt{\frac{2}{\alpha a}} \leq \frac{2}{\sqrt{\alpha a}}.$$

□

Lemma 2.4 (MAGIC Change of variable). *Let $(x(\tau), v(\tau))$ be an ion characteristic issued from $(x, w) \in [0, 1] \times \mathbb{R}_-$, and define $\varepsilon := \mathcal{L}_e(x(\tau), v(\tau))$ and $z := x(\tau)$. Then*

$$d\tau \wedge dw = \frac{dz \wedge d\varepsilon}{2 \left(\varepsilon + \frac{1}{\mu} \varphi(z) \right)^{1/2} \left(\varepsilon - \varphi(x) + \left(1 + \frac{1}{\mu} \right) \varphi(z) \right)^{1/2}}.$$

Demonstration We have $\varepsilon = \frac{(v(\tau))^2}{2} - \frac{1}{\mu} \varphi(z)$, so that

$$d\varepsilon = v(\tau) \dot{v}(\tau) d\tau - \frac{1}{\mu} \varphi'(z) dz = -v(\tau) \varphi'(z) d\tau - \frac{1}{\mu} \varphi'(z) dz \implies d\tau = -\frac{1}{v(\tau) \varphi'(z)} d\varepsilon - \frac{1}{\mu v(\tau)} dz.$$

Moreover, $\mathcal{L}_i(x(\tau), v(\tau)) = \frac{(v(\tau))^2}{2} + \varphi(z)$, so $\left(1 + \frac{1}{\mu} \right) \varphi(z) = \mathcal{L}_i(x(\tau), v(\tau)) - \mathcal{L}_e(x(\tau), v(\tau)) = \mathcal{L}_i(x, w) - \varepsilon$ and

$$w = -\sqrt{2} \left(\varepsilon - \varphi(x) + \left(1 + \frac{1}{\mu} \right) \varphi(z) \right)^{1/2} \implies = \frac{d\varepsilon + \left(1 + \frac{1}{\mu} \right) \varphi'(z) dz}{w}.$$

Then, using $v(\tau) = -\sqrt{2} \left(\varepsilon + \frac{1}{\mu} \varphi(z) \right)^{1/2}$, we get

$$d\tau \wedge dw = -\frac{1 + \frac{1}{\mu}}{w v(\tau)} d\varepsilon \wedge dz - \frac{\frac{1}{\mu}}{v(\tau) w} dz \wedge d\varepsilon = \frac{dz \wedge d\varepsilon}{v(\tau)(-w)} = \frac{dz \wedge d\varepsilon}{2 \left(\varepsilon + \frac{1}{\mu} \varphi(z) \right)^{1/2} \left(\varepsilon - \varphi(x) + \left(1 + \frac{1}{\mu} \right) \varphi(z) \right)^{1/2}}.$$

□

Remark 2.2. Notice that $\mathcal{L}_i(x(\tau), v(\tau)) = \varepsilon + \left(1 + \frac{1}{\mu}\right) \varphi(z)$. Moreover, we have $-\frac{1}{\mu} \varphi(z) = \mathcal{L}_e(z, 0)$, and $\varphi(x) = \mathcal{L}_i(z, 0)$. Then the change of variable rewrites

$$d\tau \wedge dw = \frac{dz \wedge d\varepsilon}{2(\mathcal{L}_e(x(\tau), v(\tau)) - \mathcal{L}_e(z, 0))^{1/2} (\mathcal{L}_i(x(\tau), v(\tau)) - \mathcal{L}_i(x, 0))^{1/2}},$$

and is well-defined for (τ, w) such that both square roots are real.

2.2 Boundedness estimates

2.2.1 Electronic density n_e

Lemma 2.5. The electron density n_e is bounded independently of φ . More precisely,

$$n_e(x) \leq 2\bar{c}v^* \quad \forall x \in [0, 1].$$

Demonstration Let $v \in \mathbb{R}$ such that $|v| > v^*$, with $f_{e,b}(\mathbb{R} \setminus [-v^*, v^*]) = 0$. Then

$$\mathcal{L}_e(x, v) = \frac{v^2}{2} - \frac{1}{\mu} \varphi(x) \geq \frac{v^2}{2} = \mathcal{L}_e(v, 0) \implies f_e(x, v) = 0.$$

Then we may write for all $x \in [0, 1]$ that

$$n_e(x) = \int_{v=-v^*}^{v^*} f_e(x, v) dv \leq 2\bar{c}v^*.$$

□

2.2.2 Ion density n_i

The estimates will rely on two particular cases, the we treat independently as lemmas. For a given x , we define $g_x : [0, x] \mapsto \mathbb{R}^+$ by

$$\mathcal{L}_i(x, -g_x(y)) = \mathcal{L}_i(y, 0), \quad \text{i.e.} \quad g_x(y) = (2(\varphi(y) - \varphi(x)))^{1/2}.$$

For the reader: draw y to the left of x . Then $-g_x(y)$ is the intersection of the ion characteristic ending on $(y, 0)$ with the vertical line going through x .

Lemma 2.6. Let $0 \leq y < x \leq 1$. We have

$$\mathcal{I} := \int_{v=-g_x(y)}^0 \int_{z=x_b(x,v)}^x \frac{1}{(v^2 - g_x^2(z))^{1/2}} dz dv \leq 2 \left(\frac{\beta}{\alpha}\right)^{1/4} (x - y).$$

Demonstration Let us first use Fubini's theorem to switch the order of integration. The lower bound $x_b(x, v) \geq z$ becomes an upper bound $v \leq -g_x(z)$, and we have

$$\mathcal{I} = \int_{z=y}^x \int_{v=-g_x(y)}^{-g_x(z)} \frac{1}{(v^2 - g_x^2(z))^{1/2}} dv dz = \int_{z=y}^x \int_{v=g_x(z)}^{g_x(y)} \frac{1}{(v^2 - g_x^2(z))^{1/2}} dv dz.$$

With the change of variable $w = v - g_x(z)$, and using $\frac{d}{dw} [\sinh^{-1}(\sqrt{\frac{w}{a}})] = (w^2 + 2aw)^{-1/2}$, we get

$$\mathcal{I} = \int_{z=y}^x \int_{w=0}^{g_x(y)-g_x(z)} \frac{1}{(w^2 + 2wg_x(z))^{1/2}} dw dz = \int_{z=y}^x \sinh^{-1} \left(\sqrt{\frac{g_x(y)-g_x(z)}{g_x(z)}} \right) dz.$$

Using the elementary estimate $\sinh^{-1}(a) \leq pa^{1/p}$ for all $p \in \mathbb{N}^*$, and the coarse estimate $\sqrt{\frac{a-b}{b}} \leq \sqrt{\frac{a}{b}}$, we get

$$\mathcal{I} \leq \int_{z=y}^x p \left(\frac{g_x(y)}{g_x(z)} \right)^{1/2p} dz = \int_{z=y}^x p \left(\frac{\varphi(y) - \varphi(x)}{\varphi(z) - \varphi(x)} \right)^{1/4p} dz \quad \forall p \in \mathbb{N}^*.$$

The assumption of strong concavity yields $\varphi(z) - \varphi(x) \geq -\varphi'(z)(x-z) + \frac{\alpha}{2}|x-z|^2 \geq \frac{\alpha}{2}(x-z)^2$, so that

$$\mathcal{I} \leq \left(\frac{2}{\alpha}\right)^{1/4p} (\varphi(y) - \varphi(x))^{1/4p} \int_{z=y}^x p \frac{1}{(x-z)^{1/2p}} dz = \left(\frac{2}{\alpha}\right)^{1/4p} (\varphi(y) - \varphi(x))^{1/4p} \frac{p^2}{p-1/2} (x-y)^{1-1/2p}.$$

By the strong $(-\beta)$ -convexity, we may write $\varphi(y) - \varphi(x) \leq \beta \frac{(x-y)^2}{2}$, so that

$$\mathcal{I} \leq \left(\frac{\beta}{\alpha}\right)^{1/4p} \frac{p^2}{p-1/2} (x-y) \quad \forall p \in \mathbb{N}^*.$$

Using the coarse estimates $\sinh^{-1}(a) \leq a$ and $\sqrt{\frac{a-b}{b}} \leq \sqrt{\frac{a}{b}}$, we get

$$\mathcal{I} \leq \int_{z=y}^x \sqrt{\frac{g_x(y)}{g_x(z)}} dz = \int_{z=y}^x \left(\frac{\varphi(y) - \varphi(x)}{\varphi(z) - \varphi(x)} \right)^{1/4} dz.$$

The assumption of strong concavity yields $\varphi(z) - \varphi(x) \geq -\varphi'(z)(x-z) + \frac{\alpha}{2}|x-z|^2 \geq \frac{\alpha}{2}(x-z)^2$, so that

$$\mathcal{I} \leq \left(\frac{2}{\alpha}\right)^{1/4} (\varphi(y) - \varphi(x))^{1/4} \int_{z=y}^x \frac{1}{(x-z)^{1/2}} dz = 2 \left(\frac{2}{\alpha}\right)^{1/4} (\varphi(y) - \varphi(x))^{1/4} \sqrt{x-y}.$$

By the strong $(-\beta)$ -convexity, we may write $\varphi(y) - \varphi(x) \leq \beta \frac{(x-y)^2}{2}$, so that

$$\mathcal{I} \leq 2 \left(\frac{\beta}{\alpha}\right)^{1/4} (x-y).$$

□

Lemma 2.7. *Let $0 \leq y \leq x \leq 1$, and $-v_0 < -g_x(y)$. We have*

$$\mathcal{I} := \int_{v=-v_0}^{-g_x(y)} \int_{z=y}^1 \frac{1}{(v^2 + 2(\varphi(x) - \varphi(z)))^{1/2}} dz dv \leq \frac{2\sqrt{2}}{\alpha^{1/4}} \left(v_0^2 + \beta(x-y)^2\right)^{1/4}.$$

Demonstration We first shift the v -integration from the vertical line $z = x$ to $z = y$. Let $w = w(v)$ be such that

$$\mathcal{L}_i(y, w(v)) = \mathcal{L}_i(x, v), \quad \text{i.e.} \quad w(v) = -\left(v^2 + 2(\varphi(x) - \varphi(y))\right), \quad \text{and} \quad dv = \frac{-w}{(w^2 + 2(\varphi(y) - \varphi(x)))^{1/2}} dw.$$

Then, defining $w_0 := (v_0^2 + 2(\varphi(x) - \varphi(y)))^{1/2}$, and noticing that $w(-g_x(y)) = 0$, we get

$$\mathcal{I} = \int_{w=-w_0}^0 \int_{z=y}^1 \frac{1}{(w^2 + 2(\varphi(y) - \varphi(z)))^{1/2}} dz \frac{-w}{(w^2 + 2(\varphi(y) - \varphi(x)))^{1/2}} dw.$$

Since $y \leq x$, we have $\varphi(y) \geq \varphi(x)$, and $\frac{-w}{(w^2 + 2(\varphi(y) - \varphi(x)))^{1/2}} \leq \frac{-w}{|w|} = 1$. By the strong concavity assumption, we have $\varphi(y) - \varphi(z) \geq -\varphi'(y)(z-y) + \frac{\alpha}{2}|z-y|^2 \geq \frac{\alpha}{2}(z-y)^2$, so that

$$\mathcal{I} \leq \int_{w=-w_0}^0 \int_{z=y}^1 \frac{1}{(w^2 + \alpha(z-y)^2)^{1/2}} dz dw = \int_{w=0}^{w_0} \int_{z=0}^{1-y} \frac{1}{(w^2 + \alpha z^2)^{1/2}} dz dw.$$

Using lemma (2.1), and the $(-\beta)$ -convexity, we conclude that

$$\mathcal{I} \leq 2 \frac{\sqrt{2w_0(1-y)}}{\alpha^{1/4}} = \frac{2\sqrt{2(1-y)}}{\alpha^{1/4}} (v_0^2 + 2(\varphi(x) - \varphi(y)))^{1/4} \leq \frac{2\sqrt{2}}{\alpha^{1/4}} (v_0^2 + \beta(x-y)^2)^{1/4}.$$

□

Proposition 2.1. *The density n_i is bounded. More precisely,*

$$n_i(x) \leq 2\nu \left(2 \left(\frac{\beta}{\alpha}\right)^{1/4} + 2\sqrt{2} \left(\frac{\beta}{\alpha}\right)^{1/4} + \frac{2\sqrt{2}}{\alpha^{1/4}} \sqrt{v^*} \right) \quad \forall x \in [0, 1].$$




Figure 1: Decomposition of the integral defining n_i .

The phase space is divided by the critical characteristic (in solid black). Whenever $v \leq -g_x(0)$, the characteristics (dotted green lines) are reaching the boundary with $x_b(x, v) = 0$.

Demonstration We use the symmetry of f_i to write

$$n_i(x) = \int_{v=-\infty}^{\infty} f_i(x, v) dv = 2 \int_{v=-\infty}^0 f_i(x_b(x, v), v_b(x, v)) dv = 2\nu \int_{v=-\infty}^0 \int_{t=-\infty}^0 f_e(x(t), v(t)) dt dv,$$

where $(x(t), v(t))_{t \leq 0}$ is the ion characteristic reaching $(x_b(x, v), v_b(x, v))$ at $t = 0$. Notice that the lower bounds are artificial, since the characteristic enters the support of f_e in finite time: we may use $v \geq -v^*$, and consider only times t for which $x(t) \in [0, 1]$.

We first reparametrize $(x(t), v(t))$ using the space variable. Define $z = x(t) \in [x_b, 1]$, and observe that

$$dz = \dot{x}(t) dt = v(t) dt, \quad \text{with} \quad \mathcal{L}_i(z, v(t)) = \mathcal{L}_i(x, v) \iff v(t) = -(v^2 + 2(\varphi(x) - \varphi(z)))^{1/2}.$$

Then, the density rewrites

$$n_i(x) = 2\nu \int_{v=-v^*}^0 \int_{z=x_b(x, v)}^1 \frac{f_e\left(z, -(v^2 + 2(\varphi(x) - \varphi(z)))^{1/2}\right)}{(v^2 + 2(\varphi(x) - \varphi(z)))^{1/2}} dz dv.$$

Let us show that n_i is bounded. We use the coarse estimate $f_e \leq \bar{c}$, and decompose the integral in three:

$$n_i(x) \leq 2\nu \bar{c} \left[\underbrace{\int_{v=-g_x(0)}^0 \int_{z=x_b(x, v)}^x}_{\mathcal{I}_1} + \underbrace{\int_{v=-g_x(0)}^0 \int_{z=x}^1}_{\mathcal{I}_2} + \underbrace{\int_{v=-v^*}^{-g_x(0)} \int_{z=x_b(x, v)}^1}_{\mathcal{I}_3} \right] \frac{1}{(v^2 + 2(\varphi(x) - \varphi(z)))^{1/2}} dz dv.$$

The corresponding domains are represented figure (3).

Notice that whenever $z \leq x$, we have $0 \geq 2(\varphi(x) - \varphi(z)) = -(2(\varphi(z) - \varphi(x)))^{2/2} = -g_x^2(z)$. Then, the integral \mathcal{I}_1 may be bounded using lemma (2.6) with $y = 0$:

$$\mathcal{I}_1 = \int_{v=-g_x(0)}^0 \int_{z=x_b(x, v)}^x \frac{1}{(v^2 - g_x^2(z))^{1/2}} dz dv \leq 2 \left(\frac{\beta}{\alpha} \right)^{1/4} x \leq 2 \left(\frac{\beta}{\alpha} \right)^{1/4}.$$

We use lemma (2.7) to bound \mathcal{I}_2 and \mathcal{I}_3 . In the first case, we take $y = x$ and $v_0 = g_x(0)$, and notice that $-g_x(x) = 0$. In the second case, we take $v_0 = v^*$ and $y = 0$, and notice that on $v \leq -g_x(0)$, we have $x_b(x, v) = 0$ (the velocity is low enough so that the characteristic ends on $x_b = 0$). Finally, we use that $g_x^2(0) = -2\varphi(1) \leq \beta$ to write

$$\mathcal{I}_2 \leq \frac{2\sqrt{2}}{\alpha^{1/4}} (g_x^2(0))^{1/4} \leq 2\sqrt{2} \left(\frac{\beta}{\alpha} \right)^{1/4}, \quad \text{and} \quad \mathcal{I}_3 \leq \frac{2\sqrt{2}}{\alpha^{1/4}} ((v^*)^2 + 2\varphi(x))^{1/4} \leq \frac{2\sqrt{2}}{\alpha^{1/4}} \sqrt{v^*}.$$

□

Lemma 2.8. *The density n_i is uniformly bounded away from 0. More precisely,*

$$n_i(x) \geq \frac{\sqrt{2\nu\mathfrak{c}(\bar{v}^2 - \underline{v}^2)}}{\bar{v} \left(\frac{\bar{v}^2}{2} + \beta \right)^{1/2}} \frac{\underline{v}}{\sqrt{\beta \left(1 + \frac{1}{\mu} \right)}} \quad \forall x \in [0, 1].$$

Demonstration By assumption, there exists a constant $\mathfrak{c} > 0$ such that $f_{e,b}(v) \geq \mathfrak{c} \mathbf{1}_{\{v \leq |v| \leq \bar{v}\}}$, which implies

$$f_e(x, v) \geq \mathfrak{c} \mathbf{1}_{\{\mathcal{L}_e(0, \underline{v}) \leq \mathcal{L}_e(x, v) \leq \mathcal{L}_e(0, \bar{v})\}}.$$

Notice that we may always take $\underline{v} > 0$ as long as $\bar{v} > 0$. Let us denote $\underline{\varepsilon} := \mathcal{L}_e(0, \underline{v})$ and $\bar{\varepsilon} := \mathcal{L}_e(0, \bar{v})$. For any $x \in [0, 1]$, we have

$$n_i(x) = 2\nu \int_{w \in \mathbb{R}_-} \int_{\tau=-\infty}^0 f_e(x(\tau), v(\tau)) d\tau dw, \quad \text{with} \quad \mathcal{L}_i(x(\tau), v(\tau)) = \mathcal{L}_i(x, w).$$

Notice that for $w \in [-g_x(-\bar{v}), -g_x(0)]$, the characteristic issued from (x, w) enters the domain $\{\underline{\varepsilon} \leq \mathcal{L}_e \leq \bar{\varepsilon}\}$. We will restrict the domain of integration to

$$\mathcal{D} := \{(w, \tau) \in \mathbb{R}_-^2 \mid \underline{\varepsilon} \leq \mathcal{L}_e(x(\tau), v(\tau)) \leq \bar{\varepsilon} \text{ and } 0 \leq x(\tau) \leq a\},$$

where (a, v_a) is the unique point such that $\mathcal{L}_e(a, v_a) = \mathcal{L}_e(0, \underline{v})$ and $\mathcal{L}_i(a, v_a) = 0$ (note that a depends on β). This (somehow coarse) estimate will give us trivial bounds when applying the change of variable of lemma (2.4), namely $z = x(\tau)$ and $\varepsilon = \mathcal{L}_e(x(\tau), v(\tau))$. Denoting by $[\underline{\tau}, \bar{\tau}] \subset]-\infty, 0]$ the time interval where $(x(\tau), v(\tau)) \in \mathcal{D}$, we have

$$n_i(x) \geq \int_{w=-g_x(-\bar{v})}^{-g_x(0)} \int_{\tau=\underline{\tau}}^{\bar{\tau}} 2\nu\mathfrak{c} d\tau dw = \int_{\varepsilon=\underline{\varepsilon}}^{\bar{\varepsilon}} \int_{z=0}^a \frac{2\nu\mathfrak{c}}{2 \left(\varepsilon + \frac{1}{\mu} \varphi(z) \right)^{1/2} \left(\varepsilon - \varphi(x) + \left(1 + \frac{1}{\mu} \right) \varphi(z) \right)^{1/2}} dz d\varepsilon.$$

Using remark (2.2), we are assured that the square roots are well-defined, since

$$\varepsilon \geq \underline{\varepsilon} = \mathcal{L}_e(0, \underline{v}) = \mathcal{L}_e(a, v_a) \geq \mathcal{L}_e(a, 0) \geq \mathcal{L}_e(z, 0) = -\frac{1}{\mu} \varphi(z) \quad \text{and} \quad \mathcal{L}_i(x(\tau), v(\tau)) = \mathcal{L}_i(x, w) \geq \mathcal{L}_i(x, 0).$$

Using that $\varepsilon + \frac{1}{\mu} \varphi(z) \leq \bar{\varepsilon} + 0$ and $\varepsilon - \varphi(x) + \left(1 + \frac{1}{\mu} \right) \varphi(z) \leq \bar{\varepsilon} + \beta + 0$, we obtain

$$n_i(x) \geq 2\nu \int_{\varepsilon=\underline{\varepsilon}}^{\bar{\varepsilon}} \int_{z=0}^a \frac{\mathfrak{c}}{2\bar{\varepsilon}^{1/2} (\bar{\varepsilon} + \beta)^{1/2}} dz d\varepsilon = \frac{\nu a \mathfrak{c} (\bar{\varepsilon} - \underline{\varepsilon})}{\bar{\varepsilon}^{1/2} (\bar{\varepsilon} + \beta)^{1/2}}.$$

Let us make explicit the dependance over β by estimating a using the $(-\beta)$ -convexity assumption: we have

$$\varphi(x) \geq -\beta \frac{x^2}{2} \quad \Rightarrow \quad \varphi^{-1}(y) \geq \sqrt{\frac{-2y}{\beta}}, \quad \text{so that} \quad a = \varphi^{-1} \left(-\frac{v^2}{2 \left(1 + \frac{1}{\mu} \right)} \right) \geq \frac{\underline{v}}{\sqrt{\beta \left(1 + \frac{1}{\mu} \right)}}.$$

and the uniform lower bound behaves like β^{-1} when $\beta \rightarrow \infty$. □

Lemma 2.9. *We probably can do better, but maybe not win any exponent. Suppose that there exists $v_* > 0$ such that $f_{e,b}(v) = 0$ for all $|v| \leq v_*$. Then the density f_i satisfies the uniform bound*

$$f_i(x, v) \leq 2^{9/4} \bar{c} \sqrt{\frac{\frac{\beta}{\alpha} \left(1 + \frac{1}{\mu} \right) + 1}{v_*}} \quad \forall (x, v) \in [0, 1] \times \mathbb{R}.$$

Demonstration Let $(x, v) \in [0, 1] \times \mathbb{R}_-$, and denote by $(x(t), v(t))_{t \leq 0}$ the ion characteristic going through $(x, v) = (x(-T), v(-T))$, with the convention $(x_b(x, v), v_b(x, v)) = (x(0), v(0))$. Owing to the positivity of f_e ,

$$f_i(x(0), v(0)) = \int_{t=-T}^0 f_e(x(t), v(t)) dt + f_i(x, v) \geq f_i(x, v).$$




Figure 2: Notations for the lower bound on n_i .

The coloured area corresponds to the domain $\underline{\varepsilon} \leq \mathcal{L}_e \leq \bar{\varepsilon}$, on which we know that $f_e \geq \underline{c}$. The hatched area represents the domain \mathcal{D} . The solid black line is the critical ion characteristic.

On the other hand, we use $f_i(x, v) + f_i(x, -v) = 2f_i(x(0), v(0))$ to write $f_i(x, -v) \leq 2f_i(x(0), v(0))$. This shows that it is enough to bound f_i on the boundary $\mathcal{B} := \{x \geq 0, v = 0\} \cup \{x = 0, v \leq 0\}$ to obtain a uniform bound.

Let then $(x, v) \in \mathcal{B}$. By hypothesis, $f_e(x(t), v(t))$ vanishes whenever $\mathcal{L}_e(x(t), v(t)) \notin [\mathcal{L}_e(0, v_*), \mathcal{L}_e(0, v^*)]$, and is bounded by \bar{c} otherwise. Then, we may use the reparametrization by space

$$f_i(x, v) \leq \int_{z=a(x,v)}^1 \frac{\bar{c}}{(v^2 + 2\varphi(x) - 2\varphi(z))^{1/2}} dz, \quad \text{with} \quad a(x, v) := \begin{cases} \varphi^{-1}\left(\frac{v^2 + \varphi(x) - \frac{v_*^2}{2}}{1 + 1/\mu}\right) & \text{if } \mathcal{L}_e(x, v) \leq \mathcal{L}_e(0, v_*) \\ x & \text{otherwise.} \end{cases}$$

The function a gives the smallest spatial coordinate of the characteristic $(x(t), v(t))_{t \leq 0}$ such that $f_e > 0$. Using that $\varphi(z) \leq -\alpha \frac{z^2}{2}$, we have

$$f_i(x, v) \leq \bar{c} \int_{z=a(x,v)}^1 \frac{1}{(v^2 + 2\varphi(x) - \alpha z^2)^{1/2}} dz \leq \bar{c} \min\left(\frac{1}{|v^2 + 2\varphi(x)|^{1/2}}, \frac{2}{\sqrt{\alpha a(x, v)}}\right)$$

where we used lemma (2.3) with $L := v^2 + 2\varphi(x)$.

We rely on the concavity estimate

$$\varphi(x) \geq (1-x)\varphi(0) + x\varphi(1) + \frac{\alpha}{2}x(1-x) \geq -x\beta \quad \implies \quad -\frac{y}{\beta} \leq \varphi^{-1}(y)$$

to write that for (x, v) satisfying $\mathcal{L}_e(x, v) \leq \mathcal{L}_e(0, v_*)$,

$$a(x, v) = \varphi^{-1}\left(\frac{\frac{v^2}{2} + \varphi(x) - \frac{v_*^2}{2}}{1 + 1/\mu}\right) \geq \frac{\frac{v_*^2}{2} - (\frac{v^2}{2} + \varphi(x))}{\beta(1 + 1/\mu)}.$$

Then

$$f_i(x, v) \leq \bar{c} \min\left(\frac{1}{\sqrt{|X|}}, \frac{A}{\sqrt{B-X}}\right) \quad \text{with} \quad X := \frac{v^2}{2} + \varphi(x), \quad A := \frac{2}{\sqrt{\frac{\alpha}{\beta(1+1/\mu)}}}, \quad \text{and} \quad B := \frac{v_*^2}{2}.$$

The elementary study of the function $X \rightarrow \min(|X|^{-1/2}, A(B-X)^{-1/2})$ reveals a global maximum at $X = \frac{B}{A^2+1} < B$, and we conclude to the result. \square




Figure 3: Notations for the boundedness of f_i .

The coloured area corresponds to $\mathcal{L}_e(x, v) \geq \mathcal{L}_e(0, v_*)$. The function $a(x, v)$ gives the point of the ion characteristic (in brown) where f_e vanishes.

2.3 Continuity estimates

2.3.1 Electron density n_e

Lemma 2.10. *The electronic density n_e is continuous.*

Demonstration We already know that n_e is bounded. Moreover, using the symmetry $f_e(x, v) = f_e(x, -v)$, we may write

$$n_e(x) - n_e(y) = 2 \underbrace{\int_{v=0}^{v_x(y)} f_e(x, v) dv}_{=: \mathcal{I}^+} + 2 \left(\underbrace{\int_{v=v_x(y)}^{v_x(1)} f_e(x, v) dv - \int_{v=0}^{v_y(1)} f_e(y, v) dv}_{=: \mathcal{I}^-} \right).$$

The term \mathcal{I}^+ is bounded by $\bar{c}v_x(y) = \bar{c} \left(\frac{2}{\mu} (\varphi(x) - \varphi(y)) \right)^{1/2}$, and by continuity of φ , we have $\mathcal{I}^+ \xrightarrow{y \rightarrow x} 0$. On the first integral of \mathcal{I}^- , we apply the change of variable

$$w = \left(v^2 - \frac{2}{\mu} (\varphi(x) - \varphi(y)) \right)^{1/2} \iff dv = \frac{w}{\left(w^2 + \frac{2}{\mu} (\varphi(x) - \varphi(y)) \right)^{1/2}} dw, \quad w \in [0, v_y(1)]$$

to get

$$\mathcal{I}^- = \int_{w=0}^{v_y(1)} f_e \left(x, \left(w^2 + \frac{2}{\mu} (\varphi(x) - \varphi(y)) \right)^{1/2} \right) \frac{w}{\left(w^2 + \frac{2}{\mu} (\varphi(x) - \varphi(y)) \right)^{1/2}} dw - \int_{v=0}^{v_y(1)} f_e(y, v) dv.$$

Notice that

$$f_e \left(x, \left(w^2 + \frac{2}{\mu} (\varphi(x) - \varphi(y)) \right)^{1/2} \right) = f_{e,b} \left(\frac{w^2}{2} + \frac{1}{\mu} (\varphi(x) - \varphi(y)) - \frac{1}{\mu} \varphi(x) \right) = f_e(y, w).$$

Renaming w in v , we obtain the (clearly nonpositive) expression

$$\begin{aligned}\mathcal{I}^- &= \int_{v=0}^{v_y(1)} f_e(y, v) \left(\frac{v}{\left(v^2 + \frac{2}{\mu}(\varphi(x) - \varphi(y))\right)^{1/2}} - 1 \right) dv \geq \bar{c} \int_{v=0}^{v_y(1)} \left(\frac{v}{\left(v^2 + \frac{2}{\mu}(\varphi(x) - \varphi(y))\right)^{1/2}} - 1 \right) dv \\ &= \bar{c} \left(\left(v_y(1)^2 + \frac{2}{\mu} \underbrace{(\varphi(x) - \varphi(y))}_{\geq 0} \right)^{1/2} - \left(\frac{2}{\mu}(\varphi(x) - \varphi(y)) \right)^{1/2} - v_y(1) \right) \geq -\bar{c} \left(\frac{2}{\mu}(\varphi(x) - \varphi(y)) \right)^{1/2}\end{aligned}$$

and this shows that $\mathcal{I}^- \xrightarrow{y \rightarrow x} 0$. \square

Lemma 2.11. *Let n_e^φ and n_e^ψ be the electron densities generated by potentials φ and ψ satisfying the assumptions. Then*

1. *If $f_{e,b}$ is Lipschitz-continuous with constant $[f_{e,b}]$, then*

$$|n_e^\varphi(x) - n_e^\psi(x)| \leq 2[f_{e,b}]v^* \sqrt{\frac{2}{\mu}} |\varphi(x) - \psi(x)|^{1/2} \quad \forall (x, y) \in [0, 1]^2.$$

2. *If there exists a constant $[f_{e,b}]$ such that $|f_{e,b}(x) - f_{e,b}(y)| \leq [f_{e,b}]|x^2 - y^2|$ (or equivalently, $x \rightarrow f_{e,b}(\sqrt{x})$ is a Lipschitz function, as for instance e^{-x^2}), then*

$$|n_e^\varphi(x) - n_e^\psi(x)| \leq \frac{4v^*[f_{e,b}]2}{\mu} |\varphi(x) - \psi(x)| \quad \forall (x, y) \in [0, 1]^2.$$

Demonstration We have

$$|n_e^\varphi(x) - n_e^\psi(x)| \leq 2 \int_{v=0}^{v^*} \left| f_e^\varphi(x, v) - f_e^\psi(x, v) \right| dv = 2 \int_{v=0}^{v^*} \left| f_{e,b} \left(\left(v^2 - \frac{2}{\mu} \varphi(x) \right)^{1/2} \right) - f_{e,b} \left(\left(v^2 - \frac{2}{\mu} \psi(x) \right)^{1/2} \right) \right| dv.$$

Then, if $f_{e,b}$ is Lipschitz with constant $[f_{e,b}]$, we obtain

$$|n_e^\varphi(x) - n_e^\psi(x)| \leq 2[f_{e,b}] \int_{v=0}^{v^*} \left| \left(v^2 - \frac{2}{\mu} \varphi(x) \right)^{1/2} - \left(v^2 - \frac{2}{\mu} \psi(x) \right)^{1/2} \right| dv.$$

Using lemma (2.2) yields

$$|n_e^\varphi(x) - n_e^\psi(x)| \leq 2[f_{e,b}] \int_{v=0}^{v^*} \left| -\frac{2}{\mu} \varphi(x) + \frac{2}{\mu} \psi(x) \right|^{1/2} dv = 2[f_{e,b}]v^* \sqrt{\frac{2}{\mu}} |\varphi(x) - \psi(x)|^{1/2}.$$

If $f_{e,b}(\sqrt{\cdot})$ is Lipschitz with constant $[f_{e,b}]_2$, we may directly write

$$|n_e^\varphi(x) - n_e^\psi(x)| \leq 2v^*[f_{e,b}]_2 \frac{2}{\mu} |\varphi(x) - \psi(x)|.$$

\square

2.3.2 Ion density n_i

Proposition 2.2. *The density n_i is continuous.*

Demonstration Let $0 \leq y < x \leq 1$. For convenience, we represent $n_i(x)$ (resp. $n_i(y)$) as an integral with the artificial lower bound $-g_x(-v^*) \leq -v^*$ (resp. $-g_y(-v^*)$). Then

$$\begin{aligned} n_i(x) - n_i(y) &= 2 \int_{v=-g_x(-v^*)}^0 f_i(x_b(x, v), v_b(x, v)) dv - 2 \int_{v=-g_y(-v^*)}^0 f_i(x_b(y, v), v_b(y, v)) dv \\ &= 2 \underbrace{\left[\int_{v=-g_x(-v^*)}^{-g_x(y)} f_i(x_b(x, v), v_b(x, v)) dv - \int_{v=-g_y(-v^*)}^0 f_i(x_b(y, v), v_b(y, v)) dv \right]}_{=: \mathcal{I}^-} \\ &\quad + 2 \underbrace{\int_{v=-g_x(y)}^0 f_i(x_b(x, v), v_b(x, v)) dv}_{=: \mathcal{I}^+}. \end{aligned} \quad (2.1)$$

The term \mathcal{I}^+ is clearly nonnegative, and may be addressed using our lemmas. Indeed, using the integral representation of $f_i(x_b, v_b)$ and the reparametrization by a space variable z , we have

$$\begin{aligned} \mathcal{I}^+ &= \int_{v=-g_x(y)}^0 \left[\int_{z=x_b(x, v)}^x + \int_{z=x}^1 \right] \frac{f_e(z, -(v^2 + 2(\varphi(x) - \varphi(z)))^{1/2})}{(v^2 + 2(\varphi(x) - \varphi(z)))^{1/2}} dz dv \\ &\leq \bar{c} \int_{v=-g_x(y)}^0 \int_{z=x_b(x, v)}^x \frac{1}{(v^2 - g_x^2(z))^{1/2}} dz dv + \bar{c} \int_{v=-g_x(y)}^0 \int_{z=x}^1 \frac{1}{(v^2 + 2(\varphi(x) - \varphi(z)))^{1/2}} dz dv \\ &\leq \bar{c} \left(2\sqrt{\frac{2}{\alpha}} (\varphi(y) - \varphi(x))^{1/4} \sqrt{x-y} + \frac{2\sqrt{2}}{\alpha^{1/4}} (2(\varphi(y) - \varphi(x)))^{1/2} \right), \end{aligned}$$

where we used lemma (2.6) for the first term, and lemma (2.7) for the second term (with $y = x$ and $v_0 = -g_x(y)$ under the notations of the lemma). Since φ is continuous, we deduce that $\mathcal{I}^+ \xrightarrow[y \rightarrow x]{} 0$. Taking the extreme case $y = 0$ and $x = 1$, we obtain that

$$\mathcal{I}^+ \leq \bar{c} \left(2\sqrt{\frac{2}{\alpha}} (-\varphi(1))^{1/4} + \frac{2\sqrt{2}}{\alpha^{1/4}} (-2\varphi(1))^{1/2} \right) =: K.$$

Let us now focus on \mathcal{I}^- . On the first integral, we make the change of variable

$$w = -(v^2 + 2(\varphi(x) - \varphi(y)))^{1/2} \quad v = -(w^2 + 2(\varphi(y) - \varphi(x)))^{1/2}.$$

Since $\mathcal{L}_i(x, v) = \mathcal{L}_i(y, w)$, this yields $x_b(x, v) = x_b(y, w)$ and $v_b(x, v) = v_b(y, w)$. The bounds $v \in [-g_x(-v^*), -g_x(y)]$ are exactly transported to $w \in [-g_y(-v^*), 0]$. Renaming w in v , we get

$$\mathcal{I}^- = \int_{v=-g_y(-v^*)}^0 f_i(x_b(y, v), v_b(y, v)) \left(\frac{-v}{(v^2 + 2(\varphi(y) - \varphi(x)))^{1/2}} - 1 \right) dv.$$

Since $\varphi(y) \geq \varphi(x)$, the factor of f_i is nonpositive, and so is \mathcal{I}^- . Moreover,

$$n_i(x) - n_i(y) = 2\mathcal{I}^- + 2\mathcal{I}^+ \leq 2\mathcal{I}^- + 2K \iff \mathcal{I}^- = -K + n_i(x) - n_i(y) \geq -K - |n_i|_\infty$$

and \mathcal{I}^- is bounded. The function $\frac{-v}{(v^2 + 2(\varphi(y) - \varphi(x)))^{1/2}} - 1$ converges pointwise to 0 when $x \rightarrow y$ **and f_i is almost everywhere finite**, and by Lebesgue's dominated convergence, $\mathcal{I}^- \xrightarrow[x \rightarrow y]{} 0$. Then n_i is continuous. \square

3 Results

Let $0 < \alpha \leq \beta$, and define the following convex set:

$$\mathcal{K} := \{ \varphi \in \mathcal{C}^1([0, 1], \mathbb{R}) \mid \varphi(0) = \varphi'(0) = 0, \quad \varphi \text{ } \alpha\text{-concave et } (-\beta)\text{-convex.} \}.$$

Both variation conditions rewrite

$$\forall (x, y, \gamma) \in [0, 1]^3, \quad \begin{cases} \varphi((1-\gamma)x + \gamma y) \geq (1-\gamma)\varphi(x) + \gamma\varphi(y) + \frac{\alpha}{2}\gamma(1-\gamma)|x-y|^2, \\ \varphi((1-\gamma)x + \gamma y) \leq (1-\gamma)\varphi(x) + \gamma\varphi(y) + \frac{\beta}{2}\gamma(1-\gamma)|x-y|^2. \end{cases}$$

In the case $\varphi \in \mathcal{C}^2$, it is equivalent to $\varphi'' \in [-\beta, -\alpha]$.

Lemma 3.1. \mathcal{K} is closed for the topology induced by the sup-norm $|\cdot|_\infty := \max_{[0,1]} |\cdot|$.

Demonstration Since $\mathcal{K} \subset \mathcal{C}^0([0,1], \mathbb{R})$, we know that any Cauchy sequence $(\varphi_n)_n \subset \mathcal{K}$ admits a limit $\varphi \in \mathcal{C}([0,1], \mathbb{R})$. The pointwise condition $\varphi_n(0) = 0$ and the pointwise growth conditions are preserved when $n \rightarrow \infty$. Since the family of continuous functions $(\varphi'_n)_n$ is equicontinuous, we may use Arzelà-Ascoli to extract a uniformly converging subsequence $\varphi'_{n_k} \rightarrow \varphi'_\infty$. Using that for all $(x, y) \in [0, 1]$,

$$\left| \varphi(y) - \varphi(x) - \int_x^y \varphi'_\infty(z) dz \right| \leq |\varphi(y) - \varphi_{n_k}(y)| + |\varphi(x) - \varphi_{n_k}(x)| + \int_x^y |\varphi'_\infty(z) - \varphi'_{n_k}(z)| dz \xrightarrow{n \rightarrow \infty} 0,$$

we get that φ'_∞ is the (continuous) derivative of φ . Finally, the uniform convergence $\varphi'_{n_k} \rightarrow \varphi'_\infty$ gives $\varphi'(0) = 0$. \square

We define an operator $F : \mathcal{K} \mapsto \mathcal{C}^2([0,1], \mathbb{R})$ by the solution of the Poisson problem

$$-\lambda^2 F'' = n_i[\varphi] - n_e[\varphi],$$

where n_i and n_e are the ion and electron densities obtained with the characteristics induced by φ .

Stability The function F is solution to a Poisson problem with bounded continuous source term, so it enjoys \mathcal{C}^2 regularity. The boundary condition $F_\varphi(0) = F'_\varphi(0) = 0$ are satisfied by construction. We turn to the stability of the variation estimates: owing to the boundedness of n_i and n_e stated above, we have

$$\begin{cases} -F''(x) \geq \frac{\kappa_1}{\sqrt{\beta(\kappa_2 + \beta)^{1/2}}} - \kappa_3, \\ -F''(x) \leq \frac{\kappa_4}{\alpha^{1/4}} (\beta^{1/4} + \kappa_5) \end{cases}$$

where

$$\kappa_1 := \frac{\nu \sqrt{2} \underline{c} (\bar{v}^2 - \underline{v}^2) \underline{v}}{\lambda^2 \bar{v} \sqrt{1 + \frac{1}{\mu}}}, \quad \kappa_2 := \frac{\bar{v}^2}{2}, \quad \kappa_3 := \frac{2 \bar{c} v^*}{\lambda^2}, \quad \kappa_4 := \frac{4 \nu (1 + \sqrt{2})}{\lambda^2}, \quad \kappa_5 := 4 \sqrt{2} v^*.$$

A sufficient condition for the stability $F(\mathcal{K}) \subset \mathcal{K}$ is to obtain $\alpha \leq -F'' \leq \beta$. We may not always have a solution $(\alpha, \beta) \dots$ It depends on the coefficients.