

Multivariable Calculus

Avery Karlin

Fall 2015

Contents

1	Multivariable Calculus Basics	3
1.1	Green's Theorem	3
1.2	Generalized FTC	3
2	Real Number Set	3
2.1	Definitions of Number Structures	3
2.1.1	\mathbb{N}	3
2.1.2	\mathbb{Z}	4
2.1.3	\mathbb{Q}	4
2.1.4	\mathbb{R}	4
2.2	Basic Structure of \mathbb{R}^1	4
2.3	Properties of \mathbb{R}^n	5
2.4	Basic Axioms for \mathbb{R}^1	5
2.5	The Separation Axiom	6
2.5.1	Proof of Irrationals	6
3	Sequence Theorems	6
3.1	The Least Upper Bound Theorem	6
3.1.1	Proof	6
3.2	Bounded Monotone Sequence Theorem	7
3.2.1	Proof	7
3.3	Archimedean Property	7
3.3.1	Proof	7
3.3.2	Consequences	7
3.4	Sunrise Lemma	7
3.4.1	Well-Ordering Property	8
3.4.2	Proof	8
3.5	Bolzano-Weierstrass Theorem	8
4	Extreme Value Theorem	8

Teacher: Stern

1 Multivariable Calculus Basics

Multivariable calculus is the study of the analogs, if any, of the fundamental theorem of calculus to higher dimensions, and any restrictions that may exist on higher dimensions

Let R be a simple closed curve, where simple means that each point is crossed by the curve at most once and closed means that the curve does not have a unique starting point and end point.

When integrating in 2 dimensions, we also need to pick an “interval.” In this case, R , the bounded region, would be an interval (analogous to (a, b) in single variable calculus) and C would be the boundary (analogous to a, b).

If we were to integrate a function $f(x, y)$ over R , it is denoted by:

$$\iint_R f(x, y) dA_{xy}$$

The dA_{xy} is the area element, an infinitesimally small piece of area, analogous to dx , the length element in single variable calculus.

Note that in single variable calculus, there is an implied orientation, going left to right is the positive “direction,” In multivariable calculus, it is accepted that the positive direction for the curve to go in is the counterclockwise direction, such that the opposite direction adds a negative.

1.1 Green’s Theorem

This is one of the FTC’s generalization to higher dimensions. The Green’s Theorem works with functions that take in 2 variables.

Suppose there exists $f(x, y)$ and $g(x, y)$, and a region R bounded by a positively oriented, simple closed curve C . Then Green’s theorem states that:

$$\iint_R \left(\frac{\partial g}{\partial x} - \frac{\partial f}{\partial y} \right) dA_{xy} = \int_C f dx + g dy = \int_C (f x'(t) + g y'(t)) dt$$

1.2 Generalized FTC

The goal of this course in multivariable calculus is to reach the following conclusion:

For some function ω evaluated over the region M , where ∂M is the boundary of M :

$$\int_M d\omega = \int_{\partial M} \omega$$

2 Real Number Set

2.1 Definitions of Number Structures

2.1.1 \mathbb{N}

We can define the natural number system by sets, like the following:

$$0 = \emptyset$$

And from there we introduce a succession operation:

$$n + 1 = n \cup \{n\}$$

So for example, $1 = \{0\} = \{\emptyset\}$, $2 = \{0, 1\} = \{\emptyset, \{\emptyset\}\}$, etc.

Set theory is the deepest concept in mathematics, such that it must be assumed as postulates.

2.1.2 \mathbb{Z}

Positive integers are defined as an ordered pair of natural numbers, for example, $2 = (2, 0)$, and $-2 = (0, 2)$

2.1.3 \mathbb{Q}

Rationals are defined as an infinite set of ordered pairs of integers. A rational number $q = \frac{n}{m}$, then $q = \{(n, m), (2n, 2m), (3n, 3m) \dots (-2n, -2m), (-3, -3m) \dots\}$

2.1.4 \mathbb{R}

There are several definitions of the real number system

- Each real number can be thought of as an infinite sequence in the following format:

$$(s, N, d_1, d_2, d_3 \dots)$$

Where $s = \pm 1$, $N \in \mathbb{N}$, and $d_n \in 0, 1, 2, 3, \dots, 8, 9$. It is also not the case that $d_n = d_{n+1} = \dots = 0$. If there is a terminal decimal, we express it as 9 repeated.

- Each real number forms a subdivision of \mathbb{Q} into two disjoint sets that cover the entirety of \mathbb{Q} , one of which lies entirely to the left of the other.

2.2 Basic Structure of \mathbb{R}^1

\mathbb{R} is an instance of many kinds of mathematical structures, such as:

- Field
 - A set closed under addition and multiplication (results are within set)
 - Obeys all associated laws of addition and multiplication
- Ordered Set
 - The set has an ordering that reflect the operations in the field (such that $x, y \neq 0$, then $x + y$ and $xy \neq 0$)
 - This structure is what allows for comparisons, like the $<$ function
- Metric Space
 - There exists a standard distance operation between numbers
 - In \mathbb{R} , $dist(x, y) := |x - y|$ ($:=$ means “is defined as”)
- Vector Space
 - Elements can be thought of as vectors from the origin
- Geometric Space
 - This structure means that you can measure angle in a meaningful way
 - In \mathbb{R}^1 , the angle measure can be either 0 or π
 - In higher dimensions there are more angles possible

2.3 Properties of \mathbb{R}^n

In higher dimensions, several of the properties of \mathbb{R} no longer hold. $\mathbb{R}^n, n > 1$ is not a field, and not an ordered set, but it is a vector, metric, and geometric space.

2.4 Basic Axioms for \mathbb{R}^1

\mathbb{R} is a field under $+$ and \cdot ($x, y \in \mathbb{R}$)

1. Additive closure: $x + y \in \mathbb{R}$
2. Associative Property of Addition: $x + (y + z) = (x + y) + z$
3. Communicative Property of Addition: $x + y = y + x$
4. 0 is the identity element of addition: $x + 0 = x$
5. Every element has an additive inverse: $x + (-x) = 0$
6. Multiplicative closure: $xy \in \mathbb{R}$
7. Associative Property of Multiplication: $x(yz) = (xy)z$
8. Communicative Property of Multiplication: $xy = yx$
9. 1 is the identity element of multiplication: $x(1) = x$
10. Every element (except 0) has a multiplicative inverse: $x \cdot \frac{1}{x} = 1$
 - Theorem: $x(0) = 0$
 - Proof: $x(0) = x(0 + 0)$, then we apply the distributive law, and get $x(0) = x(0) + x(0)$. Now we add $-x(0)$ to both side, and we get: $0 = x(0)$
11. Distributive Law: $x(y + z) = xy + xz$

\mathbb{R} is an ordered field and has a proper subset (aka not the entire set) \mathbb{R}^+ (the positives) such that:

1. \mathbb{R}^+ is closed under $+$ and \cdot
2. $1 \in \mathbb{R}^+, 0 \notin \mathbb{R}^+$
3. **Trichotomy Property:** for any $x \in \mathbb{R}$, x is either 0, $\in \mathbb{R}^+$ or $\notin \mathbb{R}^+$
4. Definition of $<$ and $>$:
 - $x < y$ means $y - x \in \mathbb{R}^+$
 - $x > y$ means $y < x$

2.5 The Separation Axiom

The main difference between the \mathbb{Q} and the \mathbb{R} is the separation axiom which the rationals do not have. If $\mathcal{A} \subseteq \mathbb{R}$ and $\mathcal{B} \subseteq \mathbb{R}$, and:

1. $\mathcal{A} \cap \mathcal{B} = \emptyset$
2. $\mathcal{A} \neq \emptyset, \mathcal{B} \neq \emptyset$
3. $\mathcal{A} < \mathcal{B}$ “ \mathcal{A} is to the left of \mathcal{B} ”
 - \forall (for all) $a \in \mathcal{A}, \forall b \in \mathcal{B}, a < b$

2.5.1 Proof of Irrationals

$$\mathcal{A} = \mathbb{Q}^- \cup \{0\} \cup \{q \in \mathbb{Q}^+ | q^2 < 2\}$$

$$\mathcal{B} = \{q \in \mathbb{Q}^+ | q^2 \geq 2\}$$

Then, \exists (there exists at least 1) $c \in \mathbb{R}$ such that $\mathcal{A} \leq c \leq \mathcal{B}$

We know that $\mathcal{A} \neq \emptyset$ and $\mathcal{B} \neq \emptyset$ because 0 is in \mathcal{A} and 2 is in \mathcal{B} . We also know that $\mathcal{A} \cup \mathcal{B} = \mathbb{Q}$ and $\mathcal{A} \cap \mathcal{B} = \emptyset$. As well as the fact that $\mathcal{A} < \mathcal{B}$ (all elements of \mathcal{A} are less than all elements of \mathcal{B}).

Now, if we want to find the boundary element, q_0 , which separates \mathcal{A} and \mathcal{B} . We know that $\mathcal{A} \leq q_0 \leq \mathcal{B}$. So that must mean $q_0 = \sqrt{2}$. However, $\sqrt{2} \notin \mathbb{Q}$. Therefore, we know that the set of rational numbers do not follow the separation axiom.

3 Sequence Theorems

3.1 The Least Upper Bound Theorem

If $\mathcal{A} \subseteq \mathbb{R}$ is non-empty, and is bounded above (So $\exists b_1 \in \mathbb{R}$ such that $\mathcal{A} < b_1$), then \mathcal{A} has a least upper bound, i.e. a number $b_0 \in \mathbb{R}$ such that $\mathcal{A} \leq b_0$ and for any b with $\mathcal{A} \leq b, b_0 \leq b$

$$\mathcal{A} \subseteq \mathbb{R}, \mathcal{A} \neq \emptyset, (\exists b_1 \in \mathbb{R} : \mathcal{A} \leq b_1) \rightarrow [\exists b_0 \in \mathbb{R} : \mathcal{A} < b_0, \forall b \in \mathbb{R} (\mathcal{A} \leq b \rightarrow b_0 \leq b)]$$

b_0 is known as the least possible upper bound, or the *supremum* of \mathcal{A} , we write $b_0 = \sup \mathcal{A}$.

Similarly, for any non-empty set \mathcal{A} bounded below, it has a greatest lower bound, $\inf \mathcal{A}$, called the *infimum* of \mathcal{A}

3.1.1 Proof

Define \mathcal{B} to be the set of all upper bounds of \mathcal{A} . Let $\mathcal{C} = \mathbb{R} \setminus \mathcal{B}$. Clearly \mathcal{B} is nonempty; also \mathcal{C} is non-empty because it contains $x_0 - 1$, where $x_0 \in \mathcal{A}$. By the way in which we defined \mathcal{C} , $\mathcal{B} \cap \mathcal{C} = \emptyset$. Pick any $c \in \mathcal{C}$ and $b \in \mathcal{B}$. By the definition of \mathcal{C} , $\exists x_1 \in \mathcal{A} : c < x_1$. But $x_1 \leq b$ by the definition of \mathcal{B} . Therefore $\mathcal{C} < \mathcal{B}$. By the separation postulate, $\exists b_0 \in \mathbb{R} : \mathcal{C} \leq b_0 \leq \mathcal{B}$. Note that $\mathcal{A} \setminus \{b_0\} \subseteq \mathcal{C}$. Thus, b_0 is an upper bound for \mathcal{A} . Moreover, it is the least upper bound because $b_0 \leq \mathcal{B}$.

3.2 Bounded Monotone Sequence Theorem

For any sequence $\{a_n\}$

1. If $a_n \leq a_{n+1}$ for all $n \geq 1$, and $\exists b \in \mathbb{R}$ such that $a_n \leq b$ for all $n \geq 1$, then $\lim_{n \rightarrow \infty} a_n$ exists and is less than or equal to b
2. If $a_n \geq a_{n+1}$ for all $n \geq 1$, and $\exists b \in \mathbb{R}$ such that $a_n \geq b$ for all $n \geq 1$, then $\lim_{n \rightarrow \infty} a_n$ exists and is greater than or equal to b

3.2.1 Proof

We first convert the sequence $\{a_n\}$, which is bounded by b into the set $\mathcal{A} = \{a_n | n \geq 1\}$. We know that $\mathcal{A} \neq \emptyset$ because the sequence has some terms. We also know that \mathcal{A} is bounded above by b ; $\mathcal{A} < b$

By the Least Upper Bound Theorem, $\exists b_0 = \sup \mathcal{A}$. We now show that $b_0 = \lim_{n \rightarrow \infty} a_n$. By the definition of limits, to say $b_0 = \lim_{n \rightarrow \infty} a_n$ means to say $\forall \epsilon > 0, \exists N > 0, \forall n \geq N, |a_n - b_0| < \epsilon$

If we look at the number $b_0 - \epsilon$, it is not an upper bound on \mathcal{A} because b_0 is the least upper bound and $\epsilon > 0$. Therefore, $\exists a_N > b_0 - \epsilon$. Since $\{a_n\}$ is increasing, $\forall n > N, a_n > b_0 - \epsilon$. If we rearrange the terms, we get $b_0 - a_n < \epsilon$. Therefore, b_0 (which exists by the least upper bound theorem) is the limit of a_n as $n \rightarrow \infty$.

3.3 Archimedean Property

For any positive numbers x and y , it is possible to find some $n \in \mathbb{N}$ such that $nx > y$.

$$\forall x, y > 0, \exists n \in \mathbb{N} : nx > y$$

3.3.1 Proof

Assume that $\neg \exists n : nx > y$, this is logically equivalent to $\forall n : nx \leq y$. Let $\mathcal{C} = \{nx | n \in \mathbb{N}\}$. Then $\mathcal{C} \leq y$, let $c = \sup \mathcal{C}$. We claim that $\exists N : c - \frac{1}{2}x < Nx \leq c$. This is true because if such N does not exist, then $c - \frac{1}{2}x$ would be an upper bound, but c is the least upper bound, so such N must exist. Now we've established the existence of N , let us consider $(N+1)x$. $(N+1)x = Nx + x > (c - \frac{1}{2}x) + x = c + \frac{1}{2}x > c$. But $(N+1)x \in \mathcal{C}$, so it should be $\leq c$. We have a contradiction. This shows that the original assumption is false, so $\forall x, y > 0, \exists n \in \mathbb{N} : nx > y$

3.3.2 Consequences

This property can be used to show that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$. If we consider the definition of limits, the statement is equivalent to saying that $\forall \epsilon \in \mathbb{R} > 0, \exists N \in \mathbb{N}, \forall n > N, n \in \mathbb{N}, \frac{1}{n} < \epsilon$. If we rearrange the term, we get that we need to show $1 < \epsilon N$ for any ϵ . This is true because of the Archimedean Property. $\forall n > N$, since $\epsilon > 0, 1 < \epsilon N < \epsilon n$. Therefore we know the limit is truly 0.

3.4 Sunrise Lemma

The Sunrise Lemma states for any sequence $(a_n)_{n=1}^{\infty} = 1$ in \mathbb{R} , \exists monotone subsequence $(a_{n_k})_{k=1}^{\infty}$, where $(n_k)_{k=1}^{\infty}$ is a strictly increasing sequence in \mathbb{N} and $n_k \geq k$ for all $k \in \mathbb{N}$

Vistas are points in a sequence, a_n , where $N \in \mathbb{N}$, such that $a_N > a_n$ for all $n > N$

This means that for any sequence, there exists a subset of points within, such that within that sequence, the sequence is monotone

3.4.1 Well-Ordering Property

For any set $A \subseteq \mathbb{N}$, $A \neq \emptyset$, $\min(A)$ exists

3.4.2 Proof

Case I: The set V of vistas, is infinite, such that $n_1 = \min(v)$ and $n_k = \min(V \cap n_{k-1}^\infty)$, where $k \geq 2$, then a_{n_k} is strictly decreasing

Case II:

$$n_1 = \begin{cases} 1 & \text{if } v = \emptyset \\ 1 + \max(v) & \text{if } v \neq \emptyset \end{cases}$$

$n_k = \text{choice}\{n > n_{k-1} \mid a_n \geq a_{n_{k-1}}\}$, thus $n_k \neq \emptyset$ because V is finite, thus a_{n_k} is increasing

3.5 Bolzano-Weierstrass Theorem

Every bounded sequence in \mathbb{R} has at least one convergent subsequence

4 Extreme Value Theorem

For some function $f : [a, b]$