Module-Wrestler Online Character Dictionary

Gabe Field

October 23, 2026

Abstract

We are the most modern parody of a long-established dictionary. We relate the language of characters with the language of finite-dimensional complex group modules.

Contents

1 The Translation Dictionary **2**
- 1.1 Definitions in the Language of Characters . . . 2
- 1.2 Grammatical Rules of the Language of Characters . . . 2
- 1.3 Translating Between Characters and Representations . . . 3

2 The Story of Characters, Told in Broad Strokes **5**
- 2.1 Prologue: The Birth of Characters . . . 5
- 2.2 Chapter One: Separating Modules with Inner Products . . . 6
- 2.3 Chapter Two: Orthonormality of Irreducible Characters . . . 6
- 2.4 Chapter Three: Bonus Goodies . . . 8

3 Proofs **10**

4 Some Character Tables **11**

1 The Translation Dictionary

There are categories **FinGroup** of finite groups, **FinVect** of finite-dimensional vector spaces over $\mathbb{C}$, and for each $G : \mathbf{FinGroup}$, a category $\mathbf{FinVect}^G$ of finite-dimensional complex G-modules. If $V : \mathbf{FinVect}^G$, the G-action will be denoted $G \xrightarrow{\rho_V} \mathrm{GL}(V)$ or $g \cdot x := \rho_V(g)(x)$. If it is clear, we may denote $\rho_V(g)$ by $g \cdot -$. The set of conjugacy classes will be denoted $\mathcal{CC}(G)$. For $x \in G$, its conjugacy class is denoted $\mathcal{CL}(x)$.

1.1 Definitions in the Language of Characters

Throughout, $G : \mathbf{FinGroup}$ is a group and $V : \mathbf{FinVect}^G$ is a finite-dimensional complex G-module. (Any other type data will be specified.)

(D1) **Character** of V: The function $\chi_V := \mathrm{tr} \circ \rho_V : G \to \mathbb{C}$. It is a class function (D3), so it is occasionally factored through the quotient $G \twoheadrightarrow \mathcal{CC}(G)$, where it is regarded as a map $\mathcal{CC}(G) \to \mathbb{C}$.

(D2) **Character table** of G: A listing of the characters χ_I corresponding to the finite-dimensional complex irreducible modules I of G. As characters are class functions (D3), the value on each *conjugacy class* is listed. As the irreducible modules biject with the conjugacy classes, this is a square matrix. For example, behold the character table of S_2 (it's not interesting):

Character	$(*)(*)$	$(**)$
χ_{trivial}	1	1
χ_{sign}	1	-1

Beware – although the row for χ_I encodes the character χ_I, it is missing information on the size of the conjugacy class. This information is critical to computing the inner product of two characters (D5); generically, the inner product **does not** agree with the "dot product".

(D3) **Class function**: A function $\phi : G \to A$ which is constant on conjugacy classes of G. A is typically $\mathbb{C}$.

(D4) **Kernel** of the character χ_V: The normal subgroup $\{g \in G \mid \chi_V(g) = \chi_V(1)\}$. More details at (T7).

(D5) **Inner product** on $\mathbb{C}^G$: The vector space of functions $G \to \mathbb{C}$ carries Hermitian inner product $\langle -, - \rangle$ given by

$$\langle \phi, \psi \rangle := \frac{1}{|G|} \sum_{g \in G} \phi(g) \cdot \overline{\psi(g)}$$

or, for characters $(U, V : \mathbf{FinVect}^G)$:

$$\langle \chi_U, \chi_V \rangle := \frac{1}{|G|} \sum_{g \in G} \chi_U(g) \cdot \chi_V(g^{-1})$$

because of (G3).

1.2 Grammatical Rules of the Language of Characters

I mean in this section to list some key properties that characters always satisfy. If you were to communicate using characters, these are some rules that "always appear" in your sentences! Throughout, $G : \mathbf{FinGroup}$ and $U, V : \mathbf{FinVect}^G$ unless otherwise specified. The set of conjugacy classes is denoted $\mathcal{CC}(G)$, and $\mathcal{I}(G)$ denotes the set of irreducible finite-dimensional complex G-modules, listed once per isomorphism class. Conjugacy classes are denoted $\mathcal{CL}(g)$ and centralisers by $\mathcal{CN}(g)$.

(G1) **Column orthogonality of character tables**: Listing a character table with each character along the rows and each conjugacy class along the columns,

$$X := \begin{array}{c} \\ \chi_{\text{trivial}} \\ \chi_{I_1} \\ \vdots \\ \chi_{I_{k-1}} \end{array} \begin{array}{c} \begin{array}{cccc} \mathcal{CL}(1_G) & \mathcal{CL}(g_1) & \cdots & \mathcal{CL}(g_{k-1}) \end{array} \\ \begin{pmatrix} * & * & \cdots & * \\ * & * & \cdots & * \\ \vdots & \vdots & \ddots & \vdots \\ * & * & \cdots & * \end{pmatrix} \end{array}$$

we have that each column is orthogonal under the usual dot product[1]. That is, for $g, h \in G$ with distinct conjugacy classes $\mathcal{CL}(g) \neq \mathcal{CL}(h)$,

$$\sum_{i=0}^{k-1} \chi_{I_i}(g)\overline{\chi_{I_i}(h)} = 0$$

whilst to give the square-norm of a column, we have

$$\sum_{i=0}^{k-1} |\chi_{I_i}(g)|^2 = \frac{|G|}{|\mathcal{CL}(g)|} = |\mathcal{CN}(G)|.$$

(G2) **Dimensions/degrees of irreducibles divide centre indices**: $\dim(V) = \chi_V(1) \mid [G : Z(G)]$.

(G3) **Inverses**: For $g \in G$, $\chi_V(g^{-1}) = \overline{\chi_V(g)}$.

(G4) **Kernels are normal**: $\operatorname{Ker} \chi_V$ is a normal subgroup of G.

(G5) **Frobenius reciprocity**: If $H \leq G$, $U : \mathbf{FinVect}^H$ and $V : \mathbf{FinVect}^G$, then

$$\langle \operatorname{Ind}_H^G U, V \rangle = \langle U, \operatorname{Res}_H^G V \rangle.$$

See (T1).

(G6) **Orthonormality of irreducibles**:

$$U, V \text{ both irreducible} \implies \langle \chi_U, \chi_V \rangle = \mathbf{1}_{U \simeq V} \quad \text{(indicator function)}.$$

(G7) **Orthonormal basis formed by irreducibles**: The vector space $\mathbb{C}^{\mathcal{CC}(G)}$ carries orthonormal basis $\{\chi_I \mid I \in \mathcal{I}(G)\}$. The inner product is **not** the "dot product", but rather as defined at (D5).

1.3 Translating Between Characters and Representations

Throughout, $G : \mathbf{FinGroup}$ and $U, V : \mathbf{FinVect}^G$.

(T1) **Adjunction of induction to restriction**: For a subgroup $H \leq G$, the adjunction of functors $\operatorname{Ind}_H^G \dashv \operatorname{Res}_H^G$ appears as **Frobenius reciprocity**:

$$\mathbf{FinVect}^G(\operatorname{Ind}_H^G U, V) \simeq_{\text{nat.}U,V} \mathbf{FinVect}^H(U, \operatorname{Res}_H^G V) \quad \rightsquigarrow \quad \langle \operatorname{Ind}_H^G U, V \rangle = \langle U, \operatorname{Res}_H^G V \rangle.$$

The inner product relation is a decategorification of the (categorical) adjunction.

(T2) **Commutator**: The commutator subgroup $[G, G]$ is the intersection of all one-dimensional characters.

$$[G, G] = \bigcap_{I \in \mathcal{I}(G).\, \dim I = 1} \operatorname{Ker} \chi_I.$$

[1] This inner product is *not* weighted by the sizes of the conjugacy classes.

(T3) **Dimension/Degree** of V:

$$\dim(V) = \chi_V(1).$$

(T4) **Dimension of hom-space** $\mathbf{FinVect}^G(U,V)$ (this space often goes by $\mathrm{Hom}_G(U,V)$):

$$\dim \mathbf{FinVect}^G(U,V) = \langle \chi_U, \chi_V \rangle.$$

(T5) **Equivalence of irreducible modules**: In the context where U, V are irreducible,

$$U \simeq V \iff \langle \chi_U, \chi_V \rangle = 1$$

and furthermore, in the same context,

$$U \not\simeq V \iff \langle \chi_U, \chi_V \rangle = 0.$$

See (G6) for a more concise statement.

(T6) **Equivalence of modules**:

$$U \simeq V \iff \chi_U = \chi_V.$$

(T7) **Normal subgroups**: For $N \subseteq G$,

$$N \trianglelefteq G \iff \exists \mathcal{J} \subseteq \mathcal{I}(G). N = \bigcap_{I \in \mathcal{J}} \mathrm{Ker}\, \chi_I.$$

(T8) **Frobenius reciprocity**: See (T1).

(T9) **Irreducible factors**: For U irreducible,

$$\text{a factor of } U \text{ appears in } V \iff \langle \chi_U, \chi_V \rangle \neq 0.$$

(A factor "appearing" means that $V \simeq U \oplus (\text{stuff})$.)

(T10) **Irreducibility**: See also (G6).

$$V \text{ is irreducible} \iff \langle \chi_V, \chi_V \rangle = 1.$$

(T11) **Order of centralisers**: For $g \in G$,

$$|\mathcal{CN}(g)| = \sum_{I \in \mathcal{I}(G)} |\chi_I(g)|^2$$

(T12) **Order of group**:

$$|G| = \sum_{I \in \mathcal{I}(G)} |\chi_I(1_G)|^2 = \sum_{I \in \mathcal{I}(G)} (\dim I)^2$$

2 The Story of Characters, Told in Broad Strokes

If you want, this section charts the "etymology" of each word in the dictionary!

The slogan is that **characters decategorify modules**, in analogy to how **dimension decategorifies vector spaces**.

$$\begin{array}{ccc} \text{Categorified} & \xrightarrow{\text{character}} & \text{Decategorified} \\ & & \\ \text{Module } U & \longmapsto & \text{List of numbers } \chi_U \in \mathbb{C}^G \\ \Big\downarrow{\scriptstyle \text{morphism space}} & & \Big\downarrow{\scriptstyle \text{its dimension}} \\ \text{Module } V & \longmapsto & \text{List of numbers } \chi_V \in \mathbb{C}^G \end{array} \qquad \begin{array}{c} \text{What is lost?} \\ \\ \text{Algebraic data (action } G \to \mathrm{GL}(U)) \\ \Big\downarrow{\scriptstyle \text{transformation of maps}} \\ \text{Algebraic data: (action } G \to \mathrm{GL}(V)) \end{array}$$

To exemplify the analogy:

$$\begin{array}{ccc} \text{Categorified} & \xrightarrow{\text{dimension}} & \text{Decategorified} \\ & & \\ \text{Vect. Sp. } U & \longmapsto & \text{Number } \dim U \in \mathbb{Z}_{\geq 0} \\ \Big\downarrow{\scriptstyle \text{morphism space}} & & \Big\downarrow{\scriptstyle \text{rank-nullity statements}} \\ \text{Vect. Sp. } V & \longmapsto & \text{Number } \dim V \in \mathbb{Z}_{\geq 0} \end{array} \qquad \begin{array}{c} \text{What is lost?} \\ \\ \text{Algebraic data (vectors)} \\ \Big\downarrow{\scriptstyle \text{transformation of algebraic data}} \\ \text{Algebraic data (vectors)} \end{array}$$

In both situations, **the decategorification classifies the categorification up to isomorphism** – that is, two (finite-dimensional) vector spaces are isomorphic just when they have the same dimension, and two (finite-dimensional, complex) G-modules are isomorphic just when they have the same character. At the level of objects, the decategorification retains all necessary information up to isomorphism (as one would hope of anything being granted the monacre of "decategorification"). However, passing to this "more computational invariant" has the deficiency of losing algebraic data. On decategorifying **FinVect**, we lose information on how linear maps put together vector spaces. On decategorifying $\mathbf{FinVect}^G$, we lose understanding of the actual actions of G and how these actions intertwine.

2.1 Prologue: The Birth of Characters

We now tell a fiction on the story of character theory, all in broad strokes. Once upon a time, vector spaces were decategorified by their dimension, and everyone lived happily ever after. Well, they lived up until algebraists decided to represent symmetries on vector spaces, moving us from the land **FinVect** to the various lands $\mathbf{FinVect}^G$ (for $G : \mathbf{FinGroup}$). In early and tumultuous days, many discoveries were made paralleling the story for vector spaces:

	Vector spaces	Modules
can be broken up into	one-dimensional subspaces,	irreducible submodules;
the maps between which are	effectively scalars,	effectively scalars;
and which form all objects by	direct sum,	direct sum.

After these initial algebraic gains, however, lingered important questions – "How many irreducible modules are there?", or "Where can we find these modules?", for example. To gain further ground, the category $\mathbf{FinVect}^G$ needed to be quite seriously tamed. After dusting off the history books, the algebraists attempted to decategorify $\mathbf{FinVect}^G$ by mirroring the dimension decategorification of **FinVect**. Fortunately, in the context of a group representation, the dimension appears

naturally:

$$G \xrightarrow{\rho_V} \mathrm{GL}(V) \quad \rightsquigarrow \quad \rho_V(1_G) = 1_V = \underbrace{\begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{pmatrix}}_{n \times n} \quad \rightsquigarrow \quad \dim(V) = n = \mathrm{tr}(\rho_V(1_G)).$$

This is a natural choice because the representing linear transformations $\rho_V(g), g \in G$ are the only extra data present within objects of $\mathbf{FinVect}^G$, and the only such transformation guaranteed to exist is $\rho_V(1_G) = 1_V$. The generalisation easily revealed itself – the **character** of V is the function $\chi_V : G \to \mathbb{C}$ given by $\chi_V := \mathrm{tr} \circ \rho_V$. The miracle is that this successfully decategorified away enough complexity to make tractable progress in understanding the category $\mathbf{FinVect}^G$.

2.2 Chapter One: Separating Modules with Inner Products

The preceding text is best understood as a prologue to our story – "Chapter 0", if you will. In chapter one, we begin to understand the hom-spaces $\mathbf{FinVect}^G(U, V)$ for $U, V : \mathbf{FinVect}^G$. Every part of our story takes place over the category of vector spaces, so it is not a surprise to find that $\mathbf{FinVect}^G(U, V)$ is a vector space. Recording its dimension would be a first feat, and no easy one at that. In perhaps a stroke of luck, or perhaps by arcane insight dug up from historical texts[2], one can find some inspiration in a superficial[3] look at $\mathbf{FinVect}^G(-,-)$:

$$\begin{array}{lll} \text{Categorified} & \mathbf{FinVect}^G \times \mathbf{FinVect}^G \longrightarrow \mathbf{FinVect} & \\ & (U, V) \longmapsto \mathbf{FinVect}^G(U, V), & \\ \text{character} \downarrow & & \\ \text{Decategorified} & \mathbb{C}^G \times \mathbb{C}^G \dashrightarrow \mathbb{C} & \\ & (\chi_U, \chi_V) \longmapsto ?. & \end{array}$$

The superficial symmetry is that $\mathbf{FinVect}^G(-,-)$ appears to decategorify into something resembling an inner product. There is a natural choice of inner product on the set of functions $G \to \mathbb{C}$, given by

$$\begin{aligned} \mathbb{C}^G \times \mathbb{C}^G &\xrightarrow{\langle -,- \rangle} \mathbb{C} \\ (\phi, \psi) &\longmapsto \frac{1}{|G|} \sum_{g \in G} \phi(g) \cdot \overline{\psi(g)}. \end{aligned}$$

The division by $|G|$ may be seen as a normalisation factor; this had proven useful to the algebraists prior. The miracle of chapter one is that *this inner product idea worked* and gave rise to a rich theory, eventually giving the dimension formula

$$\dim \mathbf{FinVect}^G(U, V) = \langle \chi_U, \chi_V \rangle$$

and answering some of the lingering questions prior. How these results follow is not immediate, and will take us another chapter to describe.

2.3 Chapter Two: Orthonormality of Irreducible Characters

A first observation is that

[2] The *Module-Wrestler Online Character Dictionary* keeps no good record of history.
[3] Part of the superficiality is not adequately keeping track of the opposite category.

Lemma 2.1 (Characters of direct sums decompose as sums of characters)
Let $\{V_\alpha\}_{\alpha\in\mathcal{A}}$ be objects of $\mathbf{FinVect}^G$. Then, $\chi_{\oplus_{\alpha\in\mathcal{A}}V_\alpha} = \sum_{\alpha\in\mathcal{A}}\chi_{V_\alpha}$.

Since any module decomposes uniquely into a direct sum of irreducible submodules, we should first study the characters χ_I where I is irreducible. If our inner product $\langle -,-\rangle$ really is to count dimensions of hom-spaces, then Schur's lemma tells us we should expect that

$$\begin{aligned} U \simeq V \text{ are irreducible } &\implies \mathbf{FinVect}^G(U,V) \simeq_{\mathbf{FinVect}} \mathbb{C}^1 \implies \langle\chi_U,\chi_V\rangle = 1, \\ U \not\simeq V \text{ are irreducible } &\implies \mathbf{FinVect}^G(U,V) \simeq_{\mathbf{FinVect}} 0 \implies \langle\chi_U,\chi_V\rangle = 0, \end{aligned}$$

which is to say in other words that if we enumerate the characters as χ_I, one for each isomorphism class of irreducible modules $I : \mathbf{FinVect}^G$, then these characters are orthonormal. Sure enough,

Proposition 2.2 (Orthonormality of irreducible characters)
For irreducible modules $I, J : \mathbf{FinVect}^G$,

$$\langle\chi_I,\chi_J\rangle = \mathbf{1}_{I\simeq J},$$

with the $\mathbf{1}$ denoting an indicator function. (Perhaps think of the Kronecker delta.)

Now, let us collect together one irreducible G-module from each isomorphism class into a set $\mathcal{I}(G)$ (ignore size issues; they can be fixed) and write two direct sum decompositions $U \simeq \bigoplus_{I\in\mathcal{I}(G)} I^{\oplus u_I}$ and $V \simeq \bigoplus_{I\in\mathcal{I}(G)} I^{\oplus v_I}$. Considering the universal properties of the direct sum, Schur's lemma tells us that

$$\begin{aligned} \mathbf{FinVect}^G(U,V) &\simeq \bigoplus_{I,J\in\mathcal{I}(G)} \mathbf{FinVect}^G(I,J)^{\oplus u_I v_J} \\ &\simeq \bigoplus_{I,J\in\mathcal{I}(G)} \left(\mathbb{C}^{\mathbf{1}_{I=J}}\right)^{\oplus u_I v_J} \\ \mathbf{FinVect}^G(U,V) &\simeq \bigoplus_{I\in\mathcal{I}(G)} \mathbb{C}^{\oplus u_I v_I} \\ \implies \quad \dim \mathbf{FinVect}^G(U,V) &= \sum_{I\in\mathcal{I}(G)} u_I v_I \end{aligned}$$

while if we compute the characters, the previous proposition shows us that

$$\begin{aligned} \langle\chi_U,\chi_V\rangle &= \sum_{I,J\in\mathcal{I}(G)} u_I v_J \langle\chi_I,\chi_J\rangle \\ &= \sum_{I,J\in\mathcal{I}(G)} u_I v_J \mathbf{1}_{I=J} \\ \langle\chi_U,\chi_V\rangle &= \sum_{I\in\mathcal{I}(G)} u_I v_I \end{aligned}$$

and therefore,

Theorem 2.3 (Inner products of characters decategorify hom-spaces)
For $U, V : \mathbf{FinVect}^G$, we have

$$\langle\chi_U,\chi_V\rangle = \dim \mathbf{FinVect}^G(U,V).$$

What we have seen also gives us a strategy to extract from a character $\chi_V = \sum_{i\in\mathcal{I}(G)} a_i \cdot \chi_{V_i}$ the coefficients $a_i \in \mathbb{Z}_{\geq 0}$ declaring the "multiplicities" in the direct sum $V \simeq \bigoplus_{I\in\mathcal{I}(G)} I^{\oplus a_I}$ of irreducible modules. One can either leverage orthonormality of the irreducible characters (to extract the coefficients straight away), or leverage the fact that $a_i = \dim \mathbf{FinVect}^G(I,V)$ (which is not hard to prove). Either way,

Theorem 2.4 (Irreducible characters give an orthonormal basis for all characters)
Just as the irreducible submodules give a "direct sum basis" for the finite-dimensional complex G-modules up to isomorphism, the irreducible characters give a "$\mathbb{Z}_{\geq 0}$-weighted basis" for their characters. That is, we have a correspondence

$$\mathbf{FinVect}^G_{/\simeq} \longleftrightarrow \{\text{characters of } G\}$$
$$V \simeq \bigoplus_{I \in \mathcal{I}(G)} I^{\oplus a_I} \longleftrightarrow \chi_V = \sum_{I \in \mathcal{I}(G)} a_I \cdot \chi_I.$$

Read the right way, the previous result entails that

Theorem 2.5 (Equality of characters decategorifies equivalence of modules)
For $U, V : \mathbf{FinVect}^G$, we have

$$U \simeq V \iff \chi_U = \chi_V.$$

Our result 2.7 on a "basis" for the set of characters has a linear algebraic deficiency. It would be nice to interpret this as an actual basis for a vector space, but the set of characters is not closed under scaling by $\mathbb{C}$. Rather than doing something artificial (e.g. taking the $\mathbb{C}$-span of this set), we can consider the vector space $\mathbb{C}^G$. Thinking categorically, we find that

Lemma 2.6 (Characters are class functions)
Just as conjugate (isomorphic) elements x and gxg^{-1} of a group have similar (isomorphic) representing matrices $\rho_V(x)$ and $\rho_V(g)\rho_V(x)\rho_V(g)^{-1}$, the character assigns these elements equal values. That is, $\chi_V(x) = \chi_V(gxg^{-1})$. (This decategorifies isomorphism into equality.)

This does mean that characters cannot span $\mathbb{C}^G$ for non-Abelian G. However, it also shows us a better target for their span.

Theorem 2.7 (Irreducible characters give an orthonormal basis for class functions)
The set $\{\chi_I \mid I \in \mathcal{I}(G)\}$ is an orthonormal basis for $\mathbb{C}^{\mathcal{CC}(G)}$. There is hence one irreducible character of G for each conjugacy class of G. Categorifying, there is (up to isomorphism) one irreducible module of G for each conjugacy class of G.

We can now conclude that $\mathcal{I}(G)$ is a finite set.

2.4 Chapter Three: Bonus Goodies

All good things come in threes, and the missing partner to characters and conjugacy is the left-regular representation $\mathbb{C}[G]$. A telling of this story which does not take such broad strokes would actually reveal, in showing that the irreducible characters span $\mathbb{C}^{\mathcal{CC}(G)}$ (as part of proving 2.7), that the left-regular representation appears. To take yet another broad stroke, the conclusion here is that this representation can be completely torn apart.

Proposition 2.8 (The left-regular representation summarises the irreducible modules)
The left-regular representation $\mathbb{C}[G]$ decomposes as

$$\mathbb{C}[G] \simeq \bigoplus_{I \in \mathcal{I}(G)} I^{\oplus \dim I},$$

or in the language of characters,

$$\chi_{\mathbb{C}[G]} = \sum_{I \in \mathcal{I}(G)} \dim I \cdot \chi_I.$$

Furthermore, the centre of the endomorphism algebra captures the conjugacy classes and irre-

ducible modules together:

$$|\mathcal{CC}(G)| = \dim\left(Z\left(\mathbf{FinVect}^G(\mathbb{C}[G],\mathbb{C}[G])\right)\right) = |\mathcal{I}(G)|.$$

This gives another proof, independent of 2.7, that irreducible characters biject with conjugacy classes.

If your goal is to apply representation theory to distinguish groups, the story told thus far should guide you towards finding characters. The "holy grail" in this avenue is to list the character table of any finite groups you care about (see section 4), and point at different tables to tell apart many groups. However, before you go out on such an odyssey, let us close this story with some words on induction and restriction.

Theorem 2.9 (Frobenius reciprocity)
Fix $H \leq G$ in **FinGroup** and an enumeration $\{x_\alpha H\}_{\alpha\in\mathcal{A}}$ of the cosets in G/H. The adjoint functors $\mathrm{Ind}_H^G \dashv \mathrm{Res}_H^G$ act in characters by

$$\begin{array}{ccccc} \sum_{\alpha\in\mathcal{A}} \hat{\chi}_U(x_\alpha^{-1}(-)x_\alpha) & & \{\text{char's of } G\} & \xleftarrow{\text{char.}} \mathbf{FinVect}^G & \xrightarrow{\text{char.}} \{\text{char's of } G\} \\ \uparrow & & \uparrow & \uparrow\downarrow & \downarrow \\ \chi_U & & \{\text{char's of } H\} & \xleftarrow[\text{char.}]{} \mathbf{FinVect}^H & \xrightarrow[\text{char.}]{} \{\text{char's of } H\} \end{array} \qquad \begin{array}{c} \chi_V \\ \downarrow \\ \chi_V|_H \end{array}$$

where $\hat{\chi}_U : G \to \mathbb{C}$ is given by $\hat{\chi}_U(h) = \chi_U(h)$ for $h \in H$ and $\hat{\chi}_U(g) = 0$ for $g \in G \setminus H$. This gives a formula to define induction and restriction operators on the space of class functions,

$$\begin{array}{ccc} \sum_{\alpha\in\mathcal{A}} \hat{\varphi}(x_\alpha^{-1}(-)x_\alpha) & & \mathbb{C}^{\mathcal{CC}(G)} \\ \uparrow & & \uparrow\downarrow \\ \varphi & & \mathbb{C}^{\mathcal{CC}(H)} \end{array} \qquad \begin{array}{c} \psi \\ \downarrow \\ \psi|_H\,, \end{array}$$

and the formula for ind_H^G is independent of the representatives x_α. The fact that the functors are adjoint appears in characters as the linear operators being Hermitian adjoints.

$$\mathbf{FinVect}^G\left(\mathrm{Ind}_H^G U, V\right) \simeq \mathbf{FinVect}^H\left(U, \mathrm{Res}_H^G V\right) \quad \rightsquigarrow \quad \langle \mathrm{ind}_H^G \chi_U, \chi_V\rangle_G = \langle \chi_U, \mathrm{res}_H^G \chi_V\rangle_H$$

3 Proofs

We collect together proofs of all statements in the translation dictionary and in the "story told in broad strokes" (sections 1 and 2).

4 Some Character Tables