

Does the undirforest depend drastically on the birth order?

On the concluding conjecture of S. A. Alexander, A. de Bruin and D. J. Kornet, *An Alternative Construction of Internodons: The Emergence of a Multi-level Tree of Life* (2015)

25 September 2026

Correction (27 September 2026). An external audit found that the empirical conclusion of §§3 and 5 is overstated, and the statistic I used is to blame. The two proved results (§1, §2) are unaffected.

1. **The all-pairs statistic hides lost co-membership.**
 - The reported 4–14% is the share of *all* organism pairs whose same/different status changes. Most pairs were never co-members, so that figure is small almost by construction.
 - Measured on originally co-clustered pairs, reordering within generations separated **18–32%** of them. The Jaccard overlap of co-member pairs was 0.48–0.73.
 - Single comparisons lost up to 52–61% of original co-memberships. Two independent re-computations with fresh draws agree: `../audit-2026-09-27/controls.json` (the audit’s) and `comembership_check.py` → `comembership-check.json` (this folder).
 - So “species membership almost intact” is **not** warranted. The partition is moderately robust at best.
2. **The “arbitrary order” sampler is not uniform.** `order_robustness_any.py` picks uniformly among the currently available vertices, not uniformly over linear extensions. For $a \rightarrow c$ plus an isolated b , it gives abc $1/4$, acb $1/4$ and bac $1/2$.
3. **One label is a code convention, not the source definition.** An organism that is not SD and has no SD ancestor gets a synthetic singleton label; under Definition 9 it lies in no internodon. In the audited draws, restricting to covered organisms left the co-membership metrics unchanged.

Also, merge-tree interleaving stability does not by itself preserve the internodon labels of fixed organisms. The labelled-pair persistence conjecture (note 09, §5.2) remains open.

Status.

- The merge-tree identification (§1) and the adjacent-swap locality lemma (§2) are hand proofs, each also checked exhaustively on random genealogies.
- §§3 and 5 are finite measurements on one genealogy model.
- No independent review. Whether the merge-tree identification already appears in the phylogenetics literature was not checked.

Source. S. A. Alexander, A. de Bruin and D. J. Kornet, *An alternative construction of internodons: the emergence of a multi-level tree of life*, Bulletin of Mathematical Biology, 2015. [doi:10.1007/s11538-014-0048-2](https://doi.org/10.1007/s11538-014-0048-2). The concluding conjecture (printed pp. 43–44) reads: the

construction takes “a DAG along with distinct birthdates and produced a forest... We suspect that in some sense, biological undirforests do not depend drastically on which topological order is used.”

Scripts in this folder. All are Python 3, using numpy.

- `undirforest.py`: the construction (Definitions 4–5), the join-tree sweep, and a check that the two agree.
- `swap_locality.py` → `swap-locality.json`: the check of the §2 lemma.
- `order_robustness.py` → `order-robustness.json`. Its buffered mode (`python3 order_robustness.py buffered`) → `order-robustness-buffered.json`.
- `order_robustness_any.py` → `order-robustness-any.json`: arbitrary topological orders.
- `comembership_check.py` → `comembership-check.json`: lost co-membership (added with the correction).

Answer in short.

1. The undirforest **is a merge tree** (the augmented join tree of the genealogy graph, with birth order as height). This imports an established stability theory.
2. **Swapping two consecutive, unrelated births changes at most one clade.** It changes nothing when no younger component touches both. This is proved below and checked on 4,302 swaps.
3. Under realistic birth-order uncertainty (reordering within generations), the forest changes a great deal **at the organism level** (51–87% of undirparents change), but the **internodon partition barely moves** (4–10% of organism pairs change status). **[Corrected 27 Sep 2026; see the correction box above.]**

So the conjecture is true “in some sense”: for clades counted one swap at a time, and for species partitions. It is false for parent pointers and for the *size* of the one clade that changes.

1. The undirforest is a merge tree

By Definition 4, x is an undirancestor of y iff an undirected path from x to y avoids x ’s elders: **y lies in x ’s connected component of G restricted to organisms born no earlier than x .** By Corollary 6, y ’s undirparent is its youngest undirancestor.

Now sweep through the organisms from youngest to oldest, adding each to a union-find of the undirected genealogy. When x is added, every current component adjacent to x merges with it. The *oldest vertex so far* of each such component receives x as parent. Then:

- that vertex lies in x ’s component of $G[\text{born} \geq x]$;
- no younger x' has it in its component, since x' would have merged it earlier.

So the parent assigned is the youngest undirancestor. This is exactly the **augmented join tree** (merge tree) of the genealogy graph with height = birth rank, in the sense of Carr–Snoeyink–Axen.

Check: the brute-force Definition-4/5 construction and the join-tree sweep agree on 300 random genealogies with the default order and on 400 under random topological orders ([undirforest.py](#)).

Consequence (established literature, not new): merge trees are stable in the **interleaving distance**, which is bounded by the sup-norm change of the height function (Morozov, Beketayev and Weber, “Interleaving distance between merge trees”, *TopoInVis* 2013). Changing the topological order changes each organism’s rank by at most its displacement. The undirforests of two orders are therefore within interleaving distance **max displacement** of each other, in rank units. That is a precise, already-proved “does not depend drastically” statement, *for a metric that measures merge heights*. Combinatorial (edit-type) distances are a different matter. A 2025 preprint, [arXiv 2508.02352](#), classifies merge-tree edit distances by their stability under

vertex perturbation; only its title was seen here. §2 and §3 measure what happens combinatorially in this setting.

2. Locality of one adjacent swap

Lemma. Let u and v be consecutive in the birth order (u older), with no parent-child edge between them, so swapping them keeps a topological order. Let Y be the set of organisms younger than both, and classify the connected components of $G[Y]$ as:

- S : those adjacent to both u and v ;
- A : adjacent to u only;
- B : adjacent to v only.

Then:

1. **If S is empty, the undirforest is unchanged.**
2. **If S is non-empty,** exactly these parent pointers change: the oldest vertex of each component in S ($v \rightarrow u$); v ($u \rightarrow w$); and u ($w \rightarrow v$). Here w is u 's former parent. In the clade family, **exactly one clade is replaced**: the inner clade $\{v\} \cup S \cup B$ becomes $\{u\} \cup S \cup A$. The outer clade $\{u, v\} \cup S \cup A \cup B$, and every other clade, are unchanged.

Proof. In the sweep, v is added immediately before u .

- *When S is empty,* u 's merges and v 's merges involve disjoint sets of components. u and v are not adjacent, and their merged components do not touch. Both orders therefore assign the same parents and leave the same union-find partition with the same oldest vertices. The rest of the sweep is identical.
- *When S is non-empty,* the original order assigns:
 - parent v to the oldest vertices of S and B ;
 - then parent u to those of A ;
 - and parent u to v , since the merged component touches u .

The swapped order assigns:

- u to the oldest vertices of S and A ;
- v to those of B ;
- v to u .

In both orders the partition after the pair is the same; only its oldest vertex differs (u originally, v after the swap). So the vertex w that later receives it as a child is the same. All other assignments coincide. The subtrees of all other vertices contain both of u , v or neither, so their clades are unchanged. ■

Check: on 4,302 valid adjacent swaps in random genealogies, the forest changed exactly when S was non-empty (4,302 of 4,302), and every change replaced exactly one clade (1,637 of 1,637) ([swap-locality.json](#)).

Corollary. Any two topological orders are joined by a sequence of adjacent swaps of incomparable consecutive organisms. Bubble sort does it, swapping only differently-ordered pairs, which are incomparable. The number of swaps is the number of differently-ordered pairs. So the clade families of the two undirforests differ by at most $2 \times$ **(number of swaps whose S is non-empty)** clades, in the Robinson-Foulds sense.

What stays drastic. The replaced clade can be large: A and B may contain whole descendant subtrees. In the random test genealogies, the changed clade averaged 58% of the population. One swap moves at most one clade, but possibly a big one.

3. Measured on genealogies with a permanent split

These are sexual Wright-Fisher-type generations. One ancestral deme of size N splits permanently into two demes at generation 6, with 14 generations in all. Each child has two parents with probability “sex”, otherwise one. Organisms of the same generation are mutually incomparable, so *every within-generation reordering is a valid topological order*: this is the realistic birth-order uncertainty. For each setting: 6 genealogies $\times$ 12 random reorderings, compared with a reference order ([order-robustness.json](#)).

N	sex	organisms	undirparents changed	clades not shared	internodon pairs changed	internodons (ref / alt)
6	0.3	132	51%	32%	8.0%	12.8 / 13.5
6	1.0	132	78%	58%	6.3%	9.5 / 9.8
12	0.3	264	56%	44%	6.1%	32.0 / 32.7
12	1.0	264	83%	70%	6.2%	20.7 / 19.4
24	0.3	528	59%	49%	4.4%	63.0 / 63.5
24	1.0	528	85%	74%	6.0%	39.8 / 42.1

(The full table, including sex = 0.7, is in the JSON.)

With a buffer. The finite-window SD property counts lineages that would die out later, and the paper’s permanent splits need the infinite future. So the buffered variant evaluates only the first 10 generations and simulates 4–12 further generations after them.

Internodons in the evaluated part fall from 11–21 to about 5–15. The pair-change rate stays at **4–10%**, while 70–87% of undirparents change ([order-robustness-buffered.json](#)).

Reading. Birth-order uncertainty *within* a generation rewires most organism-level undirparents. That is expected: the forest’s chains through a generation follow birth order. Yet it leaves species membership (internodon partition) almost intact. **[Corrected 27 Sep 2026; see the correction box above.]** That is the paper’s “in some sense”: robust where biology looks (the partition into internodons), not where the construction does its bookkeeping (who is whose undirparent).

4. An observation about the definitions

Checking whether internodons respect a permanent deme split turned up a feature of Definitions 7–9, not a code error.

- **How it happens.** The pre-split ancestral segment can continue unbranched into one daughter deme. A childless (hence non-SD) organism of the *other* daughter then joins the internodon of its youngest SD ancestor, which lies on that ancestral segment.
- **The result.** An internodon can contain organisms from both sides of a permanent split, through non-SD attachment. Whether Kornet’s internodons (equivalent by Theorem 20) should allow this is a question about the intended treatment of non-SD organisms. It is recorded here and not resolved.
- **The finite window adds noise.** Organisms that die childless inside it are non-SD.

Status and limits

- **Proved here:** the identification with the join tree, and the adjacent-swap locality lemma with its corollary. Both are hand proofs, each also checked exhaustively on random genealogies. Whether the merge-tree identification appears in the phylogenetics literature was not checked.
- **Imported, established:** stability of merge trees in the interleaving distance (Morozov et al. 2013). It was not re-proved here, and its exact hypotheses (vertex-valued height on a graph) should be confirmed before the corollary is quoted.
- **Measured, finite:** internodon robustness on one genealogy model. The paper’s infinite biosphere and permanent-split data are approximated by a finite window with a buffer.
- **Not done:**
 - reorderings *across* generations (for example overlapping generations with continuous birthdates);
 - the paper’s second suggestion, extending the construction down to gene trees.

5. Arbitrary topological orders

[order_robustness_any.py](#) → `order-robustness-any.json`. Same genealogy model; the alternative orders are random topological orders of the whole parent DAG (not uniform over linear extensions; see the correction box) (the continuous-birthdate case, where uncertainty spans generations). 5 genealogies × 10 orders.

N	sex	undirparents changed	clades not shared	internodon pairs changed	mean rank displacement (share of n)
6	0.3	57%	50%	14.5%	9.0%
6	1.0	83%	76%	10.7%	5.5%
12	0.3	62%	55%	10.4%	9.7%
12	1.0	87%	79%	11.9%	5.8%

Internodon co-membership moves about twice as much as under within-generation reorderings (10–14% of pairs against 4–8%), but most of it still survives. The conjecture’s “in some sense” therefore depends on how much birth-order uncertainty is admitted:

- **within generations:** robust;
- **across generations:** moderately robust. Neither case is robust at the level of parent pointers.