

Clinical Cognition Deep Dive (Attending Physician Perspective)

Case: 23-year-old male with post-COVID bilateral hydropneumothorax, pneumomediastinum, subcutaneous emphysema, and respiratory failure

Module CC — Clinical Cognition Deep Dive

1. Executive Clinical Summary

A previously healthy 23-year-old food delivery worker developed COVID-19 pneumonia approximately one month before presentation. Initial HRCT demonstrated severe bilateral viral pneumonia with a crazy-paving pattern together with pneumomediastinum and extensive subcutaneous emphysema, suggesting alveolar rupture (Macklin effect).

After apparent virological recovery (COVID negative after 20 days), he developed recurrent cough, dyspnea and eventually sudden deterioration with hemoptysis and severe hypoxemia. Chest radiograph demonstrated left pneumothorax requiring emergency ICD insertion.

Subsequent HRCT revealed:

- Bilateral loculated hydropneumothoraces
- Partial lower lobe collapse
- Persistent bilateral infective air-space disease
- Right lower lobe bulla
- ICD in situ

Despite chest drainage he continued to deteriorate because of bilateral loculated pleural collections, eventually requiring thoracic surgical opinion for thoracoscopy and decortication.

Overall this represents severe post-COVID pulmonary structural destruction complicated by persistent alveolar air leak and infected pleural disease.

2. Problem Representation

Young male

↓

Recent severe COVID pneumonia

↓

Diffuse alveolar injury

↓

Alveolar rupture

↓

Pneumomediastinum + subcutaneous emphysema

↓

Subpleural bulla formation

↓

Bilateral hydropneumothorax

↓

Persistent air leak

↓

Secondary bacterial pneumonia

↓

Respiratory failure

3. Semantic Qualifiers

Young

Acute on chronic

Post-viral

Structural lung injury

Severe hypoxemia

Bilateral disease

Persistent air leak

Complicated pleural disease

Secondary infection

Rapid physiological deterioration

4. Illness Script

Epidemiology

Young adults with severe COVID may rarely develop:

- diffuse alveolar damage
- spontaneous pneumothorax
- pneumomediastinum
- persistent air leak syndrome

especially when extensive peripheral lung involvement exists.

Predisposing Factors

Recent COVID pneumonia

Diffuse alveolar injury

Peripheral lung involvement

Formation of fragile subpleural bullae

Possible excessive coughing

Pathophysiology

Diffuse viral inflammation

↓

Alveolar rupture

↓

Air dissects along bronchovascular bundles

↓

Macklin effect

↓

Pneumomediastinum

↓

Subcutaneous emphysema

↓

Pleural rupture

↓

Pneumothorax

↓

Persistent bronchopleural communication

↓

Hydropneumothorax

↓

Pleural infection

↓

Loculations

↓

Restrictive lung mechanics

↓

Respiratory failure

Clinical Features

Progressive dyspnea

Productive cough

Hemoptysis

Chest pain

Hypoxemia

Reduced air entry

Tachypnea

Tachycardia

5. Clinical Timeline

One month prior

COVID positive

↓

HRCT:

- Crazy paving
- Pneumomediastinum
- Subcutaneous emphysema

↓

Medical treatment

↓

COVID negative

↓

Recurrent cough



Dyspnea



Acute collapse



Left pneumothorax



Emergency ICD



Persistent dyspnea



CT:

Bilateral loculated hydropneumothoraces



Thoracic surgery referral



Clinical deterioration

6. Differential Diagnosis

Most likely

1. Post-COVID persistent alveolar air leak syndrome ★★★★★

Supported by:

- pneumomediastinum
 - subcutaneous emphysema
 - pneumothorax
 - bulla
-

2. Secondary bacterial necrotizing pneumonia ★★★★★☆

Supported by

Leukocytosis

Hydropneumothorax

Persistent infiltrates

3. Bronchopleural fistula ★★★★★☆

Persistent air leak despite ICD

Hydropneumothorax

4. Empyema with air-fluid level ★★★★★☆

Loculated collections

Persistent infection

Less likely

Pulmonary embolism

Tuberculosis

Vasculitis

Lung abscess

Malignancy

7. Interpretation of Investigations

HRCT (Initial)

Crazy paving

Diffuse alveolar damage

Severe COVID

Pneumomediastinum

Subcutaneous emphysema

Suggests early alveolar rupture.

HRCT Follow-up

Hydropneumothorax

Loculations

Lower lobe atelectasis

Persistent infiltrates

Bulla

Demonstrates evolution from viral injury into structural pulmonary destruction.

D-Dimer 1940

May indicate

COVID inflammation

Thrombosis risk

Fibrinolysis

Not diagnostic of PE alone.

CRP Negative

Interesting discordance.

Could indicate

Resolving viral inflammation

Localized pleural infection

Prior antibiotic exposure

CBC Trend

Initial leukocytosis

↓

Normalization

↓

Hemoglobin fall

Likely due to:

Repeated blood sampling

Hemoptysis

Critical illness

Inflammation

8. Clinical Reasoning

The pivotal diagnostic clue is **pneumomediastinum before pneumothorax**.

This suggests alveolar rupture rather than primary pleural disease.

Later:

persistent air leakage

↓

formation of bullae

↓

hydropneumothorax

↓

infection

↓

loculations

↓

failure of conservative drainage

This explains why chest tube alone could not completely resolve the pathology.

9. Management Appraisal

Strengths

Prompt oxygen therapy

Emergency ICD insertion

Serial imaging

Pulmonology referral

Broad-spectrum antibiotics

Supportive care

Thoracic surgery consultation

Areas for Optimization

Assessment for persistent bronchopleural fistula

Pleural fluid analysis (cell count, Gram stain, culture, biochemistry)

Pleural ultrasound to define septations

CT-guided drainage if indicated

Consider intrapleural fibrinolytic therapy (selected cases)

Earlier thoracic surgery evaluation once loculations became evident

Formal VTE evaluation because of elevated D-dimer

Pulmonary rehabilitation planning after stabilization

10. Learning Pearls

COVID can produce spontaneous pneumothorax even without mechanical ventilation.

Pneumomediastinum often precedes pneumothorax via the Macklin effect.

Persistent dyspnea after COVID should prompt evaluation for structural complications.

Loculated hydropneumothorax frequently requires more than simple chest tube drainage.

Young patients can develop severe post-COVID pulmonary sequelae despite eventual viral clearance.

11. Cognitive Bias Check

Anchoring Bias

Assuming recurrent dyspnea was simply persistent COVID pneumonia.

Avoided by: obtaining repeat imaging.

Premature Closure

Stopping after identifying pneumothorax.

Avoided by: repeat CT demonstrating bilateral loculated disease.

Availability Bias

Attributing all respiratory symptoms to COVID.

Avoided by: considering secondary bacterial infection and pleural complications.

Search Satisfaction

Finding one pneumothorax could have halted further investigation.

Avoided through: HRCT revealing bilateral pathology and bullous disease.

12. Systems Thinking

Primary pathology: Diffuse alveolar damage

↓

Mechanical complication: Air leak syndrome

↓

Pleural complication: Hydropneumothorax

↓

Infectious complication: Secondary pneumonia



Mechanical restriction: Loculations + atelectasis



Physiological consequence: Hypoxemic respiratory failure

13. Expert Attending Assessment

This case exemplifies the progression from severe viral lung injury to a complex post-COVID air leak syndrome. The chronology of pneumomediastinum followed by pneumothorax strongly supports alveolar rupture as the initiating event. Persistent bilateral loculated hydropneumothoraces, combined with evidence of bullous disease and ongoing respiratory compromise, suggest a continuing bronchopleural communication complicated by secondary infection.

Initial emergency management was appropriate, including oxygen therapy, intercostal drainage, antibiotics, and multidisciplinary referral. However, the persistence of bilateral loculated collections indicates that definitive management would likely require advanced pleural intervention, including thoracoscopy with decortication and evaluation for bronchopleural fistula.

From a clinical reasoning standpoint, this case highlights the importance of repeatedly reassessing evolving pulmonary pathology rather than attributing persistent respiratory symptoms solely to viral pneumonia. Sequential imaging, awareness of post-COVID structural complications, and early multidisciplinary collaboration are critical to optimizing outcomes in similar patients.

Final Teaching Points

Working Diagnosis

Post-COVID diffuse alveolar injury complicated by:

- Persistent alveolar air leak syndrome
 - Pneumomediastinum
 - Bilateral loculated hydropneumothorax
 - Secondary bacterial pneumonia
 - Possible bronchopleural fistula
 - Acute hypoxemic respiratory failure
-

Key Clinical Reasoning Takeaways

- Recognize pneumomediastinum as an early marker of alveolar rupture.
- Consider structural complications when dyspnea recurs after apparent COVID recovery.
- Persistent air leak after ICD should prompt evaluation for bronchopleural fistula.
- Bilateral loculated hydropneumothorax often necessitates surgical intervention beyond chest tube drainage.
- Sequential reassessment and multidisciplinary management are essential in complex post-COVID pulmonary sequelae.