

HARW Beta 15.0

Dummy Title eleven

--Manuscript Draft--

Manuscript Number:	
Full Title:	Dummy Title eleven
Article Type:	Original Article
Corresponding Author:	* Amy Author Springer Heidelberg, GERMANY
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Springer
Corresponding Author's Secondary Institution:	
First Author:	* Amy Author
First Author Secondary Information:	
Order of Authors:	* Amy Author
Order of Authors Secondary Information:	
Author Comments:	Test
Suggested Reviewers:	

Role of perforin secretion from CD8⁺ T-cells in neuronal cytotoxicity in multiple sclerosis 31

Daidi Zhao ^{1#}, Fuqiang Feng ^{2#}, Cong Zhao ¹, Fang Wu ³, Chao Ma ⁴, Yanan Bai ¹, Jun Guo ^{1*}, Hongzeng Li ^{1*}

1-Department of Neurology, Tangdu Hospital, Fourth Military Medical University, Xi'an, Shaanxi, 710038, China

2-Department of Neurosurgery, Tangdu Hospital, Fourth Military Medical University, Xi'an, Shaanxi, 710038, China

3-Department of Neurology, Xi'an Children's Hospital, Xi'an, Shaanxi, 710003, China

4-Department of Cardiology, Tangdu Hospital, Fourth Military Medical University, Xi'an, Shaanxi, 710038, China

Daidi Zhao and Fuqiang Feng contributed equally to this study.

* Corresponding authors

Email: lihongzeng_fmму@163.com

Fax & Tel. +86 02984777435.

Conflict of interest: The authors have no conflict of interest and no outside funding to declare.

Abstract

Multiple sclerosis is the most prevalent autoimmune disease of the central nervous system characterized by the presence of inflammation, myelin damage. While the immune system initiates the autoimmune response it remains unclear how it orchestrates neuronal damage. In previous studies, the presence of autoreactive CD4⁺ and CD8⁺ T lymphocytes, together with other inflammatory cells and cytokines in active MS lesions, suggest an autoimmune pathogenesis. In an extension of those studies, here we show that the role of CD8⁺ T-cell in MS and confirmed the perforin is

an important factor. Furthermore, we determined that the CD4⁺ T-cell can enhance CD8⁺ T-cell cytotoxicity to neurons by induction intense inflammation.

Keywords: Multiple sclerosis, CD4⁺ T-cell, CD8⁺ T-cell, Perforin, neurodegeneration

Introduction

Although multiple sclerosis (MS) was first diagnosed in 1894, the underlying pathomechanism of the disease remains unknown, and little evidence has been published regarding the efficacy of radical treatment options. Studies have demonstrated that focal demyelinating lesions are not restricted to the white matter in MS, but also affect the gray matter and normal-appearing white matter, suggesting that both axonal and neuronal loss are major causes for irreversible neurological disability[1].

Inflammation is present at all stages of MS, leading to the predominant infiltration of macrophages, followed by CD8⁺ T-cells and lower numbers of CD4⁺ T-cells, B cells, and plasma cells[2]. Progression of the disease results in diffuse inflammatory T-cell and B cell infiltration, microglia and astrocyte activation, and diffuse myelin reduction and axonal injury. These findings are associated with more pronounced atrophy of the gray and white matter.

Research has indicated that CD8⁺ T-cells exhibit oligoclonal expansion and are the most abundant T-cells in CNS lesions of patients with MS[3-6]; however, the function of these cells in MS remains unclear and somewhat controversial. The pathogenic role of CD8⁺ T-cells in MS first became evident in a study in which an EAE model was developed using the self-protein myelin basic protein (MBP) [7]. Various CNS-specific and therapeutically induced CD8⁺ T-cell responses have also been observed in patients with MS[8-11].

Accumulating evidence has suggested that neurons indirectly interact with T-cells via microglia in the CNS[12]. In a recent study, researchers recorded T-cell adhesion *in vivo* using PLP139-151-specific T-cells, observing a direct effect of T-cells on neurons

1 in mouse brain slices[13]. The authors observed that T-cells directly attacked neurons,
2 resulting in glutamate receptor-mediated toxicity, which in turn resulted in calcium
3 overload and neuronal death. However, as brain sections contain multiple rather than
4 isolated neurons, it is difficult to determine the exact process by which damage to
5 other brain cells occurs.
6
7
8
9

10 In our previous study, we observed that myelin basic protein (MBP)-induced
11 lymphocytes can be applied directly to neurons of the brain *in vitro*[14]. Most studies
12 that have addressed the immunobiological aspects of multiple sclerosis (MS) in
13 animal models of experimental autoimmune encephalomyelitis (EAE) have focused
14 on CD4+ T-cells as the main orchestrators of pathogenesis and regulation. In 1984,
15 Takenaka et al. transferred CD4+ T-cells in murine animals, observing that adoptive
16 transfer of CD4+T-cells isolated from myelin antigen-primed animals is sufficient to
17 induce disease in animal models of EAE[15]. In 1989, Waldmann et al. reported that
18 CD4+ T-cells from peripheral blood attack myelin in the CNS, leading to the
19 development of MS. Therefore, studies have aimed to prevent disease recurrence and
20 progression by blocking or inhibiting CD4+ T-cells in animal models of MS. Further
21 studies have aimed to selectively remove CD4 + T-cells from the blood to treat MS in
22 both the acute and recurrent stages[16]. In 2013, Zaguia et al. reported that
23 NK-associated molecules expressed by human CD4+ T-cells confer
24 oligodendrocyte-directed cytotoxicity. A significantly elevated proportion of *ex vivo*
25 peripheral blood CD4+ T-cells from patients with MS express NKG2C when
26 compared with controls, suggesting a novel mechanism via which
27 infiltrating CD4+ T-cells may contribute to tissue injury in MS[17].
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 Although studies have revealed that T-cells contribute to neuronal apoptosis and
49 axonal degeneration in MS, the precise mechanism by which such damage occurs
50 remains unclear, as some damage to the gray matter occurs in the very early stages of
51 MS, despite the normal appearance of the white matter. Both CD4+ and CD8+ T-cells
52 play important roles in MS, though their relationship to neuronal damage remains
53 unclear. In the present study, we aimed to clarify the function of T-cell subgroups in
54
55
56
57
58
59
60
61
62
63
64
65

1 the pathogenesis of MS in order to provide an experimental basis for preventing the
2 loss of nerve function in patients with MS.

3 **Methods**

4 **EAE animal models**

5
6
7
8
9 Adult male Lewis rats were injected with myelin basic protein (MBP, Sigma)
10 emulsified with Complete Freund's Adjuvant (CFA). Plantar injections of 100 µg
11 MBP were administered on each side (200 µg per rat). The state of EAE was
12
13
14
15
16 monitored each day.

17 **Cell cultures**

18
19
20
21 Purified CD4⁺ or CD8⁺ T-cells were isolated from heparinized peripheral blood
22
23 (EAE animals and normal animals) via negative selection using RosetteSep
24
25 (STEMCELL technologies, Grenoble, France), followed by a Ficoll density-gradient
26
27 centrifugation. Purity of the isolated CD4⁺ or CD8⁺ T-cells within the T-cell
28
29 population usually is between 97.8%–99.4%.
30

31
32
33 CD4⁺ cells were cultured with APC cells and stimulated with MBP87-99
34
35 [Val(87)-His-Phe-Phe-Lys-Asn-Ile-Val-Thr- Pro-Arg- Thr-Pro(99)], while CD8⁺ cells
36
37 were directly cultured with MBP87-99. These cells were then co-cultured with
38
39
40 neurons.
41

42 **ELISA**

43
44
45 Concentrations in cell culture supernatants were measured via enzyme-linked
46
47 immunosorbent assay (ELISA) using the BD OptEIA™ Set Mouse kit (BD
48
49 Biosciences, San Diego, USA), in accordance with manufacturer instructions.
50

51 **Western blot analysis**

52
53
54 Stimulated cells were lysed on ice in lysis buffer (15 mM PBS, 2% NP-40, 0.2% SDS,
55
56
57 10 mM EDTA, 1% protease inhibitor cocktail, 1% phosphatase inhibitor cocktail).
58
59
60 The solution was then centrifuged at 14,000 g for 15 min at 4°C, following which the
61
62
63
64
65

1 supernatant was collected and protein concentration was evaluated via BSA assay.
2 Proteins were resolved on 10% acrylamide SDS-polyacrylamide gel electrophoresis
3 and then electro-transferred to nitrocellulose membranes for Western blot (WB)
4 analysis. The membranes were blocked in a 5% milk solution in TBS (0.1% Tween 20)
5 and incubated overnight at 4°C with primary antibodies for Bax (1:1000 dilution,
6 Santa Cruz Biotechnology, Santa Cruz, CA, USA). Samples underwent incubation
7 with HRP-conjugated secondary anti-body for 2 h at 25 °C, followed by
8 chemiluminescence reaction performed via the use of electrochemiluminescence and
9 film exposure. Expression of β -actin was used as a loading control.
10
11
12
13
14
15
16
17
18
19
20
21

22 **TUNEL**

23
24 The apoptosis of stimulated neurons with was evaluated using a TUNEL Apoptosis
25 Assay Kit (Roche,TUN11684817), in accordance with manufacturer instructions.
26
27
28

29 **MTT**

30
31 Stimulated cells were cultured in 96-well plates. After 24 h, 20 μ l MTT solution (5
32 mg/ml prepared with PBS, pH = 7.4) was added, and cells were incubated for an
33 additional 4 h. The supernatant was carefully extracted, and 150 μ l DMSO was
34 added to stop the reaction. The OD value was tested at 490 nm.
35
36
37
38
39
40
41

42 **Statistical analysis**

43
44 Values are expressed as the mean \pm SE. Statistics as indicated in the figure legends
45 were calculated using GraphPad Prism version 5.01 for Windows (GraphPad Software,
46 San Diego, USA). A *p* value of <0.05 was statistically significant.
47
48
49
50
51
52
53

54 **Results**

55 **Differential cytotoxic effects of CD4⁺ and CD8⁺ T-cells on neurons**

56
57 To address which type of T-cells play an important role in neuronal cell death in MS,
58
59
60
61
62
63
64
65

we co-cultured T-cells stimulated by MBP peptide *in vitro* (MBP group) or T-cells from EAE models (EAE group) with neurons. The degree of neuronal cell death was examined after 24 hours via TUNEL staining. Our results revealed that CD8⁺ T-cells were more toxic to neurons than CD4⁺ T-cells in both the MBP and EAE, although these effects were more pronounced in the MBP group (Figure 1). These data demonstrate that CD8⁺ T-cells, but not CD4⁺ T-cells, play a major role in neuronal cell death in MS.

The expression of Bax in neurons co-cultured with different T-cell types

To further confirm that CD8⁺ T-cells play a greater role than CD4⁺ T-cells in neuronal cell death associated with MS, we examined the expression of Bax in neurons co-cultured with different T-cell types. We observed that, compared with controls, the level of Bax was slightly increased when neurons were co-cultured with CD4 in the MBP group, though a greater increase was observed in neurons co-cultured with CD8 (Figure 2 A and B). Similar results were observed in the EAE group (Figure 2 C and D). These data suggest that CD8⁺ T-cells provide a greater contribution to neuronal cell death than CD4⁺ T-cells in MS.

Expression of various factors secreted by different T-cell types co-cultured with neurons

In order to determine which factors secreted by each type of T-cell provided the greater contribution to neuronal cell death, we examined the secretion of various factors via ELISA. We observed significant increases in interleukin 17 (IL-17) secretion by CD4⁺ T-cells in both the MBP group and EAE group (Figure 3 A). Further, we observed increases in the secretion of perforin (Figure 3 B), granzyme B (Figure 3 C), and IFN- γ (Figure 3 D) by CD8⁺ T-cells. These results suggest that perforin, IL-17, and/or IFN- γ may play a primary role in neuronal cell death in MS.

Direct effect of various factors on neuronal cell death

To examine which of these factors was associated with damage to neurons, we examined their effects on neuronal cell death via MTT and Western blotting. Our findings indicated that perforin played a more important role than other factors: The MTT assay revealed that the viability of neurons treated with perforin was worse than that of other groups (Figure 4 A). Furthermore, Bax expression was significantly higher in neurons treated with perforin than under other conditions (Figure 4 B, $*P < 0.05$). These data suggest that perforin was more toxic to neurons.

Discussion

In the present study, neurons were co-cultured with CD4⁺ T-cells and CD8⁺ T-cells separated from EAE or MBP models to determine which T-cell type provides the greatest contribution to neuronal cell death in MS. We observed that CD8⁺ was associated with a significantly greater degree of apoptosis than CD4⁺ T-cells, and that the combination of CD8⁺ and CD4⁺ cells enhanced this apoptotic effect. Thus, these findings suggest that CD8⁺ T-cells may act to directly induce neuronal apoptosis via action on both neurons and axons in the CNS, potentially providing new insight into the pathogenesis of MS.

In the present study, we observed that simultaneous application of CD8⁺ and CD4⁺ T-cells resulted in increased neural apoptosis relative to application of CD8⁺ T-cells alone[3]. Several previous studies have suggested that CD8⁺ and CD4⁺ T-cells exhibit functional interactions in the context of MS, although a growing body of evidence suggests that these interactions play a regulatory role in the pathogenesis of MS and MS-like disease. Jiang et al. reported that CD8⁺ T-cells modulated disease in an EAE model via actions on CD4⁺ T-cells. Previous researchers have also demonstrated that CD4⁺ T-cell vaccination protocol-mediated protection against EAE disease is dependent on the presence of Qa-1-restricted CD8⁺ T-cells[18]. Other studies have suggested that these CD8⁺ T-cells may induce cell death in GA-loaded

1 target T-cells and limit the proliferation of CD4⁺ T-cells[19], thereby ameliorating the
2 effects of EAE. We speculate that these effects may be result of inflammatory
3 cytokine secretion by CD4⁺ T-cells. Effector cytokines from CD4⁺ T-cells may also
4 act to enhance CD8 cytotoxic functions, thereby amplifying inflammatory cascades in
5 the CNS. However, further studies are required to determine whether the role of
6 CD8⁺ T-cells in MS is primarily pathological or regulatory.
7

8 In the present study, we evaluated levels of various cytokines (granzyme B, perforin,
9 IL-17 and IFN- γ) in co-cultures of neurons and either CD8⁺ or CD4⁺ T-cells. We
10 observed that increased levels of perforin resulted in increased neuronal cell death,
11 suggesting that perforin secretion by CD8⁺ T-cells may play an important role in
12 neuronal cell death. Previous studies have indicated that IFN- γ reduces disease
13 severity by mediating MBP-specific CD8⁺ T-cell-driven activity. Recent work has
14 suggested that IL-10-producing CD8⁺ T-cells also act to diminish disease
15 pathology[20, 21]. IFN- γ - and IL-17-producing CD8⁺ T-cells can be recruited into
16 the CNS when responding to apoptotic T-cell associated self-epitopes[22]. In the
17 present study, we observed that increased levels of these cytokines were not
18 associated with cytotoxic effects on neurons, further supporting the notion that they
19 may act to neutralization cytotoxicity. Furthermore, previous studies have reported
20 that CD8⁺ IL-17-secreting T-cell numbers are significantly elevated in acute CNS
21 lesions in patients with MS[23]. Giovanni et al. reported that patients with MS exhibit
22 higher levels of circulating CD8⁺CD56-perforin⁺ T-cells than controls, suggesting
23 that dysregulation of CD8⁺CD56-perforin⁺ T-cells may play a role in the
24 development of MS[24]. Consistent with these findings, perforin was associated with
25 increased neuronal cell death in the present study, suggesting an important role for
26 perforin secretion from CD8⁺ cells in MS.
27

28 In conclusion, our findings indicate that perforin-secreting CD8⁺ T-cells play a major
29 role in MS, and that CD4 may enhance CD8 cytotoxicity to neurons by inducing
30 inflammation. However, the precise roles of CD4 and CD8 in the pathophysiology of
31 MS remain to be elucidated.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

1. Bellavista E, Santoro A, Galimberti D, Comi C, Luciani F, Mishto M (2014) Current understanding on the role of standard and immunoproteasomes in inflammatory/immunological pathways of multiple sclerosis. *Autoimmune diseases* 2014:739705
2. Popescu BF, Lucchinetti CF (2012) Pathology of demyelinating diseases. *Annual review of pathology* 7:185-217
3. Booss J, Esiri MM, Tourtellotte WW, Mason DY (1983) Immunohistological analysis of T lymphocyte subsets in the central nervous system in chronic progressive multiple sclerosis. *Journal of the neurological sciences* 62:219-232
4. Babbe H, Roers A, Waisman A, Lassmann H, Goebels N, Hohlfeld R, Friese M, Schroder R, Deckert M, Schmidt S, Ravid R, Rajewsky K (2000) Clonal expansions of CD8(+) T cells dominate the T cell infiltrate in active multiple sclerosis lesions as shown by micromanipulation and single cell polymerase chain reaction. *The Journal of experimental medicine* 192:393-404
5. Jacobsen M, Cepok S, Quak E, Happel M, Gaber R, Ziegler A, Schock S, Oertel WH, Sommer N, Hemmer B (2002) Oligoclonal expansion of memory CD8+ T cells in cerebrospinal fluid from multiple sclerosis patients. *Brain : a journal of neurology* 125:538-550
6. Junker A, Ivanidze J, Malotka J, Eiglmeier I, Lassmann H, Wekerle H, Meinl E, Hohlfeld R, Dornmair K (2007) Multiple sclerosis: T-cell receptor expression in distinct brain regions. *Brain : a journal of neurology* 130:2789-2799
7. Huseby ES, Liggitt D, Brabb T, Schnabel B, Ohlen C, Goverman J (2001) A pathogenic role for myelin-specific CD8(+) T cells in a model for multiple sclerosis. *The Journal of experimental medicine* 194:669-676
8. Baughman EJ, Mendoza JP, Ortega SB, Ayers CL, Greenberg BM, Frohman EM, Karandikar NJ (2011) Neuroantigen-specific CD8+ regulatory T-cell function is deficient during acute exacerbation of multiple sclerosis. *Journal of autoimmunity* 36:115-124
9. Crawford MP, Yan SX, Ortega SB, Mehta RS, Hewitt RE, Price DA, Stastny P, Douek DC, Koup RA, Racke MK, Karandikar NJ (2004) High prevalence of autoreactive, neuroantigen-specific CD8+ T cells in multiple sclerosis revealed by novel flow cytometric assay. *Blood* 103:4222-4231
10. Karandikar NJ, Crawford MP, Yan X, Ratts RB, Brenchley JM, Ambrozak DR, Lovett-Racke AE, Frohman EM, Stastny P, Douek DC, Koup RA, Racke MK (2002) Glatiramer acetate (Copaxone) therapy induces CD8(+) T cell responses in patients with multiple sclerosis. *The Journal of clinical investigation* 109:641-649
11. Tennakoon DK, Mehta RS, Ortega SB, Bhoj V, Racke MK, Karandikar NJ (2006) Therapeutic induction of regulatory, cytotoxic CD8+ T cells in multiple sclerosis. *Journal of immunology* 176:7119-7129
12. Mott RT, Ait-Ghezala G, Town T, Mori T, Vendrame M, Zeng J, Ehrhart J, Mullan M, Tan J (2004) Neuronal expression of CD22: novel mechanism for inhibiting microglial proinflammatory cytokine production. *Glia* 46:369-379

13. Elong Ngono A, Pettre S, Salou M, Bahbouhi B, Soullou JP, Brouard S, Laplaud DA (2012) Frequency of circulating autoreactive T cells committed to myelin determinants in relapsing-remitting multiple sclerosis patients. *Clinical immunology* 144:117-126
14. Zhang QY, Huang JH, Li HZ, Guo HT, Zhong YQ, Wang YM, Pei JM (2008) Myelin-basic protein-reactive specific CD4+ and CD8+ NK lymphocytes induce morphological changes in neuronal cell bodies and myelin sheaths: implications for multiple sclerosis. *Archives of medical research* 39:45-51
15. Takenaka A, Kaneko K, Mori R, Itoyama Y (1984) Adoptive transfer of experimental allergic encephalomyelitis with lectin-activated spleen cells. *Journal of the neurological sciences* 66:255-261
16. Waldmann H (1989) Manipulation of T-cell responses with monoclonal antibodies. *Annual review of immunology* 7:407-444
17. Zaguia F, Saikali P, Ludwin S, Newcombe J, Beauseigle D, McCrea E, Duquette P, Prat A, Antel JP, Arbour N (2013) Cytotoxic NKG2C+ CD4 T cells target oligodendrocytes in multiple sclerosis. *Journal of immunology* 190:2510-2518
18. Jiang H, Braunstein NS, Yu B, Winchester R, Chess L (2001) CD8+ T cells control the TH phenotype of MBP-reactive CD4+ T cells in EAE mice. *Proceedings of the National Academy of Sciences of the United States of America* 98:6301-6306
19. Tyler AF, Mendoza JP, Firan M, Karandikar NJ (2013) CD8(+) T Cells Are Required For Glatiramer Acetate Therapy in Autoimmune Demyelinating Disease. *PloS one* 8:e66772
20. Ji Q, Percheret A, Goverman JM (2010) Viral infection triggers central nervous system autoimmunity via activation of CD8+ T cells expressing dual TCRs. *Nature immunology* 11:628-634
21. Trandem K, Zhao J, Fleming E, Perlman S (2011) Highly activated cytotoxic CD8 T cells express protective IL-10 at the peak of coronavirus-induced encephalitis. *Journal of immunology* 186:3642-3652
22. Lolli F, Martini H, Citro A, Franceschini D, Portaccio E, Amato MP, Mechelli R, Annibali V, Sidney J, Sette A, Salvetti M, Barnaba V (2013) Increased CD8+ T cell responses to apoptotic T cell-associated antigens in multiple sclerosis. *Journal of neuroinflammation* 10:94
23. Tzartos JS, Friese MA, Craner MJ, Palace J, Newcombe J, Esiri MM, Fugger L (2008) Interleukin-17 production in central nervous system-infiltrating T cells and glial cells is associated with active disease in multiple sclerosis. *The American journal of pathology* 172:146-155
24. Giovanni F, Domenico P, Alessandro M, Raffaele I, Viviana N, Katia PA, Paola BA (2011) Circulating CD8+CD56-perforin+ T cells are increased in multiple sclerosis patients. *Journal of neuroimmunology* 240-241:137-141

Figure Legends

Figure 1 Differences in the cytotoxicity of CD4+ and CD8+ T-cells to neurons.

CD4+, CD8+, or both CD4+ and CD8+ T-cells were co-cultured with neurons for 24

1 h. These T-cells were stimulated by MBP or harvested from EAE models. Neuronal
2 viability was assessed via TNUEL staining (Bar = 100 μ m). Data from three
3 independent experiments were expressed as mean \pm SEM and analyzed using a
4 one-way ANOVA (*P < 0.05).
5
6
7
8
9

10
11
12 **Figure 2 Expression of Bax in neurons co-cultured with different T-cell types. (A)**

13 The expression of Bax in neurons co-cultured with different T-cell types for 24h
14 stimulated with MBP. T-cells from control rats were stimulated with 50 μ g/ml MBP
15 on the first day, and another 100 μ g/ml MBP was added on the fourth day. The
16 different T-cells were then co-cultured with neurons. The control condition refers to
17 T-cells stimulated with no related peptide. (B) Relative quantification of (A). Data
18 were expressed as mean \pm SEM from at least three independent experiments and
19 analyzed using a one-way ANOVA (*P < 0.05). (C) The expression of Bax in neurons
20 co-cultured for 24 h with different T-cells from EAE models or stimulated with MBP.
21 (D) Relative quantification of (C). Data were expressed as mean \pm SEM from at least
22 three independent experiments and analyzed using a one-way ANOVA (*P < 0.05).
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 **Figure 3 The expression of various factors secreted by different T-cell types when**
42 **co-cultured with neurons for 24 h.** Different T-cells were co-cultured with neurons
43 for 24 h, and the expression of IL-17 (A), perforin (B), granzyme B (C), and IFN- γ (D)
44 was examined via ELISA. Data were expressed as mean \pm SEM from at least three
45 independent experiments and analyzed using a one-way ANOVA (*P < 0.05).
46
47
48
49
50
51
52
53
54
55

56 **Figure 4 Direct cytotoxic effects of various factors on neurons. (A)** Different
57 factors (1.5 μ g/ml) were added to neurons, and the neuronal viability was examined
58
59
60
61
62
63
64
65

via MTT. Data were expressed as mean \pm SEM from at least three independent experiments and analyzed using a one-way ANOVA (*P < 0.05). (B) Different factors (1.5 μ g/ml) were added to neurons, and the expression of Bax was examined via Western blotting.