

## Bibliography and References Cited

- [1] Mobley, D. L., Chodera, J. D., and Dill, K. A.: On the use of orientational restraints and symmetry corrections in alchemical free energy calculations. *J. Chem. Phys.* 125: 084902, 2006.
- [2] Mobley, D. L., Chodera, J. D., and Dill, K. A.: Confine and release: Obtaining correct free energies in the presence of protein conformational change. *J. Chem. Theor. Comput.* 3: 1231–1235, 2007.
- [3] Shirts, M. R., Mobley, D. L., Chodera, J. D., and Pande, V. S.: Accurate and efficient corrections for missing dispersion interactions in molecular simulations. *J. Phys. Chem. B.* 111: 13052–13063, 2007.
- [4] Chodera, J. D., Singhal, N., Pande, V. S., Dill, K. A., and Swope, W. C.: Automatic discovery of metastable states for the construction of Markov models of macromolecular conformational dynamics. *J. Chem. Phys.* 126: 155101, 2007.
- [5] Michael R. Shirts, D. L. M. and Chodera, J. D.: Alchemical free energy calculations: Ready for prime time? *Annu. Rep. Comput. Chem.* 3: 41–59, 2007.
- [6] Shirts, M. R. and Chodera, J. D.: Statistically optimal analysis of samples from multiple equilibrium states. *J. Chem. Phys.* 129: 124105, 2008.
- [7] Nilmeier, J. P., Crooks, G. E., Minh, D. D. L., and Chodera, J. D.: Nonequilibrium candidate Monte Carlo: A new tool for equilibrium simulation. *Proc. Natl. Acad. Sci. USA.* 108: E1009, 2011.
- [8] Chodera, J. D. and Shirts, M. R.: Replica exchange and expanded ensemble simulations as gibbs sampling: Simple improvements for enhanced mixing. *J. Chem. Phys.* 135: 194110, 2011.
- [9] Chodera, J. D., Mobley, D. L., Shirts, M. R., Dixon, R. W., Branson, K. M., and Pande, V. S.: Free energy methods in drug discovery and design: Progress and challenges. *Curr. Opin. Struct. Biol.* 21: 150, 2011.
- [10] Wang, K., Chodera, J. D., Yang, Y., and Shirts, M. R.: Identifying ligand binding sites and poses using GPU-accelerated Hamiltonian replica exchange molecular dynamics. *J. Comput. Aided Mol. Des.* 27: 989, 2013.
- [11] Alexov, E. and Gunner, M. R.: Incorporating protein conformational flexibility into pH-titration calculations: Results on T4 lysozyme. *Biophys. J.* 74: 2075–2093, 1997.
- [12] Alexov, E. and Gunner, M. R.: A pragmatic approach to structure based calculation of coupled proton and electron transfer in proteins. *Biochim. Biophys. Acta.* 1458: 63–87, 2000.
- [13] Georgescu, R. E., Alexov, E. G., and Gunner, M. R.: Combining conformational flexibility and continuum electrostatics for calculating  $pK_a$ s in proteins. *Biophys. J.* 83: 1731–1748, 2002.
- [14] Song, Y., Mao, J., and Gunner, M. R.: MCCE2: Improving protein  $pK_a$  calculations with extensive side chain rotamer sampling. *J. Comput. Chem.* 30: 2231–2247, 2009.
- [15] Song, Y. and Gunner, M. R.: Using multiconformation continuum electrostatics to compare chloride binding motifs in  $\alpha$ -amylase, human serum albumin, and Omp32. *J. Mol. Biol.* 387: 840–846, 2009.
- [16] Zheng, Z., Dutton, P. L., and Gunner, M. R.: The measured and calculated affinity of methyl and methoxy substituted benzoquinones for the QA site of bacterial reaction centers. *Proteins.* 78: 2638–2654, 2010.
- [17] Warncke, K., Gunner, M. R., Braun, B. S., Gu, L., Yu, C., Bruce, J. M., and Dutton, P. L.: Influence of hydrocarbon tail structure on quinone binding and electron-transfer performance at the QA and QB sites of the photosynthetic reaction center protein. *Biochem.* 33: 7830–7841, 1994.
- [18] Zhang, X. and Gunner, M. R.: Affinity and activity of non-native quinones at the Q(B) site of bacterial photosynthetic reaction centers. *Photosynth. Res.* 120: 181–196, 2013.
- [19] Nielsen, J. E., Gunner, M. R., and Garcia-Moreno, E. B.: The  $pK_a$  cooperative: a collaborative effort to advance structure-based calculations of  $pK_a$  values and electrostatic effects in proteins. *Proteins.* 79: 3249–3259, 2011.
- [20] Stanton, C. L. and Houk, K. N.: Benchmarking  $pK_a$  prediction methods for residues in proteins. *J. Chem. Theory Comput.* 4: 951–966, 2008.
- [21] Szakács, Z., Béni, S., Varga, Z., Örfi, L., Kéri, G., and Noszál, B.: Acid-base profiling of imatinib (Gleevec) and its fragments. *J. Med. Chem.* 48: 249, 2005.
- [22] American Cancer Society, . Cancer Facts & Figures 2014, 2014.
- [23] Zhang, J., Yang, P. L., and Gray, N. S.: Targeting cancer with small molecule kinase inhibitors. *Nat. Rev. Cancer.* 9: 28–39, 2009.
- [24] Deininger, M., Buchdunger, E., and Druker, B. J.: The development of imatinib as a therapeutic agent for chronic myeloid leukemia. *Blood.* 105: 2640–2653, 2005.
- [25] Stegmeier, F., Warmuth, M., Sellers, W. R., and Dorsch, M.: Targeted cancer therapies in the twenty-first century: Lessons from imatinib. *Clin. Pharm. & Therap.* 87: 543–552, 2010.
- [26] Longley, D. B. and Johnston, P. G.: Molecular mechanisms of drug resistance. *J. Pathol.* 205: 275–292, 2005.
- [27] The story of Gleevec. <http://www.innovation.org/index.cfm/StoriesofInnovation/InnovatorStories/>

The\_Story\_of\_Gleevec. Accessed: 2014-05-03.

- [28] Shirts, M. R., Mobley, D. L., and Brown, S. P. Free energy calculations in structure-based drug design. In *Structure Based Drug Design*. Cambridge University Press, 2009.
- [29] Paul, S. M., Mytelka, D. S., Dunwiddie, C. T., Persinger, C. C., Munos, B. H., Lindborg, S. R., and Schacht, A. L.: How to improve R&D productivity: The pharmaceutical industry's grand challenge. *Nature Rev. Drug Discovery*. 9: 203, 2010.
- [30] Schneider, G.: Virtual screening: an endless staircase? *Nature Rev. Drug Discovery*. 9: 273–276, 2010.
- [31] ten Brink, T. and Exner, T. E.:  $pK_a$  based protonation states and microspecies for protein-ligand docking. *J. Comput. Aided Mol. Des.* 24: 935–942, 2010.
- [32] Onufriev, A. V. and Alexov, E.: Protonation and  $pK$  changes in protein-ligand binding. *Quart. Rev. Biophys.* 46: 181–209, 2013.
- [33] Seeliger, M. A., Nagar, B., Frank, F., Cao, X., Henderson, M. N., and Kuriyan, J.: c-Src binds to the cancer drug imatinib with an inactive Abl/c-Kit conformation and a distributed thermodynamic penalty. *Structure*. 15: 299–311, 2007.
- [34] Lin, Y.-L., Meng, Y., Jiang, W., and Roux, B.: Explaining why Gleevec is a specific and potent inhibitor of Abl kinase. *Proc. Natl. Acad. Sci. USA*. 110: 1664–1669, 2013.
- [35] Shan, Y., Seeliger, M. A., Eastwood, M. P., Frank, F., Xu, H., Jensen, M. O., Dror, R. O., Kuriyan, J., and Shaw, D. E.: A conserved protonation-dependent switch controls drug binding in Abl kinase. *Proc. Natl. Acad. Sci. USA*. 106: 139–144, 2009.
- [36] Aleksandrov, A. and Simonson, T.: A molecular mechanics model for imatinib and imatinib:kinase binding. *J. Comput. Chem.* 31: 1550–1560, 2010.
- [37] Aleksandrov, A. and Simonson, T.: Molecular dynamics simulations show that conformational selection governs the binding preferences of imatinib for several tyrosine kinases. *J. Biol. Chem.* 285: 13807–13815, 2010.
- [38] Park, M.-S., Gao, C., and Stern, H. A.: Estimating binding affinities by docking/scoring methods using variable protonation states. *Proteins: Structure, Function, and Bioinformatics*. 79: 304–314, 2010.
- [39] Mongan, J., Case, D. A., and McCammon, J. A.: Constant pH molecular dynamics in a generalized born implicit solvent. *J. Comput. Chem.* 25: 2038–2048, 2004.
- [40] Khandogin, J. and Brooks, C. L.: Constant pH molecular dynamics with proton tautomerism. *Biophys. J.* 89: 141–157, 2005.
- [41] Stern, H. A.: Molecular simulation with variable protonation states at constant pH. *J. Chem. Phys.* 126: 164112, 2007.
- [42] Donnini, S., Tegeler, F., Groenhof, G., and Grubmüller, H.: Constant pH molecular dynamics in explicit solvent with  $\lambda$ -dynamics. *J. Chem. Theor. Comput.* 7: 1962–1978, 2011.
- [43] Goh, G. B., Hulbert, B. S., Zhou, H., and Brooks III, C. L.: Constant pH molecular dynamics of proteins in explicit solvent with proton tautomerism. *Proteins*. 82: 1319–1331, 2014.
- [44] Chen, Y. and Roux, B.: Constant-pH hybrid nonequilibrium molecular dynamics-Monte Carlo simulation method. *J. Chem. Theor. Comput.* 11: 3919–3931, 2015.
- [45] Lee, J., Miller, B. T., and Brooks, B. R.: Computational scheme for pH-dependent binding free energy calculation with explicit solvent. *Protein Sci.* 25: 231–243, 2015.
- [46] Wang, L., Wu, Y., Deng, Y., Kim, B., Pierce, L., Krilov, G., Lupyan, D., Robinson, S., Dahlgren, M. K., Greenwood, J., Romero, D. L., Masse, C., Knight, J. L., Steinbrecher, T., Beuming, T., Damm, W., Harder, E., Sherman, W., Brewer, M., Wester, R., Murcko, M., Frye, L., Farid, R., Lin, T., Mobley, D. L., Jorgensen, W. L., Berne, B. J., Friesner, R. A., and Abel, R.: Accurate and reliable prediction of relative ligand binding potency in prospective drug discovery by way of a modern free-energy calculation protocol and force field. *J. Am. Chem. Soc.* 137: 2695–2703, 2015.
- [47] Dullweber, F., Stubbs, M. T., Musil, D., Stürzebecher, J., and Klebe, G.: Factorising ligand affinity: A combined thermodynamic and crystallographic study of trypsin and thrombin inhibition. *J. Mol. Biol.* 313: 593–614, 2001.
- [48] Stubbs, M. T., Reyda, S., Dullweber, F., Möller, M., Klebe, G., Dorsch, D., Mederski, W. W. K. R., and Wurziger, H.: pH-dependent binding modes observed in trypsin crystals: Lessons for structure-based drug design. *ChemBioChem*. 02-03: 246, 2002.
- [49] Czdrowski, P., Sotriffer, C. A., and Klebe, G.: Protonation changes upon ligand binding to trypsin and thrombin: Structural interpretation based on  $pK_a$  calculations and ITC experiments. *J. Mol. Biol.* 367: 1347–1356, 2007.
- [50] Steuber, H., Czdrowski, P., Sotriffer, C. A., and Klebe, G.: Tracing changes in protonation: A prerequisite to factorize thermodynamic data of inhibitor binding to aldose reductase. *J. Mol. Biol.* 373: 1305–1320, 2007.
- [51] Dashi, D. S., Meng, Y., and Roitberg, A. E.: pH-replica exchange molecular dynamics in proteins using a

- discrete protonation method. *J. Phys. Chem. B.* 116: 8805–8811, 2012.
- [52] Khandogin, J. and Brooks III, C. L.: Constant pH molecular dynamics with proton tautomerism. *Biophys. J.* 89: 141–157, 2005.
- [53] Tembe, B. L. and McCammon, J. A.: Ligand-receptor interactions. *Computers & Chemistry.* 8: 281–283, 1984.
- [54] Michel, J. and Essex, J.: Hit identification and binding mode predictions by rigorous free energy simulations. *J. Med. Chem.* 51: 6654–6664, 2008.
- [55] Michel, J. and Essex, J. W.: Prediction of protein–ligand binding affinity by free energy simulations: assumptions, pitfalls and expectations. *J. Comput. Aided Mol. Des.* 24: 639–658, 2010.
- [56] Gallicchio, E. and Levy, R. M.: Advances in all atom sampling methods for modeling protein–ligand binding affinities. *Curr. Opin. Struct. Biol.* 21: 161–166, 2011.
- [57] Yank: A gpu-accelerated python framework for exploring algorithms for alchemical free energy calculations. <http://getyank.org>.
- [58] Eastman, P., Friedrichs, M., Chodera, J. D., Radmer, R., Bruns, C., Ku, J., Beauchamp, K., Lane, T. J., Wang, L.-P., Shukla, D., Tye, T., Houston, M., Stitch, T., and Klein, C.: OpenMM 4: A reusable, extensible, hardware independent library for high performance molecular simulation. *J. Chem. Theor. Comput.* 9: 461, 2012.
- [59] Openmm: A high performance toolkit for molecular simulation. <http://openmm.org>.
- [60] Minh, D. D. L. and Chodera, J. D.: Estimating equilibrium ensemble averages using multiple time slices from driven nonequilibrium processes: theory and application to free energies, moments, and thermodynamic length in single-molecule pulling experiments. *J. Chem. Phys.* 134: 024111, 2011.
- [61] Minh, D. D. L. and Chodera, J. D.: Optimal estimators and asymptotic variances for nonequilibrium path-ensemble averages. *J. Chem. Phys.* 131(13): 134110, 2009.
- [62] Seeliger, M. A., Young, M., Henderson, M. N., Pellicena, P., King, D. S., Falick, A. M., and Kuriyan, J.: High yield bacterial expression of active c-Abl and c-Src tyrosine kinases. *Protein Sci.* 14: 3135–3139, 2005.
- [63] Shelley, J. C., Cholleti, A., Frye, L. L., Greenwood, J. R., Timlin, M. R., and Uchiyama, M.: Epik: A software program for pKa prediction and protonation state generation for drug-like molecules. *J. Comput. Aided Mol. Des.* 21: 681–691, 2007.
- [64] Milletti, F., Storchi, L., and Cruciani, G.: Predicting protein  $pK_a$  by environment similarity. *Proteins.* 50: 484–495, 2009.
- [65] Jakalian, A., Bush, B. L., Jack, D. B., and Bayly, C. I.: Fast, efficient generation of high-quality atomic charges. AM1-BCC model: I. method. *J. Comput. Chem.* 21(2): 132–146, 2000.
- [66] Jakalian, A., Jack, D. B., and Bayly, C. I.: Fast, efficient generation of high-quality atomic charges. AM1-BCC model: II. parameterization and validation. *J. Comput. Chem.* 23(16): 1623–1641, 2002.
- [67] Qian, B., Raman, S., Das, R., Bradley, P., McCoy, A. J., Read, R. J., and Baker, D.: High-resolution structure prediction and the crystallographic phase problem. *Nature.* 450(7167): 259–264, November 2007.
- [68] Wang, C., Bradley, P., and Baker, D.: Protein–Protein Docking with Backbone Flexibility. *Journal of Molecular Biology.* 373(2): 503–519, October 2007.
- [69] Prinz, J.-H., Wu, H., Sarich, M., Keller, B., Fischbach, M., Held, M., Chodera, J. D., Schütte, C., and Noé, F.: Markov models of molecular kinetics: Generation and validation. *J. Chem. Phys.* 134: 174105, 2011.
- [70] Crooks, G. E.: Measuring thermodynamic length. *Phys. Rev. Lett.* 99(10): 100602, 2007.
- [71] Shenfeld, D. K., Xu, H., Eastwood, M. P., Dror, R. O., and Shaw, D. E.: Minimizing thermodynamic length to select intermediate states for free-energy calculations and replica-exchange simulations. *Phys. Rev. E.* 80(4): 046705, 2009.
- [72] Swope, W. C., Andersen, H. C., Berens, P. H., and Wilson, K. R.: A computer simulation method for the calculation of equilibrium constants for the formation of physical clusters of molecules: Application to small water clusters. *J. Chem. Phys.* 76(1): 637–649, 1982.
- [73] Lyubartsev, A. P., Martsinovskii, A. A., Shevkunov, S. V., and Vorontsov-Velyaminov, P. N.: New approach to Monte Carlo calculation of the free energy: Method of expanded ensembles. *J. Chem. Phys.* 96: 1776–1783, 1992.
- [74] Tan, Z.: Optimally adjusted mixture sampling and locally weighted histogram analysis. *J. Comput. Graph. Stat.* in press.
- [75] Bain, J., Plater, L., Elliott, M., Shpiro, N., Hastie, C. J., McLauchlan, H., Klevernic, I., Arthur, J. S. C., Aless, D. R., and Cohen, P.: The selectivity of protein kinase inhibitors: A further update. *Biochem. J.* 408: 297–315, 2007.
- [76] Fabian, M. A., Biggs, W. H., Treiber, D. K., Atteridge, C. E., Azimioara, M. D., Benedetti, M. G., Carter, T. A., Ciceri, P., Edeen, P. T., Floyd, M., Ford, J. M., Galvin, M., Gerlach, J. L., Grotzfeld, R. M., Herrgard, S., Insko, D. E., Insko, M. A., Lai, A. G., Lélías, J.-M., Mehta, S. A., Milanov, Z. V., Velasco, A. M., Wodicka, L. M.,

- Patel, H. K., Zarrinkar, P. P., and Lockhart, D. J.: A small molecule–kinase interaction map for clinical kinase inhibitors. *Nature Biotech.* 23: 329–336, 2005.
- [77] Karaman, M. W., Herrgard, S., Treiber, D. K., Gallant, P., Atteridge, C. E., Campbell, B. T., Chan, K. W., Ciceri, P., Davis, M. I., Edeen, P. T., Faraoni, R., Floyd, M., Hunt, J. P., Lockhart, D. J., Milanov, Z. V., Morrison, M. J., Pallares, G., Patel, H. K., Pritchard, S., Wodicka, L. M., and Zarrinkar, P. P.: A quantitative analysis of kinase inhibitor selectivity. *Nature Biotech.* 26: 127–132, 2008.
- [78] Parton, D. L., Grinaway, P. B., Hanson, S. M., Beauchamp, K. A., and Chodera, J. D.: Ensembler: Enabling high-throughput molecular simulations at the superfamily scale. *PLoS Computational Biology*. in press.
- [79] We will start with AMBER family protein forcefields ff99sb-ildn-phi [86] and ff99sb-ildn-nmr [87], which have been shown to give excellent agreement with NMR data [88] and the generalized AMBER small molecule forcefield (GAFF) [89]. We will initially use the TIP3P [90] and TIP4P-Ew [91] explicit solvent models.
- [80] Parton, D. L., Hanson, S. M., Rodríguez-Laureano, L., Albanese, S. K., Jeans, C., Gradia, S., Seeliger, M. A., and Chodera, J. D.: A panel of recombinant human kinase domain constructs for automated bacterial expression. *bioRxiv*. <http://dx.doi.org/10.1101/038711> pp, 2016.
- [81] Meyer, B. and Peters, T.: Nmr spectroscopy techniques for screening and identifying ligand binding to protein receptors. *Angew. Chem. Int. Ed. Engl.* 42: 864–890, 2003.
- [82] Lavinder, J. L., Hari, S. B., Sullivan, B. J., and Magliery, T. J.: High-throughput thermal scanning: a general, rapid dye-binding thermal shift screen for protein engineering. *J. Am. Chem. Soc.* 131: 3794–3795, 2009.
- [83] Baker, B. M. and Murphy, K. P.: Evaluation of linked protonation effects in protein binding reactions using isothermal titration calorimetry. *Biophys. J.* 71: 2049–2055, 1996.
- [84] Song, C. W., Griffin, R., and Park, H. J. *Cancer Drug Discovery and Development*, chapter Influence of Tumor pH on Therapeutic Response, pp 21–42. Humana Press, Totowa, New Jersey, 2006.
- [85] Levinson, N. M. and Boxer, S. G.: Structural and spectroscopic analysis of the kinase inhibitor bosutinib and an isomer of bosutinib binding to the Abl tyrosine kinase domain. *PLoS One*. 7: e29828, 2012.
- [86] Nerenberg, P. and Head-Gordon, T.: Optimizing protein-solvent force fields to reproduce intrinsic conformational preferences of model peptides. *J. Chem. Theor. Comput.* 7: 1220–1230, 2011.
- [87] Li, D.-W. and Brüschweiler, R.: NMR-based protein potentials. *Angew. Chemie*. 38: 6930–6932, 2010.
- [88] Beauchamp, K. A., Lin, Y.-S., Das, R., and Pande, V. S.: Are protein force fields getting better? a systematic benchmark on 524 diverse NMR measurements. *J. Chem. Theor. Comput.* 8: 1409–1414, 2012.
- [89] Wang, J., Wolf, R. M., Caldwell, J. W., Kollman, P. A., and Case, D. A.: Development and testing of a general AMBER force field. *J. Comput. Chem.* 25: 1157–1174, 2004.
- [90] Jorgensen, W. L., Chandrasekhar, J., Madura, J. D., Impey, R. W., and Klein, M. L.: Comparison of simple potential functions for simulating liquid water. *J. Chem. Phys.* 79: 926, 1983.
- [91] Horn, H. W., Swope, W. C., Pitara, J. W., Madura, J. D., Dick, T. J., Hura, G. L., and Head-Gordon, T.: Development of an improved four-site water model for biomolecular simulations: TIP4P-Ew. *J. Chem. Phys.* 120: 9665, 2004.