Acceleration model with 4 parameters

$$a(k-1) = -\overline{c}_1 a(k) + \overline{c}_2 \left( v(k+1) - v(k) \right) \tag{1}$$

$$a(k-1) = -\overline{c}_3 a(k) + \overline{c}_4 (s(k+1) - s(k) - dt \ v(k))$$
(2)

Model in matrix notation

$$\begin{bmatrix} a(k-1) \\ a(k-1) \end{bmatrix} = \begin{bmatrix} -a(k) & v(k+1) - v(k) & 0 & 0 \\ 0 & 0 & -a(k) & s(k+1) - s(k) - dt \ v(k) \end{bmatrix} \begin{bmatrix} \overline{c}_1 \\ \overline{c}_2 \\ \overline{c}_3 \\ \overline{c}_4 \end{bmatrix}$$
(3)

Add x and y components

$$\begin{bmatrix} a_x(k-1) \\ a_x(k-1) \\ a_y(k-1) \\ a_y(k-1) \end{bmatrix} = \begin{bmatrix} -a_x(k) & v_x(k+1) - v_x(k) & 0 & 0 \\ 0 & 0 & -a_x(k) & s_x(k+1) - s_x(k) - dt & v_x(k) \\ 0 & -a_y(k) & v_y(k+1) - v_y(k) & 0 & 0 \\ 0 & 0 & -a_y(k) & s_y(k+1) - s_y(k) - dt & v_y(k) \end{bmatrix} \begin{bmatrix} \overline{c}_1 \\ \overline{c}_2 \\ \overline{c}_3 \\ \overline{c}_4 \end{bmatrix}$$
(4)

For easier implementation split into two models

$$\begin{bmatrix} a_x(k-1) \\ a_y(k-1) \end{bmatrix} = \begin{bmatrix} -a_x(k) & v_x(k+1) - v_x(k) \\ -a_y(k) & v_y(k+1) - v_y(k) \end{bmatrix} \begin{bmatrix} \overline{c}_1 \\ \overline{c}_3 \end{bmatrix}$$
 (5)

$$\begin{bmatrix} a_x(k-1) \\ a_y(k-1) \end{bmatrix} = \begin{bmatrix} -a_x(k) & s_x(k+1) - s_x(k) - dt \ v_x(k) \\ -a_y(k) & s_y(k+1) - s_y(k) - dt \ v_y(k) \end{bmatrix} \begin{bmatrix} \overline{c}_2 \\ \overline{c}_4 \end{bmatrix}$$
 (6)

Mind dump

$$\Rightarrow$$
 (7)

$$a_x(k-1) = -\bar{c}_1 a_x(k) + \bar{c}_2 \left( v_x(k+1) - v_x(k) \right) \tag{8}$$

$$a_x(k-1) = -\overline{c}_3 a_x(k) + \overline{c}_4 \Big( s_x(k+1) - s_x(k) - dt \ v_x(k) \Big)$$
(9)

$$a_y(k-1) = -\bar{c}_1 a_y(k) + \bar{c}_2 \left( v_y(k+1) - v_y(k) \right) \tag{10}$$

$$a_y(k-1) = -\overline{c}_3 a_y(k) + \overline{c}_4 \Big( s_y(k+1) - s_y(k) - dt \ v_y(k) \Big)$$
(11)

Extended forumal with added features (7 parameters)

$$v(k+1) = v(k) + c_0 v(k-1) + c_1 a(k) + c_2 a(k-1)$$
(12)

$$s(k+1) = s(k) + c_3 s(k-1) + dt \ v(k) + c_4 v(k-1) + c_5 a(k) + c_6 a(k-1)$$

$$(13)$$

(14)

$$c_2a(k-1) = v(k+1) - v(k) - c_0v(k-1) - c_1a(k)$$
(15)

$$c_6a(k-1) = s(k+1) - s(k) - c_3s(k-1) - dt \ v(k) - c_4v(k-1) - c_5a(k)$$

$$(16)$$

$$a(k-1) = \frac{v(k+1) - v(k)}{c_2} - \frac{c_0}{c_2}v(k-1) - \frac{c_1}{c_2}a(k)$$
(18)

$$a(k-1) = \frac{s(k+1) - s(k)}{c_6} - \frac{c_3 s(k-1)}{c_6} - \frac{dt}{c_6} v(k) - \frac{c_4}{c_6} v(k-1) - \frac{c_5}{c_6} a(k)$$
(19)

(20)

Imagine all coefficient have dashes

$$a(k-1) = c_2(v(k+1) - v(k)) - c_0v(k-1) - c_1a(k)$$
(21)

$$a(k-1) = c_6(s(k+1) - s(k) - dt \ v(k)) - c_3s(k-1) - c_4v(k-1) - c_5a(k)$$
(22)

(23)

Equation for the model

$$a(k-1) = -c_0 v(k-1) - c_1 a(k) + c_2 (v(k+1) - v(k))$$
(24)

$$a(k-1) = -c_3 s(k-1) - c_4 v(k-1) - c_5 a(k) + c_6 (s(k+1) - s(k) - dt \ v(k))$$
(25)

(26)

Here we solve the following system:

$$\begin{bmatrix} a(k-1) \\ a(k-1) \end{bmatrix} = \begin{bmatrix} v(k-1) & -a(k) & v(k+1) - v(k) & 0 & 0 & 0 & 0 \\ 0 & 0 & -s(k-1) & -v(k-1) & -a(k) & +(s(k+1) - s(k) - dt \ v(k)) \end{bmatrix} \begin{vmatrix} c_0 \\ \overline{c}_1 \\ \overline{c}_2 \\ \overline{c}_3 \\ \overline{c}_4 \\ \overline{c}_5 \\ \overline{c}_6 \end{vmatrix}$$
(27)

We add the x and y components (imagine the matrix also contains the indices)

$$\begin{bmatrix} a_x(k-1) \\ a_x(k-1) \\ a_y(k-1) \\ a_y(k-1) \end{bmatrix} = \begin{bmatrix} v(k-1) & -a(k) & v(k+1) - v(k) & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -s(k-1) & -v(k-1) & -a(k) & +(s(k+1) - s(k) - dt & v(k)) \\ v(k-1) & -a(k) & v(k+1) - v(k) & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -s(k-1) & -v(k-1) & -a(k) & +(s(k+1) - s(k) - dt & v(k)) \end{bmatrix} \begin{bmatrix} c_0 \\ \overline{c}_1 \\ \overline{c}_2 \\ \overline{c}_3 \\ \overline{c}_4 \\ \overline{c}_5 \\ \overline{c}_6 \end{bmatrix}$$
(28)

We split the model into two models for easier implementation

$$\begin{bmatrix} a_x(k-1) \\ a_y(k-1) \end{bmatrix} = \begin{bmatrix} v_x(k-1) & -a_x(k) & v_x(k+1) - v_x(k) \\ v_y(k-1) & -a_y(k) & v_y(k+1) - v_y(k) \end{bmatrix} \begin{bmatrix} \overline{c}_0 \\ \overline{c}_1 \\ \overline{c}_2 \end{bmatrix}$$
(29)

$$\begin{bmatrix} a_x(k-1) \\ a_y(k-1) \end{bmatrix} = \begin{bmatrix} -s_x(k-1) & -v_x(k-1) & -a_x(k) & s_x(k+1) - s_x(k) - dt & v_x(k) \\ -s_y(k-1) & -v_y(k-1) & -a_y(k) & s_y(k+1) - s_y(k) - dt & v_y(k) \end{bmatrix} \begin{bmatrix} \overline{c}_3 \\ \overline{c}_4 \\ \overline{c}_5 \\ \overline{c}_6 \end{bmatrix}$$
(30)