

The Kuramoto-Sivashinsky Equation

Anders, Elisabeth og Espen

April 1, 2014

Abstract

This is the abstract. Write smart things here.

Introduction

The Kuramoto-Sivashinsky equation,

$$u_t + u_{xx} + u_{xxxx} + uu_x = 0 \quad (1)$$

is one of the simplest partial differential equations that exhibits complicated dynamics in both time and space, which is why the equation has been the attention for a lot of research. The equation was developed by two scientists at the same time in 1977 [1]. Gregory Sivashinsky determined an equation for a laminar flame front, while Yoshiki Kuramoto modeled a diffusion-induced chaos using the same equation. Because of this, the equation is named Kuramoto-Sivashinsky. The KS-equation also models the motion of a fluid going down a vertical wall, e.g. solitary pulses in a falling thin film. [2]

The reason for the complex behaviour comes from the second- and fourth-order derivatives in (1). While the second-order term acts as an energy source and has a destabilizing effect, the fourth-order term has a stabilizing effect. In addition to this, the nonlinear term transfers energy from low to high wave numbers. [3] The KS-equation is a stiff equation, i.e. an equation where numerical methods for solving it are numerically unstable, unless the step size is extremely small. u_{xxxx} is the main reason for this as it leads to rapid variation in the solution.

Numerical results

Initial conditions

In the solution of the KS-equation we had periodic boundary conditions, i.e. $u(0, t) = u(L, t)$. We also used L-periodic initial conditions. We experienced that a common initial condition used in several other reports was

$$u(x, 0) = \cos\left(\frac{x}{16}\right)\left(1 + \sin\left(\frac{x}{16}\right)\right). \quad (2)$$

We also tried the initial condition

$$u(x, 0) = \frac{1}{\sqrt{2}} \sin(x) - \frac{1}{8} \sin(2x), \quad (3)$$

which worked well. The L-periodic initial conditions is customarily taken [4] to satisfy

$$\int_0^L f(x) dx = 0, \quad (4)$$

which both of our initial conditions satisfy. The same article also states that for L-periodic initial data, a unique solution for (1) exists, and is bounded as $t \rightarrow \infty$. The bound has been proven to be smaller than $O(L^{8/5})$. In our numerical tests, with $t = 5000$, the initial condition (2) did indeed not exceed the bound, nor did (3).

References

- [1] Scott Arthur Gasner, Fall 2004, *Integrating the Kuramoto-Sivashinsky equation: A simulation of the hopping state*
http://terminus.sdsu.edu/thesis_repository/ScottGasner_2004_Fall_MS_Comp_Sci.pdf,
 03/31-2014
- [2] Mehrdad Lakestani and Mehdi Dehghan, February 2012, *Numerical solutions of the generalized Kuramoto-Sivashinsky equation using B-spline functions*,
<http://www.sciencedirect.com/science/article/pii/S0307904X11004082>,
 03/31-2014
- [3] Marjan Uddin and Sardar Ali, January 2013, *RBF-PS method and Fourier Pseudospectral method for solving stiff nonlinear partial differential equations*,
<http://www.naturalspublishing.com/files/published/ed38fj6n3xt187.pdf>,
 03/31-2014
- [4] Andrew Spratley, March 2010, *Kuramoto-Sivashinsky equation: A PDE with chaotic solutions*, <http://www.dtic.mil/dtic/tr/fulltext/u2/a228590.pdf>,
 03/31-2014