
Stage-1 markers (n = 14): quantified 5 (1 with isomers),
p < 0.05 in the same direction 1, q < 0.05 (adjusted) 1
Val-Leu (231.1701, 4.15) [1/2, 3.92 min]
Val-Leu (231.1701, 4.15) [2/2, 3.72 min]
LysoPE(20:4/0:0) (502.2934, 15.81)
Leu-Thr-Ser (320.1823, 2.68) [1/2, 2.87 min]
Glutaminylproline (244.1295, 2.58)
5-Hydroxyomeprazole (362.1248, 8.79)
Leu-Thr-Ser (320.1823, 2.68) [2/2, 2.37 min]
Leu-Leu-Arg (401.2877, 7.90)
2-Iminobiotin (244.1115, 5.74)
Phe-Ile-Arg (435.2716, 8.11)
N-Acetyltryptophan (249.0923, 6.34)
Leu-Leu-Tyr (neurotensin 11-13) (408.2497, 9.23)
Inosine isomer (269.0880, 2.01)
Glu-Ile-Val (360.2132, 2.57)
2-Amino-3-{[2-(pyridin-4-yl)ethyl]sulfanyl}propanoic acid (249.0661, 12.45)
(3S)-3-[(2S)-2-Amino-3-hydroxybutyl]pyrrolidin-2-one (173.1286, 7.41)
AUC 0.49, p 0.91, q 0.95
AUC 0.76, p 0.00014, q 0.048
AUC 0.59, p 0.2, q 0.43
AUC 0.35, p 0.035, q 0.55
AUC 0.60, p 0.17, q 0.74
AUC 0.51, p 0.84, q 0.93
Detected, < detection filter
Detected, < detection filter
Detected, < detection filter
Not detected
Not detected
Not detected
Not detected
Not detected
Not detected
Not detected
same direction, p < 0.05
not significant
−1.0
−0.5
0.0
0.5
1.0
1.5
2.0
log2 fold change BC/Control in the independent cohort (unadjusted)