

Medical Isotope Production at Berkeley

Andrew S. Voyles

19 May 2017

2017 Nuclear Physics Summer School in Oslo

Overview

- Berkeley is currently leading a targeted experimental campaign to address these needs:
 - (n,p) production cross sections – UCB
 - Stacked-target charged particle excitation functions – LBNL / LANL
 - Tunable quasi-monoenergetic neutron source – LBNL

^{64}Cu and ^{47}Sc (n,p) Cross-Section Measurements for Medical Radionuclide Production

UCB

Some perspective

Nuclear Data Needs and Capabilities for Applications

May 27-29, 2015

Lawrence Berkeley National Laboratory,
Berkeley, CA USA



Some perspective

Isotope Production Needs

1. Charged-particle reactions for the need

- Charged-particle reactions for the production of medical isotopes at low energies ($E < 30$ MeV):
 - $^{45}\text{Sc}(\text{p},\text{n})^{45}\text{Ti}$; $^{52}\text{Cr}(\text{p},\text{n})^{52}\text{Mn}$; $^{54}\text{Fe}(\text{d},\text{n})^{55}\text{Co}$; $^{67}\text{Zn}(\text{p},\alpha)^{64}\text{Cu}$; $^{72}\text{Ge}(\text{p},\text{n})^{72}\text{As}$, $^{74}\text{Se}(\text{d},\text{n})^{75}\text{Br}$;
 - $^{86}\text{Sr}(\text{p},\text{n})^{86}\text{Y}$; $^{120}\text{Te}(\text{p},\text{n})^{120}\text{I}$

100

- $^{59}\text{Co}(\text{p},3\text{n})^{57}\text{Ni}$, $^{75}\text{As}(\text{p},3\text{n})^{75}\text{Se}$, $^{82}\text{Br}(\text{p},\text{x})^{82}\text{Se}$, $^{119}\text{In}(\text{p},\text{x})^{119}\text{Sn}$, ^{122}Tl
- $^{121}\text{Sb}(\text{p},3\text{n})^{119}\text{Te}/^{119}\text{Sb}$, $^{133}\text{Cs}(\text{p},3\text{n})^{133}\text{Ba}$
- $^{55}\text{Mn}(\text{p},4\text{n})^{52}\text{Fe}$, $^{71}\text{Ga}(\text{p},4\text{n})^{68}\text{Ge}$, $^{133}\text{Cs}(\text{p},5\text{n})^{128}\text{Ba}$
- $^{127}\text{I}(\text{p},6\text{n})^{122}\text{Xe}$
- $^{119}\text{Br}(\text{p},\text{x})^{72}\text{Se}$, $^{119}\text{In}(\text{p},\text{x})^{119}\text{Sn}$, ^{122}Tl
- $^{121}\text{Sb}(\text{p},\text{xn})^{119}\text{Te}/^{119}\text{Sb}$, $^{133}\text{Cs}(\text{p},\text{xn})^{133}\text{Ba}$
- $^{68}\text{Zn}(\text{p},\text{xn})^{64}\text{Cu}$
- $^{68}\text{Zn}(\text{p},2\text{p})^{67}\text{Cu}$, $^{124}\text{Xe}(\text{p},2\text{p})^{123}\text{I}$
- $^{127}\text{Xe}(\text{p},\text{pn})^{123}\text{Xe}$
- (p,x) reaction on $^{94-98}\text{Mo}$ for imp
- $^{107}\text{Ag}(\text{p},\text{xn})^{103}\text{Pd}$
- $^{116}\text{Cd}(\text{a},3\text{n})^{117m}\text{Sn}$; $^{192}\text{Os}(\text{a},3\text{n})^{193}$

3. Nuclear data needed for radionuclides | fission neutrons:

- $^{36}\text{S}(\text{n},\text{x})^{32}\text{Si}$
- $^{37}\text{Cl}(\text{n},\text{x})^{32}\text{Si}$, $^{37}\text{Cl}(\text{n},\text{x})^{32}\text{Si}$
- $^{62}\text{Zn}(\text{n},\text{x})^{67}\text{Cu}$, $^{62}\text{Zn}(\text{n},\text{x})^{67}\text{Cu}$, ^{70}Z
- $^{229}\text{Ra}(\text{n},2\text{n})^{225}\text{Ra}$
- $^{232}\text{Th}(\text{n},\text{x})^{225}\text{Ac}$, $^{232}\text{Th}(\text{n},\text{x})^{227}\text{Ac}$
- $^{32}\text{S}(\text{n},\text{p})^{32}\text{P}$; $^{47}\text{Ti}(\text{n},\text{p})^{47}\text{Ca}$, $^{62}\text{Zn}(\text{n},\text{p})^{64}\text{Cu}$, $^{149}\text{Sm}(\text{n},\text{p})^{149}\text{Pr}$, $^{153}\text{Eu}(\text{n},\text{p})^{153}\text{Sr}$, $^{162}\text{Tm}(\text{n},\text{p})^{162}\text{Er}$, $^{175}\text{Lu}(\text{n},\text{p})^{175}\text{Yb}$, $^{177}\text{Lu}(\text{n},\text{p})^{175}\text{Yb}$, $^{177}\text{Lu}(\text{n},\text{p})^{175}\text{Yb}$
- $^{68}\text{Zn}(\text{y},\text{p})^{67}\text{Cu}$; $^{100}\text{Mo}(\text{y},\text{n})^{99}\text{Mo}$; 10

4. High-energy photon-induced reactions

- $^{68}\text{Zn}(\text{y},\text{p})^{67}\text{Cu}$; $^{100}\text{Mo}(\text{y},\text{n})^{99}\text{Mo}$; 10

5. Nuclear data

- $^{100}\text{Mo}(\text{d},\text{p}2\text{n})$
- $^{100}\text{Mo}(\text{n},2\text{n})$
- $^{100}\text{Mo}(\text{p},\text{pn})$ - data on long-lived

4. Need small uncertainties on all dosimetry reactions

In standard benchmark neutron field
 $^{31}\text{P}(\text{n},\text{p})$, $^{10}\text{B}(\text{n},\text{X})\text{a}$, $^{54}\text{Fe}(\text{n},\text{a})$, $^{23}\text{Na}(\text{n},2\text{n})$, $^{186}\text{W}(\text{n},\text{y})$, $^{56}\text{Mn}(\text{n},\text{y})$, and $^{62}\text{Zn}(\text{n},\text{p})$
 ary
 $^{113}\text{Cd}(\text{n},\text{f})$ and $^{238}\text{U}(\text{n},2\text{n})$

1. Charged-particle reactions for the production of medical isotopes at low energies ($E < 30$ MeV):

- $^{45}\text{Sc}(\text{p},\text{n})^{45}\text{Ti}$; $^{52}\text{Cr}(\text{p},\text{n})^{52}\text{Mn}$; $^{54}\text{Fe}(\text{d},\text{n})^{55}\text{Co}$; $^{67}\text{Zn}(\text{p},\alpha)^{64}\text{Cu}$; $^{72}\text{Ge}(\text{p},\text{n})^{72}\text{As}$, $^{74}\text{Se}(\text{d},\text{n})^{75}\text{Br}$;
- $^{86}\text{Sr}(\text{p},\text{n})^{86}\text{Y}$; $^{120}\text{Te}(\text{p},\text{n})^{120}\text{I}$

o $\text{Cr}(\text{n},\text{X})$ - resonance near 1 keV in particular:

- Nuclear data needed for Super Heavy Element (SHE) target
 - $^{248}\text{Cm}(\text{n},\text{y})$ low energy resonances
 - $^{249}\text{Bk}(\text{n},\text{y})$

$^{32}\text{S}(\text{n},\text{p})^{32}\text{P}$, $^{47}\text{Ti}(\text{n},\text{p})^{47}\text{Ca}$, $^{64}\text{Zn}(\text{n},\text{p})^{64}\text{Cu}$; $^{67}\text{Zn}(\text{n},\text{p})^{67}\text{Cu}$; $^{89}\text{Y}(\text{n},\text{p})^{89}\text{Sr}$, $^{105}\text{Pd}(\text{n},\text{p})^{105}\text{Rh}$; $^{149}\text{Sm}(\text{n},\text{p})^{149}\text{Pm}$, $^{153}\text{Eu}(\text{n},\text{p})^{153}\text{Sm}$, $^{158}\text{Tb}(\text{n},\text{p})^{158}\text{Gd}$; $^{161}\text{Dy}(\text{n},\text{p})^{161}\text{Tb}$; $^{166}\text{Er}(\text{n},\text{p})^{166}\text{Ho}$; $^{169}\text{Tm}(\text{n},\text{p})^{169}\text{Er}$; $^{175}\text{Lu}(\text{n},\text{p})^{175}\text{Yb}$; $^{177}\text{Hf}(\text{n},\text{p})^{177}\text{Lu}$

^{67}Tc and $^{94}\text{Mo}(\text{a},\text{X})^{97}\text{Ru}$

Gamma yield spectrum for incident neutron energies in the 1 heat generation rates:

- ^{131}I ($\text{Te}(\text{n},\text{y})$ (production of ^{131}I))
- $^{100}\text{Mo}(\text{n},\text{y})$
- $^{103}\text{Ru}(\text{n},\text{y})$
- $^{103}\text{Rh}(\text{n},\text{y})$
- $^{154}\text{Eu}(\text{n},\text{y})$
- $^{155}\text{Eu}(\text{n},\text{y})$
- $^{141}\text{Ce}(\text{n},\text{y})$
- $^{140}\text{Ce}(\text{n},\text{y})$
- $^{89}\text{Sr}(\text{n},\text{y})$

Note: * indicates stable isotopes

Excitation function for incident neutron energies from therm

- $^{177}\text{Lu}(\text{n},\text{y})$, $^{154}\text{Eu}(\text{n},\text{y})$, $^{156}\text{Eu}(\text{n},\text{y})$, $^{77}\text{As}(\text{n},\text{y})$, $^{72}\text{As}(\text{n},\text{y})$

2. Validation data in 30 keV MAC

- $^{238}\text{U}(\text{n},\text{y})$

6. IRMM Exploratory Study of Validation Data in ^{252}Cf Standard Neutron Benchmark Field

- Issues with existing $^{197}\text{Au}(\text{n},\text{y})$ due to room return
- Issues with existing $^{90}\text{Zr}(\text{n},2\text{n})$ due to Th contamination
- Issue with existing $^{96}\text{Zr}(\text{n},2\text{n})$ due to $^{94}\text{Zr}(\text{n},\text{y})$ contribution

Inertial Confinement Fusion Data Needs

- Accurate, temperature-dependent fusion reactivity for light ions is of primary importance to describe thermonuclear burn.
- $d(\text{t},\text{a}), t(\text{t},\text{a}), \text{a}(\text{t},\text{a}), d(\text{d},\text{t}), p, d(\text{d},^3\text{He}), n, d(^3\text{He}, \text{a})$

4. Need small uncertainties on all dosimetry reactions

Xe dopants to probe ablation front instabilities.

$\text{Br}(\text{d},2\text{n})\text{Kr}$ to probe ablator/cold fuel and ablator/hot core mix.

Alpha particle induced reactions to probe hot core mix: ^6Li , ^9Be , ^{10}B (best one), ^{12}C , ^{14}N , ^{16}O , ^{19}F , ^{20}Ne , ^{23}Na , ^{24}Mg , ^{27}Al .

4. Gamma-ray diagnostics for performance and ablator/fuel instabilities.

- Total yield from d-t fusion γ branching ratio at 17.6 MeV.
- $^{12}\text{C}(\text{n},\text{n}'\gamma)$ 4.4 MeV time-integrated emission provides hydrocarbon areal densities (remaining mass). Cross section at 14 MeV must be accurate.
- Does $^{13}\text{C}(\text{n},\text{n}'\gamma)$ have strong emission near 4 MeV? If not, then a useful mix diagnostic is possible.

5. Solid Radiochemistry Diagnostic (SRC) is currently an NIF diagnostic complementary to $^{12}\text{C}-\gamma$ GRH detection (CH pr).

- Ratio of $^{198}\text{Au}/^{196}\text{Au}$ from the activated hohlraum.
- ($\text{n},\text{y})/(\text{n},2\text{n})$: low energy neutrons/14 MeV neutrons.

Highlight from WTTC 2016: $^{54}\text{Fe}(\text{p},\alpha)^{51}\text{Mn}$

- $^{117}\text{Sn}(\text{n},\text{n}')$, covering energy response 0.3 – 3.0 MeV

- Data to support new evaluations

- $^{23}\text{Na}(\text{n},\text{y})$, discrepant in fast neutron region, > 100 keV
- $^{23}\text{Na}(\text{n},2\text{n})$
- $^{27}\text{Al}(\text{n},2\text{n})$

- Address discrepancies:

10. Gamma Emission Probabilities

- $^{103m}\text{Rh} \rightarrow \text{X-ray emission}$

12. Uncertainty in recoil spectrum

- Recoil spectrum char:
- ^{66}Ga , ^{71}Ga , ^{75}As
- Fe isotopes
- Validate/test use of ca uncertainty component

Cross-Section Systematics

$$R = N_T \sigma(\bar{E}) \phi(\bar{E})$$

$$N_\gamma = N_D \epsilon_\gamma I_\gamma$$

$$= \epsilon_\gamma I_\gamma \frac{N_T \sigma(\bar{E}) \phi(\bar{E})}{\lambda} (1 - e^{-\lambda t_i}) e^{-\lambda t_d} (1 - e^{-\lambda t_c})$$

$$\sigma(\bar{E}) = \frac{N_\gamma \lambda}{N_T \epsilon_\gamma I_\gamma \phi(\bar{E}) (1 - e^{-\lambda t_i}) e^{-\lambda t_d} (1 - e^{-\lambda t_c})}$$

Medical Applications

- Emerging medical radionuclides
 - ^{64}Cu ($t_{1/2} = 12.7$ hr) – 61% β^+ to ^{64}Ni , 39% β^- to ^{64}Zn
 - ^{47}Sc ($t_{1/2} = 3.35$ d) – β^- to ^{47}Ti , with 159-keV γ

Promising Prospects for ^{44}Sc -/ ^{47}Sc -Based Theragnostics: Application of ^{47}Sc for Radionuclide Tumor Therapy in Mice

Cristina Müller¹, Maruta Bunka^{2,3}, Stephanie Haller¹, Ulli Köster⁴, Viola Groehn⁵, Peter Bernhardt^{6,7},
Nicholas van der Meulen², Andreas Türler^{2,3}, and Roger Schibli^{1,8}

¹Center for Radiopharmaceutical Sciences ETH-PSI-USZ, Paul Scherrer Institute, Villigen-PSI, Switzerland; ²Laboratory of Radiochemistry and Environmental Chemistry, Paul Scherrer Institute, Villigen-PSI, Switzerland; ³Laboratory of Radiochemistry and Environmental Chemistry, Department of Chemistry and Biochemistry University of Bern, Bern, Switzerland; ⁴Institut Laue-Langevin, Grenoble, France; ⁵Merck and Cie, Schaffhausen, Switzerland; ⁶Department of Radiation Physics, The Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden; ⁷Department of Medical Physics and Medical Bioengineering, University Hospital, Gothenburg, Sweden; and ⁸Department of Chemistry and Applied Biosciences, ETH Zurich

In Vivo Evaluation of Pretargeted ^{64}Cu for Tumor Imaging and Therapy


Michael R. Lewis, PhD¹; Mu Wang, MD¹; Donald B. Axworthy, BS²; Louis J. Theodore, PhD²; Robert W. Mallet, BS²; Alan R. Fritzberg, PhD²; Michael J. Welch, PhD¹; and Carolyn J. Anderson, PhD¹

¹Mallinckrodt Institute of Radiology, Washington University School of Medicine, St. Louis, Missouri;
and ²NeoRx Corporation, Seattle, Washington


The UC Berkeley High Flux Neutron Generator

Commissioning:
25 July 2015

2.45 MeV neutrons, 10^8 n/s/cm²
Extraction Plate: 5mm diameter





The UC Berkeley High Flux Neutron Generator





Foils Used	Metal Purity	Abundance (a/o)	Foil Density (mg/cm ²)
^{nat} In	> 99.999%	¹¹³ In (4.29%), ¹¹⁵ In (95.71%)	365.5
^{nat} Zn	> 99.99%	⁶⁴ Zn (49.17%)	714.1
^{nat} Ti	99.999%	⁴⁷ Ti (7.44%)	450.6


The UC Berkeley High Flux Neutron Generator




The UC Berkeley High Flux Neutron Generator




Relative Activation Measurements




Relative Activation Measurements




Relative Activation Measurements



What's going on???




Neutron Energy Spread




Neutron Energy Spread

H. Liskien *et al.*, Nucl Data Tables, vol 11, 2973




Neutron flux profile modeled
in target and monitor foil,
using MCNP6





Neutron Energy Spread

H. Liskien *et al.*, Nucl Data Tables, vol 11, 2973




Neutron flux
in target and
using MCNP6




$$F(E') = \frac{\int_0^{E'} \sigma(E) \frac{d\phi}{dE} dE}{\int_0^{E_{\max}} \sigma(E) \frac{d\phi}{dE} dE}$$

E_{\max} = Maximum energy neutron subtended by foil
 $F(E')$ = Fraction of Total Reactions induced up to energy E'

Neutron Energy Spread



Before...



Neutron Energy Spread





Table 3. Results of cross section measurement.

Reaction	$\sigma(E_n = 2.7645 \text{ MeV}) \text{ (mb)}$
$^{64}\text{Zn}(\text{n},\text{p})^{64}\text{Cu}$ (relative to ^{113}In)	$45.953 \pm 3.351,$ $46.493 \pm 2.805,$ 46.9 ± 3.189
$^{64}\text{Zn}(\text{n},\text{p})^{64}\text{Cu}$ (relative to ^{115}In)	$49.716 \pm 3.335,$ $49.011 \pm 2.698,$
$^{47}\text{Ti}(\text{n},\text{p})^{47}\text{Sc}$ (relative to ^{115}In)	$25.901 \pm 1.7089,$

After!



Neutron Energy Spread







Table 3. Results of cross section measurement.

Reaction	$\sigma(E_n = 2.7645 \text{ MeV}) (\text{mb})$
$^{64}\text{Zn}(\text{n},\text{p})^{64}\text{Cu}$ (relative to ^{113}In)	$45.953 \pm 3.351,$ $46.493 \pm 2.805,$ 46.9 ± 3.189
$^{64}\text{Zn}(\text{n},\text{p})^{64}\text{Cu}$ (relative to ^{115}In)	$49.716 \pm 3.335,$ $49.011 \pm 2.698,$
$^{47}\text{Ti}(\text{n},\text{p})^{47}\text{Sc}$ (relative to ^{115}In)	$25.901 \pm 1.7089,$

After!




Future Work



Phase 1 current upgrade complete!

Monoenergetic Beam: 10^{3-4} n/s/cm²



Target	Nuclide	Threshold (keV)	% abundance	Strongest γ -ray branch (keV)	# seen in a 0.5% Ge detector	Rate in a 0.5% detector
32S	32P	985	95%	0	0.00E+00	0.00
47Ti	47Ti	0	7.44%	159.381	1.24E+05	1.44
64Zn	64Cu	0	49.20%	511	5.03E+05	5.82
67Zn	67Cu	0	4.04%	184.577	5.45E+04	0.63
89Y	89Sr	726	100%	908.96	1.25E+02	0.00
105Pd	105Rh	0	22.33%	318.9	1.27E+05	1.47
149Sm	149Pm	291	11.24%	285.94	4.47E+03	0.05
153Eu	153Sm	25	52.19%	103.18	1.36E+05	1.58
159Tb	159Gd	190	100%	363.543	1.58E+05	1.83
161Dy	161Tb	0	18.89%	75.57	2.66E+04	0.31
166Er	166Ho	1079	33.50%	1379.4	4.39E+03	0.05
169Tm	169Er	0	100%	109.8	1.87E+01	0.00
175Lu	175Yb	0	97.40%	396.3	1.88E+05	2.18
177Hf	177Lu	0	18.60%	208.4	3.77E+04	0.44

Stacked-target Charged Particle Excitation Functions

LBNL / LANL

$^{51,52}\text{Mn}$ - Motivation


- Emerging medical radionuclides
 - ^{51}Mn ($t_{1/2} = 46$ min, 97% β^+) – short-lived PET tracer for metabolic studies
 - ^{52}Mn ($t_{1/2} = 5.6$ d, 29% β^+) – long-lived PET tracer for neuron tracking, immune studies

Preparation and *in vivo* characterization of $^{51}\text{MnCl}_2$ as PET tracer of Ca^{2+} channel-mediated transport

Stephen A. Graves¹, Reinier Hernandez¹, Hector F. Valdovinos¹, Paul A. Ellison¹, Jonathan W. Engle^{1*}
Todd E. Barnhart¹, Weibo Cai^{1,2,3}, Robert J. Nickles^{1*}


- Manganese has well-established biochemistry and uptake via DOTA-based chelation



Almost no Fe(p,x) XS measurements exist – can use these to probe spin physics in the $A \approx 50$ region





Methdology

88-Inch Cyclotron







L. Bernstein




Two overlapping stacks:
 $E_p = 55 \rightarrow 21 \text{ MeV}, 25 \rightarrow 11 \text{ MeV}$

- 25 μm -thin ${}^{\text{nat}}\text{Fe}$, ${}^{\text{nat}}\text{Cu}$, ${}^{\text{nat}}\text{Ti}$ foils in 0.1" Al frames




- Dosimetry: IAEA charged particle beam monitor reactions:
 - ${}^{\text{nat}}\text{Ti}(\text{p},\text{x})^{48}\text{V}$
 - ${}^{\text{nat}}\text{Cu}(\text{p},\text{x})^{62,63,65}\text{Zn}$


nds.iaea.org/medical/monitor_reactions.html




0.5% initial beam dispersion




2% initial beam dispersion



0.5% initial beam dispersion




2% initial beam dispersion




$$F(E') = \frac{\int_0^{E'} \sigma(E) \frac{d\phi}{dE} dE}{\int_0^{E_{max}} \sigma(E) \frac{d\phi}{dE} dE}$$


E_{max} = Maximum energy proton subtended by foil
 $F(E')$ = Fraction of Total Reactions induced up to energy E'



^{52m}Mn (2+)/ ^{52g}Mn (6+) vs. Energy for $^{56}\text{Fe}(\text{p},\alpha\text{n})$
TALYS Level Density Models 1-6 (default spin cut-off)



^{52m}Mn (2+)/ ^{52g}Mn (6+) vs. Energy for $^{56}\text{Fe}(\text{p},\alpha\text{n})$
TALYS Level Density Model CT + FG



Results consistent with $R \approx 1$ at high energy.


At low energy, results are ambiguous due to energy straggling.

Measurements @ LANL – Nb(p,x)

- $^{nat}\text{Nb}(p,4n)^{90}\text{Mo}$ is a high-priority objective as a new proton beam dosimetry standard for $E_p \approx 50 - 100$ MeV





Measurements @ LANL – Nb(p,x)




Measurements @ LANL – Nb(p,x)


- LBNL: 5 – 55 MeV / A, LANL: 45 – 100 MeV p⁺
- Complementary measurements explore reaction dynamics in different energy regimes, overlap region of 45-55 MeV builds confidence and consistency in results.



Measurements @ LANL – Nb(p,x)




Measurements @ LANL – Nb(p,x)



Tunable Quasi-Monoenergetic Neutron Source


LBNL

$^7\text{Li}(\text{p},\text{n})$ Neutron Sources



$^7\text{Li}(\text{p},\text{n})$ Neutron Sources


88-Inch Cyclotron



Berkeley
UNIVERSITY OF CALIFORNIA


$^7\text{Li}(\text{p},\text{n})$ Neutron Sources

88-Inch Cyclotron




$^7\text{Li}(\text{p},\text{n})$ Neutron Sources

88-Inch Cyclotron





Berkeley
UNIVERSITY OF CALIFORNIA

$^7\text{Li}(\text{p},\text{n})$ Neutron Sources



$^7\text{Li}(\text{p},\text{n})$ Neutron Sources



Summary

Demonstrated ability to measure $R_{\text{spin-cut}}$ in excitation function studies for emerging medical radioisotopes

- Already completed: ^{64}Cu and ^{47}Sc (n,p), Fe(p,x), Zr(d,x), Nb(p,x)
- Upcoming targets: ^{86}Sr (p,x) ^{86}Y , La(p,x) $^{134,135}\text{Ce}$, ^{177}Hf (n,p) ^{177}Lu
 - ^7Li (p,n) quasi-monoenergetic neutron source development
- Possible future candidates: Access targets previously fielded by β^+ -Oslo in the A \approx 50,90, rare earth regions via (p,xn), (α ,xn)



Tusen takk!