

Manuscript Number:	PBIOLOGY-D-25-00453R2
Full Title:	Optimal cues for transmission investment in malaria parasites
Short Title:	Optimal cues for transmission investment in malaria parasites
Article Type:	Research Article
Keywords:	malaria; transmission investment; phenotypic plasticity; cue perception
Corresponding Author:	Avril Wang Massachusetts Institute of Technology Cambridge, MA UNITED STATES OF AMERICA
First Author:	Avril Wang
Order of Authors:	Avril Wang Megan Ann Greischar, PhD Nicole Mideo, PhD
Abstract:	The timing of investment into reproduction is a key determinant of lifetime reproductive success (fitness). Many organisms exhibit plastic, i.e., environmentally-responsive, investment strategies, raising the questions of what environmental cues trigger responses and why organisms have evolved to respond to those particular cues. For malaria parasites (<i>Plasmodium</i> spp.), investment into the production of specialized transmission stages (versus stages that replicate asexually within the host) is synonymous with reproductive investment and also plastic, responding to host- and parasite-derived factors. Previous theory has identified optimal plastic transmission investment strategies for the rodent malaria parasite, <i>Plasmodium chabaudi</i>, as a function of the time since infection, implicitly assuming that parasites have perfect information about the within-host environment and how it is changing. We extend that theory to ask which cue(s) should parasites use? Put another way, which cue(s) maximize parasite fitness, quantified as host infectiousness during acute infection? Our results show that sensing a parasite-associated cue, e.g., the abundance of infected red blood cells or transmission stages, allows parasites to achieve fitness approaching that of the optimal time-varying strategy, but only when parasites perceive the cue non-linearly, responding more sensitively to changes at low densities. However, no single cue can recreate the best time-varying strategy or allow parasites to adopt terminal investment as the infection ends, a classic expectation for reproductive investment. Sensing two cues—log-transformed infected and uninfected red blood cell abundance—enables parasites to accurately track the progression of the infection, permits terminal investment, and recovers the fitness of the optimal time-varying investment strategy. Importantly, parasites that detect two cues more efficiently exploit hosts, resulting in higher virulence compared with those sensing only one cue. Finally, our results suggest a potential tradeoff between achieving an optimal transmission investment strategy in a given host environment and robustness in the face of environmental or developmental fluctuations.
Suggested Reviewers:	Qixin He, PhD Assistant Professor, Purdue University heqixin@purdue.edu Paola Carrillo-Bustamante, PhD Office Head of the RKI Office for WHO Hub Partnership, Robert Koch Institut carrillo-bustamantep@rki.de Arthur Talman, PhD Postdoctoral Fellow, Wellcome Sanger Institute arthur.talman@sanger.ac.uk
Opposed Reviewers:	

Additional Information:	
Question	Response
Financial Disclosure Enter a financial disclosure statement that describes the sources of funding for the work included in this submission. Review the submission guidelines for detailed requirements. View published research articles from PLOS Biology for specific examples. This statement is required for submission and will appear in the published article if the submission is accepted. Please make sure it is accurate. Funded studies Enter a statement with the following details: • Initials of the authors who received each award • Grant numbers awarded to each author • The full name of each funder • URL of each funder website • Did the sponsors or funders play any role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript? Did you receive funding for this work?	Yes
Please add funding details. as follow-up to "Financial Disclosure Enter a financial disclosure statement that describes the sources of funding for the work included in this submission. Review the submission guidelines for detailed requirements. View published research articles from PLOS Biology for specific examples. This statement is required for submission and will appear in the published article if the submission is accepted. Please make sure it is accurate.	ALW was supported by a NSERC USRA grant (https://www.nserc-crsng.gc.ca/students-etudiants/ug-pc/usra-brpc_eng.asp). MAG was funded through the Cornell University College of Agriculture and Life Sciences (https://cals.cornell.edu/). NM was supported by a NSERC Discovery Grant (RGPIN-2024-04818; https://www.nserc-crsng.gc.ca/index_eng.asp). The funders did not play any role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Funded studies Enter a statement with the following details: • Initials of the authors who received each award • Grant numbers awarded to each author • The full name of each funder • URL of each funder website • Did the sponsors or funders play any role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript?	
Did you receive funding for this work?"	
Please select the country of your main research funder (please select carefully as in some cases this is used in fee calculation). as follow-up to "Financial Disclosure Enter a financial disclosure statement that describes the sources of funding for the work included in this submission. Review the submission guidelines for detailed requirements. View published research articles from PLOS Biology for specific examples. This statement is required for submission and will appear in the published article if the submission is accepted. Please make sure it is accurate. Funded studies Enter a statement with the following details: • Initials of the authors who received each award • Grant numbers awarded to each author • The full name of each funder • URL of each funder website • Did the sponsors or funders play any role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript? Did you receive funding for this work?"	CANADA - CA
Competing Interests	The authors have declared that no competing interests exist.

You are responsible for recognizing and disclosing on behalf of all authors any competing interest that could be perceived to bias their work, acknowledging all financial support and any other relevant financial or non-financial competing interests.

Do any authors of this manuscript have competing interests (as described in the [PLOS Policy on Declaration and Evaluation of Competing Interests](#))?

If yes, please provide details about any and all competing interests in the box below. Your response should begin with this statement: *I have read the journal's policy and the authors of this manuscript have the following competing interests:*

If no authors have any competing interests to declare, please enter this statement in the box: *"The authors have declared that no competing interests exist."*

* typeset

This statement is **required** for submission and **will appear in the published article** if the submission is accepted. Please make sure it is accurate and that any funding sources listed in your Funding Information later in the submission form are also declared in your Financial Disclosure statement.

Ethics Statement

I confirm that I have read and understood PLOS' requirements for an Ethics Statement.

You must include an Ethics Statement in the Methods section of your manuscript if your study involved:

- Human participants
- Human specimens or tissue
- Vertebrate animals or cephalopods
- Vertebrate embryos or tissues
- Field research

Not including a statement when required will delay the review process.

General guidance is provided below.

Please consult the guidelines on human subjects research and animal research for detailed instructions.

Format for specific study types

Human Subject Research (involving human participants and/or tissue)

- Give the name of the institutional review board or ethics committee that approved the study
- Include the approval number and/or a statement indicating approval of this research
- Indicate the form of consent obtained (written/oral) or the reason that consent was not obtained (e.g. the data were analyzed anonymously)

Animal Research (involving vertebrate animals, embryos or tissues)

- Provide the name of the Institutional Animal Care and Use Committee (IACUC) or other relevant ethics board that reviewed the study protocol, and indicate whether they approved this research or granted a formal waiver of ethical approval
- Include an approval number if one was obtained
- If the study involved *non-human primates*, add *additional details* about animal welfare and steps taken to ameliorate suffering

Field Research

Include the following details if this study involves the collection of plant, animal, or other materials from a natural setting:

- Field permit number
- Name of the institution or relevant body that granted permission

Please check the box to confirm your understanding of this policy, then read and agree to one of the statements.

Please respond:
as follow-up to "**Ethics Statement**

You must include an Ethics Statement in the Methods section of your manuscript if your study involved:

- Human participants

My study does not require an ethics statement.

- Human specimens or tissue
- Vertebrate animals or cephalopods
- Vertebrate embryos or tissues
- Field research

Not including a statement when required will delay the review process.

General guidance is provided below.
Please consult the guidelines on human subjects research and animal research for detailed instructions.

Format for specific study types

Human Subject Research (involving human participants and/or tissue)

- Give the name of the institutional review board or ethics committee that approved the study
- Include the approval number and/or a statement indicating approval of this research
- Indicate the form of consent obtained (written/oral) or the reason that consent was not obtained (e.g. the data were analyzed anonymously)

Animal Research (involving vertebrate animals, embryos or tissues)

- Provide the name of the Institutional Animal Care and Use Committee (IACUC) or other relevant ethics board that reviewed the study protocol, and indicate whether they approved this research or granted a formal waiver of ethical approval
- Include an approval number if one was obtained
- If the study involved *non-human primates*, add *additional details* about animal welfare and steps taken to ameliorate suffering

Field Research

Include the following details if this study involves the collection of plant, animal, or other materials from a natural setting:

- Field permit number
- Name of the institution or relevant body that granted permission

Please check the box to confirm your understanding of this policy, then read and agree to one of the statements. "

Data Availability

PLOS journals require authors to make all data underlying the findings described in their manuscript fully available, without restriction and from the time of

Yes - all data are fully available without restriction

publication, with only rare exceptions to address legal and ethical concerns (see the [PLOS Data Policy](#) and [FAQ](#) for further details). When submitting a manuscript, authors must provide a Data Availability Statement that describes where the data underlying their manuscript can be found.

Your answers to the following constitute your statement about data availability and will be included with the article in the event of publication. **Please note that simply stating 'data available on request from the author' is not acceptable. If, however, your data are only available upon request from the author(s), you must answer "No" to the first question below, and explain your exceptional situation in the text box provided.**

Do the authors confirm that all data underlying the findings described in their manuscript are fully available without restriction?

Please describe where your data may be found, writing in full sentences. **Your answers should be entered into the box below and will be published in the form you provide them, if your manuscript is accepted.** If you are copying our sample text below, please ensure you replace any instances of **XXX** with the appropriate details.

- If your data are all contained within the paper and/or Supporting Information files, please state this in your answer below. For example, "All relevant data are within the paper and its Supporting Information files."
- If your data are held or will be held in a public repository, include URLs, accession numbers or DOIs. For example, "All **XXX** files are available from the **XXX** database (accession

All code and associated data files are available on https://github.com/avril-microbes/Project-Plasmodium/tree/main/code_repository

number(s) XXX, XXX)." If this information will only be available after acceptance, please indicate this by ticking the box below.

- If neither of these applies but you are able to provide details of access elsewhere, with or without limitations, please do so in the box below. For example:

"Data are available from the XXX Institutional Data Access / Ethics Committee for researchers who meet the criteria for access to confidential data."

"Data are from the XXX study whose authors may be contacted at XXX."

* typeset

Additional data availability information:



Ecology & Evolutionary Biology UNIVERSITY OF TORONTO

February 9, 2025

Dear *PLOS Biology* Editorial Board,

Please consider the manuscript entitled “*Optimal cues for transmission investment in malaria parasites*” for publication in *PLOS Biology*.

All sexually-reproducing organisms face a trade-off in allocating resources between growth and reproduction. Malaria parasites are no different: a *Plasmodium* parasite infecting a host red blood cell can either replicate asexually to produce more cell-infecting stages ('growing' the infection) or produce a single, sexual stage gametocyte that can be transmitted to mosquito vectors ('reproducing' infection). Importantly, the resolution to this trade-off in malaria parasites has implications for human health since lower investment in transmission means more host-damaging parasites are produced.

Transmission investment is a plastic trait, varying over the course of infections and in response to stressors like drug treatment in ways that are predicted to be beneficial for parasites. While adaptive plasticity is a well-known solution to the challenges of living in a changing environment, there has been skepticism about the fitness benefits of observed parasite plasticity (Kochin *et al.* 2010 PLoS Biol 8:e1000524). Knowledge of parasite environmental sensing mechanisms remains limited, and it is unclear what cues parasites use to alter traits. To date, how selection should shape transmission investment has been mathematically explored by assuming parasites have perfect knowledge of the 'age' of infection. Candidate investment strategies are assumed to be functions of that age, and the fittest strategy is numerically determined under different scenarios. Since malaria parasites are unlikely to be directly tracking infection age, an important open question is: can any environmental cue (or cues) recreate the fittest strategy? If so, what is the shape of the reaction norm that maps that cue to the transmission investment phenotype? Answers to these questions could provide insight on interventions to 'trick' parasites into adopting strategies with clinical or epidemiological benefits.

In this manuscript, we develop a modelling framework that allows parasites to perceive within-host environmental cues, determine the cues malaria parasites *should* use to set transmission investment, explore why parasites sometimes fail to attain optimal strategies, and quantify what this all means for host health. We find that no single cue is sufficient to produce the fittest patterns of transmission investment (e.g., terminal investment as the infection is being cleared), but integrating two cues improves parasite fitness and can recapitulate these patterns. Interestingly, the two cues we identify as being most informative for parasite life history decisions were previously identified as most informative for medical professionals seeking to predict clinical outcomes in malaria patients (Torres *et al.* 2016 PLoS Biol 14:e1002436). We also find that the evolutionary pressure to finely tune transmission investment by integrating two cues instead of one can lead to more virulent parasites.

We believe that this manuscript will be of broad interest to readers of *PLOS Biology* since it applies canonical theory from evolutionary ecology to an important disease system and combines analysis of sophisticated within-host dynamical models with validation using pre-existing experimental data. Our model framework is readily extendable for studying plasticity in other organisms, and our results provide a guide for future empirical work that can feedback with theory to yield a more mechanistic understanding of parasite life history strategies, which are of both academic and applied relevance.

Please do not hesitate to contact us should you require any further information.

Sincerely,

Avril L. Wang, Megan A. Greischar, and Nicole Mideo

July 13, 2025

Dear Dr. Roberts,

Thank you for the opportunity to submit a revised version of our manuscript for consideration in *PLOS Biology*. We appreciate the time spent by the reviewers and editors on our paper and believe their comments have helped us greatly improve it. In particular, we have streamlined the text, reperformed model optimization, removed analyses that were not core to our main aims, and we have tried to make those aims (along with details of the model and analyses) more clear. Below you will find the reviewer's comments (**in bold**) and our responses to them (line numbers refer to the version with tracked changes).

We hope you agree that this version thoroughly addresses the reviewers' reports, and we welcome any further feedback.

Sincerely,
Avril Wang, Megan Greischar, Nicole Mideo

Reviewer #1:

There is an intrinsic trade-off between investment in with-host growth versus transmission in malaria parasites. This theoretical study explores the influence of different scenarios on the evolution of plasticity in malaria parasites. Following previous studies, the fitness of the malaria parasite is expressed as the cumulative number of transmissible stages of the parasite (gametocytes) over the duration Δ of the infection. Previous theory assumed that the parasite could modulate the investment in transmission as a function of the time since the initiation of the infection. But it is unclear if the malaria parasite is able to track time since infection. The present work explores the evolution of parasite plasticity when the parasite is unable to track time but can track other cues like the density of different types of cells. The authors show that when the parasite has access to a single cue, some cues are better than others. Plasticity based on a single cue, however, yields much lower fitness than plasticity based on time. Next, the authors consider that the parasite has access to two different cues. Using two cues increases the fitness of the parasite. Yet, when the authors introduce some variability among hosts, the optimal strategy based on a single cue may outperform the optimal strategy based on a two cues. They interpret this effect as a lower robustness of the plasticity based on two cues (versus a single cue). Finally, the authors discuss the emerging link between plasticity and parasite virulence in their model and in *Plasmodium chabaudi*.

Major comments:

There are many interesting ideas in this study. But the analysis relies exclusively on numerical approaches and some steps in the derivation of the results are unclear to me.

First, I do not understand the way plasticity is modelled. From equation (7) the investment in transmission depends on some quantity α days ago and uses a linear combination of Heaviside step functions. But a plot of this function would help a lot to illustrate the general shape of plasticity as a function of the cue. Is this a continuous function? Can it take any shape? Besides, I do not

understand how the function can react differently to the density of the log(density). If there is a threshold value above which there is a switch to a different strategy (this is what step functions are modelling) using the density or the log(density) should be the same. But I am probably misinterpreting the way plasticity is modelled (a figure explaining what is done would help). Figure 2 seemed to provide this information but I don't understand panel B and what are the parameters that are evolving to modify the shape of the reaction norms.

We would like to thank the reviewer for their thoughtful comments that have strengthened the paper. The suggestion to include a figure explaining how we modeled and optimized the reaction norms is a great one, and we have implemented it in a new panel in Figure 1 (Fig 1C). Through this figure and in the text, we have clarified how plasticity is modeled, e.g., in the results, l. 110-114,

“For each putative cue, we define its associated reaction norm using flexible splines (Fig 1C). Specifically, the shape of the reaction norm is determined by a linear combination of three basis functions comprising a cubic spline. Adjusting the weight of each function (i.e., the model coefficients) changes the shape of the reaction norm.”

See also l. 599-611 and the caption for Figure 1 for additional information on how we have parameterized and optimized the reaction norms.

Our method of using cubic splines ensures that the reaction norm is continuous. Representing a sharper curve with this method, which is necessary for a reaction norm that changes drastically at a certain threshold, either requires transforming the input variable (e.g., log-transformation) or placing internal knots (which are like the internal boundaries of a piecewise function). We have chosen the former option, given the difficulty of “knowing” the appropriate location to place internal knots. We have redrawn Figure 2B, which shows optimal reaction norms for different cues, with the x-axis on a log scale. This redrawing better emphasizes the differences that exist for reaction norms based on raw versus logged cues. (Perhaps as intuited by the reviewer, the reaction norms look essentially identical for RBC density as a cue, since this variable generally spans less than one order of magnitude in simulated and real infections.)

We note that the use of the Heaviside function only serves to constrain the ranges of the “perceived” cue values within the plausible range and does not play a role in shaping the reaction norm within the defined cue range. We have reworded the explanation of the Heaviside function in l. 613-618.

What are the black and pink bars supposed to mean on the plots?

The black and pink bars are rug plots, which represent the cue values that are “traversed” in a simulation when the parasite is adopting the optimal transmission strategy. This rug plot indicates the approximate range of cue values relevant to a typical infection. In redrawing Figure 2B, we have made it look less busy, and the rugs are more visible. We have also edited the caption to specifically point to the “horizontal bars” that are the rug plots.

Is there an intrinsic constraint linking $c(t)$ and $c(t+a)$ (for any duration of time a)? If there was, I think it would mean that even though your function assumes that "parasites sense only the magnitude of a cue" (line 473), this function may perceive the shape of the variation of the cue.

Maybe this is why the function that uses the log(density) is performing better? The authors need to be clearer about this key step of their model.

Since c is a function of some other within-host variable (for all cues other than time), there is only an indirect link between $c(t)$ and $c(t+a)$. For example, if the cue is RBC density, then the function c must be continuously differentiable with respect to RBC density – a constraint imposed by the cubic spline. However, transmission investment could vary greatly from one time step to the next if RBC density changed drastically across time steps. The rate of change in cue value is, of course, also constrained (or determined) by our within-host dynamical model, imposing a maximum rate of change. We are not sure that this indirect constraint helps to explain our model or results, and we are confident that the $\log(\text{density})$ performs better for parasite-centric cues because it allows parasites to delay transmission investment right at the start of infections. This difference in performance with logged versus raw cues at low parasite densities is more clear with our updated Figure 2B.

Second, some results are very surprising. Why do you get higher fitness with the best single cue model than with the best dual cue models in figure 3A. If one cue was useless, the parasite should evolve not to use it, and thus the two cue models should always yield fitness $>$ or $+$ to the single cue model. I wonder if the fact you use different time step size (line 664) could explain that. Because of the above two points I am unsure about the results presented in the paper. The difference between the use of the $\log(\text{density})$ is key but I don't understand how the authors got there. I like the idea that heterogeneity among hosts may induce a higher fitness cost if the parasite uses two cues but I don't know if this result is real or if there is a "problem" in the way fitness is computed.

We thank the reviewer for noticing this surprising result, and we agree that the dual cue model should always outperform the single cue model. Despite our global+local optimization approach, this result suggested that our optimization routine for dual cues sometimes failed. To remedy this, we re-ran our dual cue optimizations using another approach (methodology detailed in l. 708-730). Briefly, we first attempted to find a dual cue strategy that resembles the best single cue by essentially ignoring the second cue. We then set this strategy (i.e., the parameters describing the spline surface) as the starting point and used the L-BFGS-B algorithm to perform local optimization. Using this approach to optimization, we were able to increase the fitness of the dual cue models, resulting in all dual cues now having a higher fitness than the single cue models, as expected.

The within-host model is trying to capture many details of the biology of malaria parasites. Do you need all these details to develop your argument? My impression is most of the results you are highlighting should hold in simpler versions of your model, and with a simpler model you may be able to get a deeper understanding of the pros and cons of different forms of plasticity. More generally, because this work is very numerical, a journal like PLOS comp biol would be more appropriate than PLOS biol.

We appreciate the reviewer's point that numerical approaches are often undervalued, despite the need for such methods when models consider even ubiquitous aspects of ecology like resource limitation or top-down controls (such as via immune clearance). For example, Koella & Antia (1995, *Theoret Pop Biol*) examined plastic transmission investment using a general model of infection and immunity, an

approach that required numerical simulations despite strong simplifying assumptions like resource-independent parasite proliferation and perfect information about the time until host death or immune clearance. Thus, our model would need to be considerably simplified to become tractable to analytical methods like optimal control. Though predictions about the pros and cons of different forms of plasticity could be made from such a simplified model, it is not clear how those predictions could be generalized to diverse organisms (or to malaria parasites), all of which are constrained by limited resources. Further, previous *PLoS Biology* articles suggest the journal places value on the use of strictly numerical methods to understand system-specific evolutionary pressures, including in malaria parasites (e.g., Bushman et al. 2018). Here we go beyond numerical methods by ground-truthing inferences drawn—specifically about the features of the best dual cue combinations—with empirical data.

Nonetheless, we take the reviewer’s point that streamlining complexity where possible will enhance the impact of this study, and we have attempted to do so, e.g., by reducing the number of dual cue combinations examined and greatly shortening the semi-stochastic modelling section to focus on key takeaways.

Minor comments:

I don't understand the logic of equation (19). Why the difference in the denominator?

In streamlining our manuscript, this analysis has been removed. Though for completeness, the equation being referred to estimated the proportion of the deviation in fitness observed in our semi-stochastic model that could be attributed to a single parameter. The denominator represented the fitness deviation observed when all six parameters were varied simultaneously compared to the deterministic model, while the numerator was the fitness deviation observed with only one parameter varying. We removed this analysis since it was not core to the takeaway that there is a tradeoff between accuracy in a given environment and susceptibility to noise.

Some discussion about the ability of the malaria parasite to perceive the density (or the log(density) of cells (infected or not) is needed. The model relies on this ability and it is important to discuss the feasibility of this assumption. What type of signal could carry this information?

We thank the reviewer for this comment, which made clear to us that we had failed to establish in the introduction what is known empirically about the ability of *Plasmodium* parasites to detect and respond to their environment. We have substantially revised the introduction, and this information is now included, l. 28-38,

“Importantly, transmission investment in Plasmodium spp. is plastic, changing in response to conspecific density [11, 12], replication rate [13], parasite-conditioned media [14, 15] (which could contain infected red blood cell (iRBC)-derived vesicles [16]), RBC age structure/availability [13, 17–19], and antimalarial exposure (e.g., [13, 20]). Understanding why transmission investment is plastic and what the optimal plastic strategy is has been the subject of several prior studies [7, 13, 21–24]. In one example, using the experimentally tractable rodent malaria system, Plasmodium chabaudi, and different doses of antimalarial drugs, Schneider et al. showed that intermediate reductions in parasite density led to reduced investment in transmission

while strong reductions had the opposite effect. These results are in line with standard predictions from life history theory...”.

We have also added related text to the Discussion, e.g., l. 376-381,

“Prior studies have found that Plasmodium transmission investment varies in response to conspecific density [11, 12], parasite-conditioned media [14, 15], and infected red blood cell (iRBC)-derived vesicles [16], suggesting that parasites possess the ability to sense iRBC density. Our results support these responses being part of an adaptive strategy.”

Reviewer #2:

Wang et al. studied the intricate timing of investment into reproduction in malaria parasites. Specifically, they examined the types and number of cues sensed by the parasites' sexual reproduction investment to maximize their fitness. They found that dual cues instead of single cue are required to realize the terminal investment and logged cues are necessary for the delayed investment. Dual cue sensing also increases disease virulence. However, dual cue perceptions have tradeoffs with the fitness cost due to environmental stochasticity. Overall, I think the manuscript is very well written. I have two general questions regarding the model assumptions and some minor comments.

We greatly appreciate the reviewer's feedback, which has significantly improved the paper's readability.

Q1: It's not immediately clear to me how the time-varying optimal investment strategy is derived (Fig. 1A, Fig. 7). In the "fitness optimization" section, it only documented how the optimal trajectory is derived using single or dual-cues, but not for the scenario when investment is a function of time directly (as investment due to responses to the cues are delayed in time).

We thank the reviewer for bringing this omission to our attention. We have now made explicit in the methods how we do this (it is essentially the same as the single cue models, except no Heaviside transformation is required), l. 620-622,

“For the best time-varying reaction norm, we defined the perceived cue as the time since infection, which has a range of 0-20 days and a step size of 0.001 (same as the simulation time step). In this case,

$$c(t) = \exp(-\exp(B3(t))).”$$

Q2: When investigating the impact of environmental stochasticity on the optimal transmission investment, the authors assume the parasites use the same optimal strategy inferred from the deterministic model under varying changes of parameters, such as burst size and immune activation strengths. Is that a reasonable assumption? If the optimal strategy is a result of long-term selection from host-parasite interactions, the outcome would be the strategy that performs overall the best under varying parameter combinations instead of a specific strategy trained from the deterministic model. I wonder if a different optimal strategy can be trained while allowing for realistic variation of the six parameters.

This comment is related to another from Reviewer #4, who identified our implicit assumption that parasites switch hosts very quickly. We now make this assumption explicit, l. 253-257,

“Specifically, the optimal reaction norms identified above are associated with a particular set of model parameters (the median estimates from posterior distributions in [41]; we refer to this as the deterministic model). This implicitly assumes that hosts are all the same or that parasites are moving between hosts sufficiently quickly that they adapt to an “average” environment.”

and, l. 271-273,

“Here we are asking, if parasites are “optimized” to the average host, does environmental and developmental variation change inferences about which cues are best”?

Although it would be answering a different question than the one we have posed, it is in principle possible to find the optimal strategy in the face of variable parameters. Technically, this would be challenging for a few reasons. First, if parameters are chosen randomly over a series of infections, this will make convergence of the optimization algorithm difficult. Second, if instead a specific series of parameter sets is chosen, then more assumptions need to be made (e.g., How many parameter sets? Are infections with each parameter set equally likely?). We note that with enough parameter sets, chosen with weights defined by the posterior distributions that we have relied on, we’d expect to find the same optimal strategies as using the medians of those distributions, as we have done. Thus, we think the assumption we have made is a reasonable place to start.

Also, I'm a bit confused by why "burst size" or host immune responses are considered environmental factors. Parasite burst size is determined by parasites' genetic factors and host-parasite interactions, which are part of the overall parasite sexual and asexual reproductive strategy that is also under selection.

We have defined the host immune response as an environmental factor, given that the immune system is an aspect of the environment experienced by the parasite. We agree that parasite genetics will have a large influence on burst size, though environmental influences have also been shown. Specifically, Stürmer *et al.* (2023, *PLoS Pathog*) recently demonstrated *in vitro* that nutrient level mediates burst size in *P. falciparum*. We have amended our language throughout to capture more sources of variation (e.g., “environmental and developmental stochasticity”).

Minor comments:

- Fig 2B, legend label, "none" should be "raw". More explanation for Fig 2 is required: It seems that when using logged cues, transmission investment is very high for low densities compared to raw cues (Fig. 2B), this seems to be counterintuitive with the fig 2A result when delayed responses are observed with logged cues.

We have corrected the Fig 2B legend label to “raw” and edited the caption for clarity. In light of this comment, we have also redrawn the plots in Fig 2B, with log-transformed x-axes for easier visualization at low densities. The revised plots better illustrate that transmission investment is indeed lower for the logged cue at low densities of parasite-centric cues.

- Lines 237-243, seems to be discussion contents instead of results.

We agree. This text has been shifted to the discussion section (l. 466-474).

- Fig 5. The legend says fitness from semi-stochastic model is rarely larger than that from the deterministic model, but it's not immediately clear from violin plots of Fig 5B, especially for burst size.

In revising our paper to make it more streamlined and clear, we decided that this figure and its associated analysis (determining which parameter contributed most to the loss of fitness in the semi-stochastic model) was not central to our main message. Thus, we have removed this figure and replaced it with another that is responsive to other reviewer comments, illustrating the tradeoff between accuracy in a given environment and susceptibility to (or precision in the face of) noise.

- Fig 6. The meaning of colors and lines in B-C are not explained

We have similarly removed Figure 6 in streamlining the article. We note that the bottom panel of our new Figure 5B is similar (though not identical) to the previous Figure 6B&C, showing the consequences for transmission investment when particular cues are perceived in the face of environmental or developmental stochasticity. (We now also show the scale of the variation in those cues from the same simulations in the top panel of Figure 5.) We have been careful to describe the different coloured lines in the Figure 5 caption.

- Fig 7. Why are the regression lines required here? Seems to be unrelated to the main message you are conveying. What do the grey points represent in the figure?

Good point — the lines weren't required and we have removed them from the figure (now Figure 6 in the revised version). The grey points represent all cues that we have not colour-coded in the legend. We have added additional clarification in the figure legend to explain this.

Reviewer #3:

Wang et al present an interesting theoretical model focused on understanding the role of cues in reproductive investment using the well studied rodent plasmodium model system as a case study. The study presents a significant advancement in understanding how malaria parasites regulate transmission investment in response to their within-host environment. The authors successfully integrate life history theory with mathematical modeling, providing a robust theoretical framework that aligns with empirical observations of transmission investment plasticity. Their results are a major step forward because they relax the assumption that parasites operate with perfect knowledge of infection time and instead propose a more biologically plausible mechanism based on environmental cue perception.

One of the paper's key strengths is its rigorous theoretical approach, which systematically evaluates various putative cues and their impact on parasite fitness in a well understood system. The distinction between single and dual cue perception is particularly insightful, demonstrating the advantages of state differentiation in guiding reproductive decisions. The finding that dual cue perception increases virulence also aligns with broader evolutionary theories that link optimal resource exploitation with pathogen fitness.

The key focus on the impact of different cues on the ability of the parasite to get more optimal outcome is very original. This is a strong reason to publish the paper, since it is a major and general conceptual advance. We don't normally think about how information on environment is some sense optimally used for organisms to optimize their fitness. It's a very general problem that has not been sufficiently addressed. Also the paper is focused on reproductive effort which is a fundamental trait and moreover using the rodent malaria system is excellent since the investment in reproduction as opposed to growth is very clearly defined. Much more so than in most systems. This allows a really comprehensive analysis. Finally the fact that they are studying malaria means that they can relate their results to important practical questions such as the evolution of virulence.

Overall, I think this is a tour de force of a paper, original in concept and extremely rigorously carried out. My criticisms emerge from just how detailed it is which results in a slightly overwhelming presentation of results and details. I think the paper would be strengthened with more discussion of the key result that multiple cues are needed for the parasites to recognize the need for terminal investment. This is really interesting and although it is discussed, I think an expanded discussion would help people understanding this intuitively. There are a lot of interacting parts here in terms of cues, and the complexity of the models, so perhaps some more intuitive discussion of two of the cues individually and why they don't work on their own but do when combined would illustrate the points better. The paper is very long and perhaps pulling this key discussion into a conclusions section would help with the structure.

We would like to thank the reviewer for their encouraging feedback. In response to this comment and others, we have streamlined our manuscript, removing details and analyses that were not crucial to our main messages, and have offered more intuitive explanations whenever possible.

Another interesting advance in the paper is the discussion of emerging tradeoffs involved in multiple cue perception. Again I think the paper would be improved with a more intuitive overview of how these trade-offs emerge and where they are underpinned by assumptions. It makes sense that while the dual cue perception improves parasite fitness and enables more precise transmission strategies, it also increases susceptibility to environmental and developmental fluctuations, but I think it would have a greater impact if people can see where this emerges from intuitively.

We agree with the reviewer that the paper needed editing to provide more intuitive understanding throughout, and we hope we have achieved that. With respect to the fitness advantage of dual cue strategies and the resulting tradeoff that emerges, we have done the following. First, we have added a new paragraph at the beginning of the dual cue results section (l. 156-166), and a new panel in Figure 3, to give readers an intuitive understanding of why single cues fail to reproduce the expected optimal pattern

of transmission investment, based on the dynamics of those single cues. This sets up the expectation that two cues will be able to recreate more aspects of the optimal investment pattern and motivates the question of which set of cues is best.

Second, we have streamlined the section on environmental and developmental stochasticity, dispensing analyses that were not core to understanding the tradeoff (e.g., exploring which model parameter drives fitness losses in variable environments). With the focus of this section now squarely on dual cue strategies being more susceptible to noise, we again appeal to the dynamics of the cues to show the reader how and why this tradeoff emerges. We do this through a new paragraph at the end of that section (l. 290-303) and an updated Figure 5B, which demonstrates that variation in cue trajectories over the course of an infection gives rise to wildly different patterns of transmission investment in the dual cue, but not the single cue case. This is because deviations from the median dual cue trajectory traverse qualitatively different paths through the finely-tuned dual cue reaction norm surface (shown in the top panel of Fig 5B).

Another important result that could be highlighted more is that, dual cue perception correlates with higher virulence, as it allows parasites to exploit hosts more effectively, potentially leading to more severe infections. This has implications for understanding parasite evolution, within-host competition, and malaria transmission dynamics, particularly in response to interventions like drug treatment or vaccines. Again I think the underlying assumptions and intuitive response for this result could be highlighted more in order to make this result pop.

We agree that this is an important result! We hope that by streamlining and simplifying the section that comes directly before the one on virulence (i.e., the section on stochasticity), that these results will “pop” more. We have also been careful to define virulence throughout the manuscript and have edited the paragraph in the discussion that relates to these results (l. 420-430).

All of this advice is given in the hope that the paper can have a broader impact beyond the rodent malaria model system community. I think the general insights that can be gained here have the potential to influence more general reproductive allocation and optimality studies. At the moment the paper is very detailed, which is a key strength, this is rigorous well grounded theory, but the intuition behind some of the big results is a little buried. Highlighting this will allow the paper to have a bigger impact. I would suggest a major rewrite of the discussion with either new paragraphs that give specific examples that allow the intuitive explanations for why individual cues don't lead to terminal investment and a broader implications outside of the model rodent malaria system, or perhaps an intuitive explanation and wider implication sentence in each paragraph.

We thank the reviewer for this incredibly helpful advice. We hope that by expanding the results section to incorporate more intuitive explanations of key inferences, in combination with removing unnecessary detail and analyses, we have produced a manuscript that is easier to read, with the potential for greater impact. With respect to the discussion, we have made a number of changes. Here we highlight three. First, we provided additional information linking our modelling result on the benefit of perceiving parasite-centric cues to the broader, empirical literature (l. 377-381),

“studies have found that Plasmodium transmission investment varies in response to conspecific density [11, 12], parasite-conditioned media [14, 15], and infected red blood cell (iRBC)-derived

vesicles [16], suggesting that parasites possess the ability to sense iRBC density. Our results support these responses being part of an adaptive strategy.”

Second, we rewrote the discussion section detailing the benefits and drawbacks of sensing multiple cues. To give readers a more intuitive understanding of why dual cue perception is beneficial, we provided a simplified explanation on why perceiving a single cue cannot allow for terminal investment (l. 387-390)

“Since most within-host variables go up and down, or vice versa, over time (e.g., iRBC densities rise and then fall due to resource limitation or immunity), one cue cannot provide enough detail to unambiguously establish “where” a parasite is in the course of infection. Combining cues solves this problem.”

Third, we expanded our discussion of sensing multiple cues in order to make connections to the broader literature on phenotypic plasticity (l. 399-410)

“There is a growing appreciation that, across diverse organisms, plasticity is often expressed in response to multiple cues that allow for more finely-tuned strategies [70–72]. Our work adds support to theories suggesting that the adaptive benefits of phenotypic plasticity are enhanced when cues provide non-redundant information [73] and enable organisms to sense distinct states that correspond to specific optimal phenotypes [74]. Yet it is also widely appreciated that integrating multiple cues presents challenges, including the additional metabolic burden of sensing another cue [75] and the multiplicative effect of noise when cues are non-additive [72]. Our complex (decidedly non-additive) dual cue reaction norms present a trade-off: they allow parasites to more accurately recreate optimal strategies in a given environment while rendering them more susceptible to stochastic environmental and developmental change, i.e., host-to-host variability.”

Minor comments

While the model focuses on *P. chabaudi*, extrapolating these findings to *P. falciparum*, the primary human malaria parasite, requires caution. *P. falciparum* has a longer gametocyte maturation period, which could influence cue reliability and transmission investment dynamics.

We agree with the reviewer that extrapolating our findings to *P. falciparum* requires caution. We have noted the discrepancy between *P. chabaudi* and *P. falciparum* sexual developmental timeline in the discussion and explained how those differences could affect cue reliability (l. 436-440),

*“The human malaria parasite *P. falciparum* has a longer sexual development period of 9-12 days [85] and this longer lag between cue perception and phenotype expression will reduce cue reliability [86]. Thus, we anticipate the value of sensing gametocytes to decrease even further in this species.”*

The model tracks the dynamics of red blood cells (RBCs) and various parasite stages within the host over a 20-day infection period. to identify the best strategies for maximizing transmission fitness, the study employs a combination of local and global optimization algorithms: I think it would be useful to explain the broader uses of the dual methods of the L-BFGS-B algorithm (Limited-memory Broyden-Fletcher-Goldfarb-Shanno with Box constraints) and Global Optimization through the Differential Evolution (DE) algorithm, implemented in the DEoptim

package. This hybrid approach helps identify globally optimal reaction norms and could be used more widely.

We agree that taking a hybrid approach has greatly facilitated our search for global optimums. However, we would like to clarify that the suitable choice of optimization algorithm varies across models and requires empirical determination. Therefore, we are hesitant to recommend the L-BFGS-B and DE algorithms for other optimization problems broadly.

Reviewer #4:

This was an interesting paper that applied a variety of mathematical approaches to explore the question of parasite plasticity in infection, using the reproductive investment of malaria parasites as a case study in this question. The authors find that cue perception mediates parasite life history and fitness and they perform optimization over several hypotheses for cue perception and identify key infection variables that singly or jointly determine the optimal cue. They go even a step further and explore optimal strategies in the context of local vs. global optimum, considering host and infection parameter heterogeneity. I enjoyed reading this paper, and think that it demonstrates interesting biological insights but I am afraid that there is too much going on and some parts need attention, clarification and streamlining. Specific comments are below:

We thank the reviewer for their positive feedback. We agree that there was too much going on, and we believe this revised version is much improved and clearer.

Overall:

This paper relies a lot on results from past work by the same authors, particularly for the model construction and motivation to explore the variable-dependent strategy of reproductive investment. I would like to see more details on the relevant conclusions from this previous work whenever it is cited in this paper, so as to make the paper more self-contained, and highlight more the rationale for the new approach adopted here. Especially in the introduction and beginning of results, e.g. line 82, it seems the reference to previous work is too quick and insufficiently informative. I would really suggest to highlight more the results from the previous work, the use of time as a proxy for "environmental quality" over infection, but a kind of arbitrary proxy, that in different infections could be inaccurate, so there is a basis for exploring concrete infection variables as more informative and parameter-transcending cues.

The reviewer makes an excellent point about time being an arbitrary proxy that is likely to be inaccurate in different infections. We have added text to the introduction to highlight this point, l. 55-63,

“In addition to assuming that parasites can, in effect, “tell time”, this work also assumes all infections proceed identically, a limitation common to models of age-specific reproductive investment [28]. In reality, time would be a poor proxy for environmental conditions since the pace of infection changes with parasite dose [29, 30], host immunity [30, 31], and myriad other

factors. Understanding the links between transmission investment and within-host environmental cues is necessary to validate modelling results with experimental evidence and for predicting the influence of novel perturbations, like vaccination, on infection dynamics and outcomes.”

In response to this comment, we also updated Figure 5A (comparative fitness of cues in the deterministic and semi-stochastic models) to include parasites sensing time. This plot makes clear that under varying conditions, parasites that sense time perform in the bottom half of the cues we tested, a result we now highlight in the text (l. 282-284).

Finally, we also broadened the introduction to rely less on previous theoretical work and include more detail on what is known empirically about the environmental influences on transmission investment in malaria parasites (l. 28-37).

I am not sure also I understand the big aim of this paper. Is the terminal investment the big issue here? The authors are trying to find 'internal' infection mechanisms for terminal investment in reproduction? E.g. the aim articulated in the lines 135-140?

We apologize for having been unclear about our main aims. Our goal was to understand what cues parasites *should* sense when making decisions about transmission investment. While terminal investment is an aspect of the “ideal” transmission investment strategy outlined in prior work, it is not the only aspect that we are looking for. For instance, delaying transmission investment at the start of infections leads to much higher fitness gains than does terminal investment at the end of them. Throughout the manuscript, we have tried to better signpost our aims, e.g., in the abstract,

“We extend that theory to ask which cue(s) should parasites use? Put another way, which cue(s) maximize parasite fitness, quantified as host infectiousness during acute infection?”

and in the introduction, l. 74-84

*“We use numerical optimization to investigate which within-host environmental cue or cues malaria parasites “should” sense for determining transmission investment (i.e., which cues give rise to the highest parasite fitness) and the optimal shape of the relationship between each cue and transmission investment (i.e., the optimal reaction norms). We assess the consequences of perceiving different cues on infection dynamics and disease severity and quantify the robustness of different transmission investment strategies in the face of environmental and developmental stochasticity. Our study further characterizes plasticity in *P. chabaudi* life history, identifies the cues we expect parasites to be using, and introduces a mathematical framework that explicitly incorporates cue perception that could be broadly applied to other organisms.”*

In my view, the paper would benefit from shortening and streamlining and clearer focus. As it is now, a lot of results are presented, they are all interesting, but somehow it's easy to miss the forest for the trees, the findings are all complex and hard to understand together, the model, cue definition, the optimization, the interpretation of the joint cues, the environmental variation, the optimization over environmental variation. The authors could be more selective and decide to divide the big story in two parts that would be able to breathe more and speak more in separation. This is a suggestion.

We completely agree with the reviewer that our original paper was trying to cover too much ground. We decided to remove analyses and text that were not central to our key takeaways, that (i) sensing two cues enables parasites to state differentiate, make better transmission investment “decisions”, and achieve higher fitness, (ii) there is a trade-off between accuracy in a given environment and sensitivity to environmental noise, and (iii) the evolution of plastic transmission investment will influence virulence.

To streamline the paper, we have:

- 1) Focused on a subset of dual cues instead of all possible pairs of single cues (i.e., only combinations of log-transformed cues given their higher performance in the single cue optimizations, see Fig 3B and l. 173-175).
- 2) Removed a figure panel (formerly, 5B) and analysis of which parameter in the semi-stochastic model explained the fitness decline, since it was not central to our takeaway about the tradeoff (ii above). We have replaced that subfigure with another showing the dynamics of the best single and dual cues and their associated patterns of transmission investment, which help explain how and why that tradeoff emerges.
- 3) Removed several paragraphs and another figure (formerly Figure 6) that investigated the mechanism by which dual cue perception sensitizes parasites against environmental stochasticity. We think the intuitive explanation offered by our new Figure 5B is sufficient.

Structure:

- The paper starts with results, but the model is very crucial to understand the results. I wonder if it would be possible to move the materials and methods before the results. Specifically, for unfamiliar readers, the model is not presented in sufficient detail and especially the key mechanisms that drive the end of infection, e.g. the role of resource limitation and immunity. Immunity is not mentioned until much later. This makes it difficult to understand what are the key players mathematically.

We assume that we are limited in restructuring the article, as per PLOS Biol guidelines, although we are open to editorial feedback on this. Nevertheless, we acknowledge the reviewer’s point that there was insufficient detail about the model. In light of this comment, we have substantially revised the first two paragraphs of the results, which describe the model. We have also added a new plot to Figure 1 that illustrates our approach to modeling reaction norm evolution (Figure 1C).

- Some part that is currently in the discussion, e.g. lines 417-428 would be more useful to be put earlier, perhaps in the introduction. Again the rationale for the phenotypic plasticity modeled in this way is important to be highlighted more.

We thank the reviewer for the suggestion and agree that there was insufficient background in the introduction on what is known empirically about *Plasmodium* spp. transmission investment and environmental influences. As noted above, we now include this detail, l. 28-37,

“Importantly, transmission investment in Plasmodium spp. is plastic, changing in response to conspecific density [11, 12], replication rate [13], parasite-conditioned media [14, 15] (which could contain infected red blood cell (iRBC)-derived vesicles [16]), RBC age structure/availability [13, 17–19], and antimalarial exposure (e.g., [13, 20]). Understanding why

transmission investment is plastic and what the optimal plastic strategy is has been the subject of several prior studies [7, 13, 21–24]. In one example, using the experimentally tractable rodent malaria system, Plasmodium chabaudi, and different doses of antimalarial drugs, Schneider et al. [13] showed that intermediate reductions in parasite density led to reduced investment in transmission while strong reductions had the opposite effect. These results are in line with standard predictions from life history theory for malaria parasites...”

- I also think the measures that are focus of the investigation, like total transmission fitness or virulence should be stated before the results are presented. It's hard to track the meaning of these different measures, without them being defined.

Great point. We have added the definition of fitness in the first paragraph of the Results, l. 99-102, “*We quantify the fitness of parasites sensing various cues using cumulative transmission potential, which measures the summed probability of infecting mosquito vectors across a 20-day infection period, assuming a constant biting rate and mosquito abundance [23].*”

We note that our operational definition of virulence is given at the start of the section “Dual cue perception leads to higher virulence” (l. 307-311). We are also careful in the abstract and result to equate virulence with infection severity, though we have also made this explicit (l. 315-319, also see response below).

- The model variables also need to be explained in a more reader-friendly manner. For example, start with the uninfected cells, then infected cells, etc. the model without cue- in a clearly-titled subsection of the methods, and then the subsection with the cue highlighting its formulation, and its exact place in the classical model. But please put the equations before and the cue formulation later. As it is now, the cue comes before the other model elements (equations 10-11..).

Thank you for the suggestions. We have rearranged the Methods section such that the model without a cue is described first, followed by a new subsection describing how transmission investment is captured.

Another suggestion is with regards to the order of presentation of the variables and behaviors that define the stages of infection (lines 160 -190) and the optimization (previous paragraphs). Now the optimization is presented first, then the stages of infection as a supporting interpretation. I think for people who read the paper, it would be more useful and feel somewhat more natural to get familiar with the model, infection behavior, stages, etc. and then build an intuitive expectation which leads up to the optimization search and confirms what one would expect from this sort of preliminary sensitivity analysis based on pure study of model behavior and coupling between variables of a dynamical system.

This is a great point and led us to reflect on how to provide the reader with an understanding of the model's behaviour and an intuition about what to expect in the dual cue models. To this end, we have added an additional panel in Figure 3 (Figure 3A) and a new paragraph at the beginning of the dual cue results section, l. 156-169,

“We found that parasites sensing a single cue cannot reproduce the expected optimal transmission investment strategy, particularly the rapid increase in transmission investment during late infection (i.e., terminal investment). Achieving terminal investment presumably requires sensing a cue that provides a unique signal at the later stages of infection, avoiding prematurely increasing investment and reducing fitness. Inspecting infection dynamics produced by parasites that sense time revealed that only RBC density provides such a distinctive late-stage signal (Fig 3A). Given the poor performance of this cue in early stages of infection, these results suggest that simultaneous perception of RBC and a parasite-centric cue could provide infection stage-specific information that enables both early reproductive restraint and terminal investment. Consistent with this expectation, in a rodent malaria experiment, including variables that reflect parasite density and host resources (specifically, replication rate and changes in RBC density) in statistical models explained substantially more variation in transmission investment than the parasite-centric variable alone [13].”

Other comments

- I am a little confused about the way immunity effects are implemented in the model. I have not seen earlier this formulation with the logarithm $(-\ln(1-N(t)))$ why not use a simple mass-action term for the interaction between immunity and infected cells? Can the authors please elaborate on the foundation origin and interpretation of this formulation?

We have added further clarification on this formulation, which is borrowed from our previous work (Kamiya et al. 2020 *PLoS Comp Biol*), l. 497-525,

“As in [41], we define N and W as the proportion of cells cleared per day by indiscriminate and targeted responses, respectively...

Because N and W have been defined as proportions of cells cleared, they can be rewritten to

reflect daily rates of clearance. For example, $N = 1 - e^{-x_N}$, where x_N is equal to the rate of clearance on a given day, and is defined as $-\ln(1 - N)$, with some rearranging of terms. A similar expression provides the daily clearance rate for targeted killing, $-\ln(1 - W)$.”

- In the definition of parasite fitness, the probability that the host is alive is not taken into account. Normally the integral would also take into account host survival, e.g. a decreasing exponential, function of total parasite load. Why is this not taken explicitly into account in equation 18? While on the other hand the authors also track the "virulence" but in a decoupled manner from transmission fitness.

The reviewer is correct that we track virulence decoupled from parasite fitness, due to growing recognition that disease-induced mortality may be too rare to serve as an important constraint on the transmission success of pathogenic organisms (e.g., Bull & Luring 2014 *PLoS Path*; Kennedy 2023 *Proc B*). In the context of malaria parasites, the overwhelming majority of infections are subclinical and nonlethal, at least in humans where data on natural infections are available (Lindblade et al. 2013 *Expert Rev Anti-Inf Therapy*). Recent modeling of *P. chabaudi* infections supports the idea that other factors, especially resource limitation, form a more important constraint on parasite fitness than host mortality (Pak et al. 2024 *Evolution*). We now clarify these points in the text, l. 315-319,

*“We note that we are using “virulence” as synonymous with infection severity and not host mortality, thus we assume no link between virulence and the duration of infection. Recent modeling suggests that resource limitation imposes a more important constraint on the evolution of transmission investment in *P. chabaudi* than does host mortality due to infection [61].”*

- I would suggest writing equation 17 in a more general form and simplifying the exponential^(log..) it becomes just $A * G(t)/(1+B * G(t))$ where A and B are just constants and then replace the particular numerical values. It's hard to see its general shape in this current formulation.

We have replaced equation 17 with a more general form, as suggested (now equation 18).

- I am not sure the comparison of the optimized reaction norms with random splines is relevant or particularly informative.

The comparison with random splines serves as a sanity check to ensure our optimization is successful. In the past, when our optimization was not successful, we found that random splines performed better than our “optimized” splines. We have left the associated text (l. 732-737) and supplementary figure (S1) in for now, but are happy to take editorial direction on removing it.

- In lines 580-586, it would be useful to state clearly the optimization framework max(...) over the model coefficients (parameters to be searched).

We now specify in the Methods that the maximum bounds are “defined in Table 3” (l. 616-617).

- In the global optimum part, I am not sure the host heterogeneity that the authors mean, is correctly referred to, when they use the term stochasticity. I would warrant them to consider more carefully the two concepts as environmental stochasticity is one thing, host heterogeneity is another. And it would be very important to define the rate of 'host switching' or 'environmental switching' and see how to local optimum compares to the global optimum. I am not sure the authors did that. In the limit the environment is not switched at all - we have the local optimum, in the other extreme where the hosts are switched very rapidly, there is no point in optimizing to any given host but to the average host. In between, things can be complicated. I wonder if in their simulations with random parameters, the authors are implicitly assuming infinitely fast rate of switching environments.

We thank the reviewer for this insightful comment, which gave us a deeper appreciation for the implicit assumption of our approach. For our deterministic model, most of our parameters are derived from the medians of posterior distributions previously generated (Kamiya et al. 2020 PLoS Comp Biol) and also used in the semi-stochastic modelling. This means that our deterministic optimization can be thought of as optimizing towards the average host when host-switching is fast, or that all hosts are the same. We know make this explicit in the text, l.253-257,

“Specifically, the optimal reaction norms identified above are associated with a particular set of model parameters (the median estimates from posterior distributions in [41]; we refer to this as

the deterministic model). This implicitly assumes that hosts are all the same or that parasites are moving between hosts sufficiently quickly that they adapt to an “average” environment.”

We also clarify our assumptions and aims with the semi-stochastic analysis, l. 271-275,

“Here we are asking, if parasites are “optimized” to the average host, does environmental and developmental variation change inferences about which cues are best? We quantified the effect of variation on cue performance using the geometric mean fitness, commonly used to assess fitness in fluctuating environments [59, 60].”

- I also would like to remark that the authors seem to not take into account the potential effect of temporal distribution of transmission events (being bitten by a vector) - it's assumed implicitly uniform over infection. This distribution could definitely have an effect on the optimal parasite strategy. So far, only the end of the infection is taken as the optimization horizon.

The reviewer is correct that we have assumed “the mosquito population is at a constant abundance and continuously biting, for the duration of the infection” (l. 642-643). In the Methods, we note that “[a]lthough this is a simplification of the evolutionary pressures faced by malaria parasites [35], it enables us to focus our investigation at the within-host level, where transmission investment “decisions” are being made and environmental cues are being perceived by parasites.” We note that we explored the effect of relaxing this assumption elsewhere (Greischar et al. 2019 *Evol*), but doing so necessitated a considerable simplification of another aspect of the biology, namely, in that work we assumed transmission investment was constant over the course of infections.

- It would be interesting, the authors mention it- but do not go deeper, to associate 'sensing' to a cost. The benefits of sensing and adapting their infection strategy, would have to be outweighing any costs for the parasite.

We agree with the reviewer that adaptive phenotypic plasticity is dependent on the benefit of plasticity outweighing the cost of perception. To that end, we have specified that one of the drawbacks of sensing multiple cues is the added metabolic burden (l. 404-406),

“Yet it is also widely appreciated that integrating multiple cues presents challenges, including the additional metabolic burden of sensing another cue [75]”

We have decided not to delve deeper into this topic, given the limited literature on the actual cost of cue sensing and the added complication that sensing different cues may have varying costs. Metabolic cost aside, we anticipate a bigger cost of sensing the “wrong” cue would be derived from a suboptimal transmission investment strategy, which we investigate in the manuscript.

- The authors mention their model choice is to depict immunologically naïve hosts. Why did they not simply vary initial conditions of their immunity variable to mimic the effect of immunological memory, in a similar way that they varied the other parameters? It seems straightforward, but maybe I am missing something. It would be good to clarify.

We decided to depict immunologically naive hosts due to the availability of immunological parameters for *Plasmodium chabaudi* in naive lab mice (Kamiya et al. 2020 *PLoS Comp Biol*). We have added additional clarifications about this choice in the Methodology (l. 522-525). We agree with the reviewer that understanding the impact of immunity on *Plasmodium* transmission investment evolution is important and could potentially be achieved by making assumptions about how immunological parameters might vary and rerunning the optimization algorithm. Given the consensus among reviewers that there was already too much in our manuscript, we think this is an interesting extension to pursue in the future, as mentioned in the discussion (l. 444-447).

- In the discussion it would be useful to clearly remark and emphasize which are the new insights on Plasmodium biology and which are broader novel insights obtained in this work.

We agree with the reviewer that the Discussion will benefit from streamlining and emphasizing further connections to the broader literature. To that end, we have made a number of changes. Here we highlight three. First, we provided additional information linking our modelling result on the benefit of perceiving parasite-centric cues to the broader, empirical literature (l. 377-381),

“studies have found that Plasmodium transmission investment varies in response to conspecific density [11, 12], parasite-conditioned media [14, 15], and infected red blood cell (iRBC)-derived vesicles [16], suggesting that parasites possess the ability to sense iRBC density. Our results support these responses being part of an adaptive strategy.”

Second, we rewrote the discussion section detailing the benefits and drawbacks of sensing multiple cues. To give readers a more intuitive understanding of why dual cue perception is beneficial, we provided a simplified explanation on why perceiving a single cue cannot allow for terminal investment (l. 387-390)

“Since most within-host variables go up and down, or vice versa, over time (e.g., iRBC densities rise and then fall due to resource limitation or immunity), one cue cannot provide enough detail to unambiguously establish “where” a parasite is in the course of infection. Combining cues solves this problem.”

Third, we expanded our discussion of sensing multiple cues in order to make connections to the broader literature on phenotypic plasticity (l. 399-410)

“There is a growing appreciation that, across diverse organisms, plasticity is often expressed in response to multiple cues that allow for more finely-tuned strategies [70–72]. Our work adds support to theories suggesting that the adaptive benefits of phenotypic plasticity are enhanced when cues provide non-redundant information [73] and enable organisms to sense distinct states that correspond to specific optimal phenotypes [74]. Yet it is also widely appreciated that integrating multiple cues presents challenges, including the additional metabolic burden of sensing another cue [75] and the multiplicative effect of noise when cues are non-additive [72]. Our complex (decidedly non-additive) dual cue reaction norms present a trade-off: they allow parasites to more accurately recreate optimal strategies in a given environment while rendering them more susceptible to stochastic environmental and developmental change, i.e., host-to-host variability.”

Optimal cues for transmission investment in malaria parasites

Avril L. Wang^{1□*}, Megan A. Greischar², Nicole Mideo¹

1 Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, Ontario, Canada

2 Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, New York, United States

□Current Address: Department of Biology, Massachusetts Institute of Technology, Cambridge, Massachusetts, United States

* avril137@mit.edu

Abstract

The timing of investment into reproduction is a key determinant of lifetime reproductive success (fitness). Many organisms exhibit plastic, i.e., environmentally-responsive, investment strategies, raising the questions of what environmental cues trigger responses and why organisms have evolved to respond to those particular cues. For malaria parasites (*Plasmodium spp.*), investment into the production of specialized transmission stages (versus stages that replicate asexually within the host) is synonymous with reproductive investment and also plastic, responding to host- and parasite-derived factors. Previous theory has identified optimal plastic transmission investment strategies for the rodent malaria parasite, *Plasmodium chabaudi*, as a function of the time since infection, implicitly assuming that parasites have perfect information about the within-host environment and how it is changing. We extend that theory to ask which cue(s) *should* parasites use? Put another way, which cue(s) maximize parasite fitness, quantified as host infectiousness during acute infection? Our results show that sensing a parasite-associated cue, e.g., the abundance of infected red blood cells or transmission stages, allows parasites to achieve fitness approaching that of the optimal time-varying strategy, but only when parasites perceive the cue non-linearly, responding more sensitively to changes at low densities. However, no single cue can recreate the best time-varying strategy or allow parasites to adopt terminal investment as the infection ends, a classic expectation for reproductive investment. Sensing two cues—log-transformed infected and uninfected red blood cell abundance—enables parasites to accurately track the progression of the infection, permits terminal investment, and recovers the fitness of the optimal time-varying investment strategy. Importantly, parasites that detect two cues more efficiently exploit hosts, resulting in higher virulence compared with those sensing only one cue. Finally, our results suggest a potential tradeoff between achieving an optimal transmission investment strategy in a given host environment and robustness in the face of environmental or developmental fluctuations.

Introduction

A key goal of life history theory is to understand how organisms should invest in reproduction over the course of their lives and explaining the substantial variation in patterns observed (e.g., [1]). Reproductive investment entails a resource allocation

1

2

3

4

tradeoff: resources allocated to growth cannot, generally, also be allocated to reproduction. Classic work predicts that reproductive investment will increase with age [2,3]. Additionally, reproductive investment is expected to depend on the state of an organism, a product of intrinsic, physiological factors and external, environmental ones [4]. Experimental data have shown that investment can be plastic, with age and state both shaping an organism’s reproductive efforts to maximize lifetime fitness (e.g., [5,6]). How organisms estimate the time remaining for reproduction—that is, what cues they utilize from their own physiology and the environment—remains an open question for many systems.

For sexually-reproducing parasites, patterns of reproductive investment can have clinical and epidemiological consequences. Protozoan parasites of *Plasmodium* spp.—the etiological agents of malaria—infect red blood cells of their host, and a given infected cell will either produce more asexual, cell-infecting stages (“growth” of an infection) or produce a single gametocyte, a sexually-differentiated stage that can be transmitted to mosquito vectors (potentially “reproducing” an infection). The proportion of infected cells that produce transmissible gametocytes, known in the malaria literature as the “conversion rate,” is a measure of investment in reproduction/transmission [7]. This trait has a direct link to infectiousness, since more gametocytes mean greater chance of onward transmission to a biting mosquito [8–10] and can influence the virulence or severity of infection, since all else equal, lower transmission investment means more host-damaging asexual parasites are produced [7].

Importantly, transmission investment in *Plasmodium* spp. is plastic, changing in response to conspecific density [11,12], replication rate [13], parasite-conditioned media [14,15] (which could contain infected red blood cell (iRBC)-derived vesicles [16]), RBC age structure/availability [13,17–19], and antimalarial exposure (e.g., [13,20]). Understanding why transmission investment is plastic and what the optimal plastic strategy is has been the subject of several prior studies [7,13,21–24]. In one example, using the experimentally tractable rodent malaria system, *Plasmodium chabaudi*, and different doses of antimalarial drugs, Schneider *et al.* [13] showed that intermediate reductions in parasite density led to reduced investment in transmission while strong reductions had the opposite effect. These results are in line with standard predictions from life history theory for malaria parasites, where mild stressors should promote reproductive restraint and strong stressors that signal imminent clearance of an infection should lead to complete investment in transmission [7], i.e., terminal investment [25]. But “stress” will vary over the course of infection—even in the absence of drug treatment—as parasites deplete host resources and hosts respond via immunity. These dynamics are thought to explain why transmission investment in malaria parasites is variable over the lifespan of infections (i.e., plastic) [11–13,22,26,27].

Mathematical modelling provides another route to understanding and predicting adaptive plasticity in parasite traits. Models assessing the fitness consequences of different patterns of transmission investment demonstrate that the best strategy involves initial reproductive restraint that facilitates within-host parasite population growth (rather than early investment in gametocytes that would have very low likelihood of onward transmission due to low densities), followed by an increase in investment once parasite densities have risen, culminating in terminal investment as the infection is ending [21,23]. This work derived the optimal transmission investment strategy by assuming that investment is a function of infection age, about which parasites have perfect knowledge. In addition to assuming that parasites can, in effect, “tell time,” this work also assumes all infections proceed identically, a limitation common

to models of age-specific reproductive investment [28]. In reality, time would be a poor proxy for environmental conditions since the pace of infection changes with parasite dose [29,30], host immunity [30,31], and myriad other factors. Therefore, understanding the links between transmission investment and within-host environmental cues will result in more biologically relevant models that can inform empirical observation of *Plasmodium* spp. transmission investment. Currently, it is unclear whether observed deviations from the theoretical optimal strategy are due to faulty theories or limitations in the cues that parasites perceive. Understanding the fitness consequences of sensing different cues can delineate those possibilities. Additionally, understanding what cues parasites should sense will enable better predictions on the influence of novel perturbations, like vaccination, on infection dynamics and outcomes.

In this study, we extend a previously-published model of within-host rodent malaria infection dynamics [23] to include transmission investment as a function of environmental cues, i.e., components of state. In our model, different cues may give rise to different early patterns of transmission investment, which will alter the trajectory of infections and feedback to influence the subsequent expression of the trait. We use numerical optimization to investigate which within-host environmental cue or cues malaria parasites “should” sense for determining transmission investment (i.e., which cues give rise to the highest parasite fitness) and the optimal shape of the relationship between each cue and transmission investment (i.e., the optimal reaction norms). We assess the consequences of perceiving different cues on infection dynamics and disease severity and quantify the robustness of different transmission investment strategies in the face of environmental and developmental stochasticity. Collectively, our study further characterizes plasticity in *P. chabaudi* life history, identifies the cues we expect parasites to be using, and introduces a mathematical framework that explicitly incorporates cue perception that could be used to generate testable hypotheses and broadly applied to other organisms.

Results

Our mathematical model tracks within-host dynamics of infection with the rodent malaria parasite *P. chabaudi* (Fig 1A). Asexual parasites, called merozoites (M), invade uninfected red blood cells (RBCs; R). The resulting infected RBC (iRBC; I) can go on to produce more merozoites that are either asexually-committed (M ; producing more merozoites if they infect an RBC) or sexually-committed (M_g ; producing a single gametocyte, G , if they infect an RBC). RBCs infected with these different types of merozoites are denoted as I and I_g , respectively. Importantly, the proportion of iRBCs that follow these different developmental routes (i.e., producing M_g) defines transmission investment. While previous approaches have treated transmission investment as either fixed or time-dependent (e.g., [21,23,32–36]), here we explicitly model it as a function of within-host environmental cues (see Materials and Methods). We select as plausible cues various measures of the density of parasites and host resources for which there is some experimental evidence to impact transmission investment in *Plasmodium* spp. (Table 1). We quantify the fitness of parasites sensing various cues using the cumulative transmission potential, which measures the summed probability of infecting mosquito vectors across a 20-day infection period, assuming a constant biting rate and mosquito abundance [23].

In our model, cue perception occurs at the beginning of asexual iRBC development (Fig 1B), determining the proportion of asexual iRBCs that produces sexually-committed merozoites. Note that this modeling choice introduces a time lag




Fig 1. Schematic of *P. chabaudi* within-host infection dynamics and fitness optimization. (A) Flow diagram of model tracking the dynamics of red blood cells (R ; RBCs), RBCs infected with asexual parasites (I), RBCs infected with sexual—but not yet mature—parasites (I_g), asexually-committed merozoites (M), sexually-committed merozoites (M_g), and gametocytes (G). (B) Temporal dynamics of cue perception. Cue perception occurs one day before the production of asexual/sexual merozoites, thereby introducing a delay between cue perception and the consequences of cue perception (i.e., production of more asexual infected RBCs or gametocytes). A further delay occurs in the sexual cycle since gametocytes of *P. chabaudi* take two days to mature [37]. Dashed circles indicate the bursting/maturation of a particular cohort of iRBCs. (C) Modeling reaction norm evolution. We model the relationship between the perceived cue and transmission investment (i.e., the reaction norm) as a flexible spline (black lines). Specifically, a cubic spline can specify any third order polynomial at a given x value as a linear combination of the columns in its basis matrix [38] (exemplified by the red, green, and blue dashed lines; combined they produce the solid, black line). For local optimization, an arbitrary starting spline is picked (left panel) and an optimization algorithm is used to adjust the relative weights of the basis functions until a fitness maximum is achieved (going from left to right). The small coloured arrows depict how the weight of a given basis function has changed from the previous panel.

between sexual commitment (as a result of cue perception) and gametocyte production, which is a departure from previous modeling work [23], but reflects empirical data on the canonical timeline of sexual commitment from the best studied system (*Plasmodium falciparum* [39]). For each putative cue, we define its associated reaction norm using flexible splines (Fig 1C). Specifically, the shape of the reaction norm is determined by a linear combination of three basis functions comprising a cubic spline. Adjusting the

spline coefficients (i.e., the weights associated with each column of the spline basis matrix) changes the shape of the reaction norm. We used a combination of local and global numerical optimization algorithms (see Materials and Methods) to adjust the spline coefficients and find the reaction norm that maximizes parasite fitness over a 20-day infection, as in previous work [23,40] (Fig 1C). We test both raw densities and logged densities as cues to expand the range of reaction norm shapes explored and because changes in infection dynamics are observed in this system when doses change by orders of magnitude [29]. For parasites sensing time, we used the same optimization process, except setting the cue range to time since infection (day 0 to 20). Finally, following [41], our within-host model includes two forms of immunity, one targeting specifically infected cells (I and I_g) and one targeting all red blood cells (I , I_g and R). Thus, while resource limitation plays some role in driving infection dynamics in this model, immunity allows for the clearance of the infection.

Table 1. Putative within-host environmental cues

Cue	Abbreviation	Experimental Evidence
Infected RBC (iRBC) density		[12, 13, 16, 24]
Asexually-committed	I	
Asexually-committed, log ₁₀	I log	
Sexually-committed	Ig	
Sexually-committed, log ₁₀	Ig log	
Total (I+Ig)	Total I	
Total (I+Ig) , log ₁₀	Total I log	
Gametocyte density		[12, 16, 24]
Gametocytes	G	
Gametocytes, log ₁₀	G log	
Resource density		[12, 13, 24]
Uninfected RBC	R	
Uninfected RBC, log ₁₀	R log	

Cue perception mediates parasite life history and fitness

Based on previous work modeling *P. chabaudi* transmission investment as a function of time [23], we expect the following as the best strategy over a 20-day infection: parasites should delay investment early on, display a rapid increase in investment when sufficient gametocytes can be produced to ensure transmission, subtly reduce investment in the face of declining resources, and ultimately rise to the highest levels of investment shortly before the end of infection (roughly, “terminal investment” [7,23]). Although the model we employ here has a few key differences from that work (see Materials and Methods), we find a similar optimal transmission investment pattern as in [23] when we assume that investment is a function of time (Fig 2A). When parasites instead respond to their environment and are constrained to perceive a single within-host cue, these same patterns of transmission investment cannot be fully recapitulated. Consequently, parasite fitness is lower in every case than when the cue is time (Fig 2A, left panel).

The environmental cues that result in the highest parasite fitness include logged parasite-centric cues (e.g., log₁₀ densities of sexually-committed iRBCs). Perceiving these cues is sufficient to produce the initial delay in transmission investment, but the delay is longer than parasites sensing time as the cue, and there is no increase in investment near the end of infection (Fig 2A, right panel). Given that parasite densities rise non-linearly at the start of infections (both simulated and real [11,42–45]), sensing

logged rather than raw densities results in reaction norms that are more sensitive to those early changes (Fig 2B), enabling parasites to increase transmission investment after the densities have reached a rough threshold. In contrast, sensing raw parasite densities results in reaction norms that essentially ignore changes at the low parasite densities characteristic of early infections (Fig 2B). The density of host resources (RBCs; raw or logged) is also a poor cue for initial stages of infection, since they change very little early on and, therefore, do not allow parasites to engage in delayed transmission investment (Fig 2A). Overall, our results broadly suggest that early transmission delay necessitates sensitivity to changes at low parasite density.




Fig 2. Cue perception mediates parasite fitness and dynamics of transmission investment. (A) On the left, fitness values achieved by parasites adopting the optimal transmission investment strategy for each cue, normalized with respect to the time-varying strategy, are displayed in descending order. On the right, the dynamics of transmission investment, given the optimal reaction norms, are plotted. Investment varies from dark blue (low values) to yellow (high values). The first four cues after time, which include measures of parasite density on a $\log_{10}$ scale, enable parasites to obtain the highest fitness and delay in transmission investment. (B) Optimal reaction norms for different cues. Each facet displays the reaction norms for raw (black, dotted lines) or log-transformed (red, solid lines) cues. The ranges of cue values “experienced” by a parasite adopting the optimal reaction norm are displayed above (raw) or below (log-transformed) the reaction norms as rug plots (horizontal bars). Based on the shape of the reaction norms, we conclude that sensing log-transformed parasite-centric cues enables more sensitive detection of (and responses to) the initial rise in parasite abundance.

Dual cue perception is necessary for terminal investment

We found that parasites sensing a single cue cannot reproduce the expected optimal transmission investment strategy, particularly the rapid increase in transmission investment during late infection (i.e., terminal investment). Achieving terminal

investment presumably requires sensing a cue that provides a unique signal at the later stages of infection. Inspecting infection dynamics produced by parasites that sense time revealed that only RBC density provides such a distinctive late-stage signal (Fig 3A). Given the poor performance of this cue in early stages of infection, these results suggest that simultaneous perception of RBC and a parasite-centric cue could provide infection stage-specific information that enables both early reproductive restraint and terminal investment. Consistent with this expectation, in a rodent malaria experiment, including variables that reflect parasite density and host resources (specifically, replication rate and changes in RBC density) in statistical models explained substantially more variation in transmission investment than the parasite-centric variable alone [13].

To investigate the fitness consequences of sensing two cues, we optimized the transmission investment strategy of parasites sensing all pairwise combinations of log-transformed cues, using two-dimensional spline surfaces and focusing on combinations of log-transformed cues given their higher performance in the single cue optimizations. Our results suggest that sensing dual cue combinations improves parasite fitness relative to sensing a single cue (Fig 3B), with parasites sensing logged RBC and logged iRBC (asexual or total) exhibiting notably higher fitness (Fig 3B). This higher fitness is conferred by transmission investment patterns that are close to the time-varying ideal, with an initial delay in transmission investment and a late infection (nearly) terminal investment before returning to a low transmission investment (Fig 3C, S2 Fig). Increasing the spline flexibility of the single cue models (to match the degrees of freedom in dual cue models) could not reproduce this same pattern (Fig 3C), confirming that dual cue perception specifically enables terminal investment and its associated fitness benefits (S3 Fig).

To better understand the optimal patterns of transmission investment, we next analyzed the reaction norm of the dual cue combination that conferred the highest fitness: logged asexual iRBC and logged RBC. The pattern of transmission investment is defined by a looping journey through the reaction norm surface (Fig 3D), confirming that the cue combination traces a unique path over the course of an infection. We categorized the reaction norm into four distinct sections/stages (Fig 3D). Stage I corresponds to the initial phase of infection, characterized by low transmission investment, low parasite density, and high RBC density (Fig 3C-D). As the infection progresses, the reaction norm enters stage II, characterized by asexual iRBC production and RBC decline. During this stage, the optimal transmission investment increases considerably, followed by subtle declines as the iRBC population continues to increase (Fig 3C-D). The transition from a low to high transmission investment observed as stage I transitions into stage II constitutes delayed transmission investment. Following peak parasite density, the reaction norm enters stage III, which is marked by decreasing iRBC population and gradual replenishment of RBCs, signaling imminent infection clearance. At this stage, transmission investment increases to peak levels (i.e., terminal investment). Finally, as the infection nears clearance and the reaction norm enters stage IV, transmission investment sharply drops as the host heads towards recovery (Fig 3C-D). We note that this drop in predicted transmission investment is likely a consequence of the constraints of the spline surface, and different strategies during this stage would have negligible impact on fitness [23]. Overall, our results suggest that each cue in this combination plays a crucial role in defining progression through infection stages I to III, with asexual iRBC density exhibiting the largest variation in early and late infection (stages I and III), and logged RBC density exhibiting the largest variation during mid-infection (Stage II; Fig 3D). This temporal staggering of iRBC and RBC dynamics (i.e., their hysteretic relationship) ensures that these infection stages can be




Fig 3. Dual cue perception increases parasite fitness and enables terminal investment. (A) RBC density uniquely marks the onset of terminal investment. In the left panel, we plot transmission investment of parasites sensing time (i.e., the ideal dynamics) and mark its key features: transmission delay (purple) and terminal investment (red). We then plot the dynamics of single cues on the right and mark the range of cue values reached during terminal investment with two red, dashed lines. The overlap of the red dashed lines with earlier cue values indicates that parasites may not be able to uniquely recognize the end of the infection. (B) The fitness of parasites (normalized with respect to the time-varying strategy) adopting the optimal strategy for each dual cue combination is indicated with a blue triangle, while the fitness obtained when perceiving the best performing single cue (of the combination) is represented with an orange circle. To obtain the fitness of a fair comparator time-varying strategy, we optimized transmission investment as a function of time using a spline with nine degrees of freedom (the same degrees of freedom used in the dual cue model), rather than the three degrees of freedom used above and previously [23]. (Note that the marginal fitness gains of using more flexible splines are small [23], but tracking two environmental cues necessitates more degrees of freedom.) (C) The pattern of transmission investment over time for parasites sensing the best dual cue combination (R log & I log, blue square) along with individual components of the cue (I log, orange circle; R log, yellow triangle; each optimized with nine degrees of freedom) and the ideal dynamic (Time, black cross) are plotted together. Parasites sensing the combination of logged asexual iRBC and logged RBC best approximate the time-varying strategy. (D) Transmission investment (colour of heatmap) is plotted as a two-dimensional reaction norm, with logged RBC and logged asexual iRBC as cues. The white line indicates the trajectory of infection, starting at the top left, for parasites adopting this optimal reaction norm, with each point representing a single day. The arrow on the white line indicates the passage of time. For (C-D), the four stages outlined in the results section are annotated with I, II, III, and IV.

tracked. As a result, dual cue perception provides parasites with the ability to “state differentiate,” to accurately monitor the progress of the infection and adjust transmission investment accordingly.

Intriguingly, the dual cue combinations that are most informative for parasites making life history decisions (top two in Fig 3B) are also the most informative for predicting patient outcomes in clinical malaria infections [46]. In both cases, two slightly out-of-phase within-host measures can uniquely define a patient’s (or simulated host’s) position on the path from health through sickness to recovery. Plotting this trajectory through disease maps (i.e., the two measures plotted against each other, as in Fig 3D) produces a looping, or hysteretic, pattern, with each infection stage occupying a unique coordinate in this two-dimensional space [46]. To further validate that forming the characteristic looping pattern is a unique feature of our identified best cue combinations, we constructed disease maps using experimental data from laboratory mice infected by a single strain of *P. chabaudi* [11, 42–45, 47] and simulated data from our models where parasite senses time. Both experimental and simulated data indicate that raw or logged RBC with logged iRBC (asexual or total in the model; only total is available in the experimental data) are the only cue combinations that generate fully separated loops (e.g., Fig 4A), while other cue combinations do not (e.g., Fig 4B and C; full data shown in S4 Fig and S5 Fig). When early and late infection stages are “condensed” in space, simulated parasites fail to state differentiate and do not delay transmission investment or terminally invest (S2 Fig). In summary, our results demonstrate that optimal transmission investment requires parasites to unambiguously state differentiate, track infection stages, and dynamically adjust investment in response to changing conditions. Our data suggest that perceiving hysteretic cues — of which a combination of RBC and logged iRBC provides a clean delineation of different infection stages — is a viable method of attaining state differentiation.

Dual cue perception exacerbates the fitness loss incurred by environmental and developmental stochasticity

The benefits of plasticity depend on the quality of information organisms obtain from environmental cues. Adaptive phenotypic plasticity is contingent on proper phenotype–environment matching, where the perceived cue is a reliable predictor of the optimal phenotype [48–52]. Theory suggests that a weak correlation between cue(s) and the optimal phenotype (i.e., low cue reliability) could disfavour plasticity relative to other strategies (i.e., local adaptation [49, 50, 53], canalization [52], bet-hedging [53]). For malaria parasites, there may be unpredictable variation in the environment (e.g., RBC availability [41, 54, 55], immunity [41, 56–58]) or in traits governing parasite development within the host [41]. Such environmental and developmental stochasticity could preclude evolution towards the strategies we predict above. Although our model allows parasites to sense components of their state, and their resulting level of transmission investment will feedback to influence that state, the results above assume that all else is equal between hosts. Specifically, the optimal reaction norms identified above are associated with a particular set of model parameters (the median estimates from posterior distributions in [41]; we refer to this as the deterministic model). This implicitly assumes that hosts are all the same or that parasites are moving between hosts sufficiently quickly that they adapt to an “average” environment.

To investigate how biologically relevant variation affects cue performance, we built a semi-stochastic model based on the work of Kamiya *et al.* [41], which employed a hierarchical Bayesian method to quantify inter-host variation in parasite burst size,




Fig 4. Trajectories of log-transformed RBC and iRBC densities form a looping pattern *in silico* and *in vivo*. Simulated (left) and experimental (right) trajectories of cues that (A) enable transmission delay and terminal investment or (B-C) fail to achieve these patterns. For each panel, the left graph displays the results from our simulations where parasite senses time with nine degrees of freedom, with the distance between each circle representing a single day and the colour of the line representing the observed transmission investment. The right graph displays the infection trajectory of laboratory *P. chabaudi* infections in mice, with the colour of the lines indicating the day post-infection. Note that asexually- and sexually-committed iRBCs were not distinguished in the experimental data. For the simulated trajectories, the first 3 days correspond to the period of delayed transmission investment and the last 5 days correspond to when terminal investment is expected. In (B), early changes in asexual iRBC density cannot be clearly separated (see the red, dotted box), therefore preventing a delayed transmission investment strategy. Similarly, in (C), late state infection stages cannot be separated from earlier time points, preventing parasites from terminally investing.

erythropoiesis (RBC replenishment), and immune strengths. For all log-transformed

single cues and dual cues, we performed 1000 20-day simulations, holding the reaction norm at the optimum from the relevant deterministic simulation, and simultaneously varying the RBC replenishment rate (ρ), parasite burst size (β), immune activation strengths (ψ_n and ψ_w) and immunity half-lives (ϕ_n and ϕ_w), sampling parameter values from the reported posterior distributions [41](S6 Fig). Each choice of parameter set will influence the within-host infection dynamics, potentially altering the trajectory of the perceived cue(s), resulting in different, expressed patterns of transmission investment and, thus, fitness. Here we are asking, if parasites are “optimized” to the average host, does environmental and developmental variation change inferences about which cues are best? We quantified the effect of variation on cue performance using the geometric mean fitness, commonly used to assess fitness in fluctuating environments [59,60].

We find that the geometric mean fitness of parasites responding to a given cue, in the face of this variation, is largely uncoupled from fitness in the deterministic model (i.e., the environment in which the strategy was optimized; Fig 5A), emphasizing that the “best” reaction norm for a certain environment does not always confer the best phenotype in another. Interestingly, all dual cues have a lower geometric mean fitness than the best-performing single cue (I log, Fig 5A). Furthermore, this result emphasizes the challenge of using time as an environmental proxy, since its geometric mean fitness in the semi-stochastic model is among the bottom half of cues. Allowing only one parameter to vary similarly tends to reduce the fitness of the best dual cue strategies disproportionately (S7 Fig). These results indicate a potential tradeoff between fine-tuning fitness in a given environment and susceptibility towards environmental and developmental stochasticity.

While sensing two cues provides more information about “where” an infection is, it could also increase the amount of perceived noise and exacerbate the negative effects of environmental/developmental stochasticity. To test whether increased noise perception reduces the fitness of dual cue-sensing parasites, we compared the trajectories of the most “robust” single cue (I log; Fig 5B left panel) and the dual cue combination with the highest deterministic model fitness (R log & I log; Fig 5B right panel). As expected, the variation in asexual iRBC density across our simulations is much lower than that of both asexual iRBC and RBC (Fig 5B top panels). While parasites sensing both types of cues delay transmission, those sensing logged RBC and iRBC display highly variable transmission investment dynamics (Fig 5B bottom right). In contrast, the transmission investment pattern of parasites sensing only logged asexual iRBC resembles that of the deterministic model (Fig 5B bottom left). These results suggest that sensing certain dual cue combinations can reduce parasite fitness by increasing noise perception that induces suboptimal transmission investment.

Dual cue perception leads to higher virulence

Given that cue perception mediates infection dynamics, it could also influence the severity of infections, or virulence. We estimated virulence using the methods described in Torres *et al.* [46], where infection dynamics are mapped in disease space, with the y-axis quantifying host health (RBC density) and the x-axis quantifying parasite load (total iRBC density). A more virulent infection is expected to produce a higher parasite load while consuming more host resources, tracing a larger loop through disease space. To evaluate the effects of cue perception on virulence, we generated 30-day disease maps of infections, allowing sufficient time for the infection to fully resolve in our model. We use the area enclosed by each disease map as a proxy for virulence, with larger areas indicating higher virulence. We note that we are using “virulence” as synonymous with infection severity and not host mortality, thus we assume no link between virulence and




Fig 5. Optimal transmission investment strategies do not always translate effectively across different environmental and developmental conditions. (A) Fitness of parasite strategies in the deterministic model is largely uncoupled from the geometric fitness of those same strategies in the semi-stochastic model. Plotted on the y axis is the geometric mean fitness of 1000 simulations of parasites adopting the optimal transmission investment strategy (sensing time from the deterministic model) using a semi-stochastic model where six parameters are randomly drawn from posterior distributions derived by Kamiya *et al.* [41]. The fitness from the deterministic model is plotted on the x-axis and all values are normalized with respect to the time-varying strategy with nine degrees of freedom (i.e., the dual cue comparator shown in Fig 3C). Note we also plot the fitness of the time-varying strategy with three degrees of freedom. Overall, dual cue-sensing parasites tend to perform worse than single cue-sensing parasites in the face of variation. The cue with the highest geometric fitness (I log) is indicated by a dotted orange box, while the cue combination with the highest deterministic model fitness (R log & I log) is bound by a blue dotted box. (B) Sensing dual cues can increase noise perception and lead to highly variable transmission investment dynamics. The top two graphs display the cue trajectories for parasites sensing the best performing cue in the semi-stochastic simulations (logged asexual iRBC density, plotted against time; left) or dual cues (logged asexual iRBC and logged RBC density, plotted against each other; right). The red lines represent individual semi-stochastic simulation results, while the thicker white line represents the dynamics from the deterministic model. The background colour indicates the optimal reaction norm: on the right, this is the same as what is plotted in Fig 3D. The bottom two graphs display the transmission investment dynamics of the same parasites. The thicker orange/blue line represents the transmission investment trajectory from the deterministic model, while the thinner grey lines represent individual semi-stochastic simulation results.

the duration of infection. Recent modeling suggests that resource limitation imposes a more important constraint on the evolution of transmission investment in *P. chabaudi* than does host mortality due to infection [61].

Parasites perceiving only a single cue trace smaller disease maps than parasites that adopt the ideal pattern of transmission investment (by sensing time; Fig 6A). Additionally, the area of the disease map is negatively correlated with parasite fitness

across all single cues (Fig 6B), suggesting that if parasites are constrained to perceive a single within-host cue, strategies that result in lower virulence would be favoured. Parasites sensing the top two dual cue combinations produce disease maps that are similar to the map traced by parasites adopting the time-varying strategy (Fig 6C) and result in higher virulence relative to parasites sensing a single cue (Fig 6D; compare x-axis values of points in B and D). Overall, these findings suggest that if parasites have limited information or have evolved to respond to only one cue (e.g., because of physiological constraints or high environmental/developmental variation), virulence will be lower, while if parasites are favoured to more finely track infection stages through dual cue perception, they will exploit host resources to a greater extent and infections will be more virulent.




Fig 6. Cue perception mediates infection severity (virulence). (A, C) display the 30-day disease maps of infections where parasites adopt the top two optimal transmission investment strategies when sensing (A) a single cue or (C) dual cues. We use the area of the map traced as a proxy for virulence [46]. The trajectory of parasites that adopt the ideal transmission investment strategy (sensing time) is shown in black. (B, D) Relationship between disease map area and fitness (normalized with respect to the time-varying strategy) for parasites that sense (B) only one cue or (D) dual cues. The top two cues for single and dual cue perceiving parasites are coloured, while the grey points represent the other (B) single or (D) dual cues. Overall, these results suggest that perceiving only a single cue generates selection for strategies that lead to lower virulence while perceiving two cues generates selection for strategies that lead to higher virulence.

Discussion

Just as with any sexually-reproducing species, investing in reproduction by producing gametocytes is critical for the fitness of *Plasmodium* spp., as this is the only parasite stage capable of being transmitted to mosquito vectors. As for other species, this investment comes with tradeoffs for malaria parasites: a given infected host cell can either produce multiple asexual progeny that can infect new host cells and have the potential to grow or maintain a given infection, or a single, sexually-differentiated gametocyte with the potential to “reproduce” an infection [62]. Thus, maximizing parasite fitness requires balancing these competing demands, the relative importance of which is likely to shift over the course of infection, as the within-host environment changes in response to parasites exploiting the host. Indeed, data show that transmission investment varies throughout infections [11, 12, 26] and appears to change in response to drug treatment in ways predicted to be beneficial for parasites (e.g., [13, 40]). While adaptive plasticity is a well-known solution to the challenge of living in a changing environment (e.g., [63]), there has been skepticism about the fitness benefits of observed parasite plasticity [64], in part because knowledge of parasite environmental sensing mechanisms remains limited. Previous mathematical modeling has treated parasites as either insensitive to environmental change (i.e., constant transmission investment [21, 32–36, 65, 66]) or as perfectly informed (time-varying investment [21, 23, 24]). We instead asked what environmental cues malaria parasites *should* use to set transmission investment, demonstrating that no single cue allows parasites to achieve the best time-varying strategy [23]. Parasites must sense two cues to recapitulate the optimal strategy but that strategy enhances the risk of parasites being “tricked” by environmental and developmental stochasticity into expressing suboptimal phenotypes. We further show that if parasites have the capacity to sense more aspects of their within-host environment, then this has negative consequences for host health.

Previous theoretical work identifies two key features of optimal transmission investment strategies for *P. chabaudi* [23]. First, there should be no investment at the start of infections, when densities are too low to permit successful infection of vectors (i.e., transmission investment delay). Second, parasites should invest all resources into transmission just before an infection ends, a “terminal investment” strategy from life history theory [7]. Our results delineate the environmental cues that can recapitulate these features. Transmission delay requires a sharp, threshold-dependent transition from no investment to an intermediate investment level as the parasite population grows. Making clear-cut decisions in response to continuous cues is common in biological systems (e.g., cell cycle commitment [67] and developmental patterning [68]) and often relies on ultrasensitivity, characterized by a steep response curve. Our results demonstrate that an ultrasensitive reaction norm where transmission investment rises sharply after a certain population size threshold—a pattern enabled by sensing parasite-centric cues on a logarithmic scale—is sufficient for transmission delay. Prior empirical studies have found that *Plasmodium* transmission investment varies in response to conspecific density [11, 12], parasite-conditioned media [14, 15], and infected red blood cell (iRBC)-derived vesicles [16], suggesting that parasites possess the ability to sense iRBC density. Our results support these responses being part of an adaptive strategy.

In addition to parasite density, *P. chabaudi* has been shown to modulate its transmission investment based on RBC availability and age structure [13, 17–19, 24]. Our results suggest that this is also part of an adaptive strategy, with the combination of iRBC and RBC cues allowing parasites to distinguish between early and late infection. Since most within-host variables go up and down, or vice versa, over time

(e.g., iRBC densities rise and then fall due to resource limitation or immunity), one cue cannot provide enough detail to unambiguously establish “where” a parasite is in the course of infection. Combining cues solves this problem. In line with this logic, researchers seeking to define a host’s trajectory over time from health through sickness and on to recovery [69] found a combination of two within-host measures (e.g., parasite density and resource abundance) that are slightly out of phase with each other can uniquely determine a patient’s position on their path for malaria infection [46]. Remarkably, though perhaps predictably, we found that the information desired by medical professionals seeking to predict patient outcomes is also the most informative for parasite life history decisions.

There is a growing appreciation that, across diverse organisms, plasticity is often expressed in response to multiple cues that allow for more finely-tuned strategies [70–72]. Our work adds support to theories suggesting that the adaptive benefits of phenotypic plasticity are enhanced when cues provide non-redundant information [73] and enable organisms to sense distinct states that correspond to specific optimal phenotypes [74]. Yet it is also widely appreciated that integrating multiple cues presents challenges, including the additional metabolic burden of sensing another cue [75] and the multiplicative effect of noise when cues are non-additive [72]. Our complex (decidedly non-additive) dual cue reaction norms present a trade-off: they allow parasites to more accurately recreate optimal strategies in a given environment while rendering them more susceptible to stochastic environmental and developmental change, i.e., host-to-host variability. Such tradeoffs may represent important constraints on the evolution of plastic reproductive investment strategies, perhaps explaining why *P. chabaudi* rarely exhibits *true* terminal investment *in vivo* (displaying only modest increases in transmission investment near infection resolution; [13, 20]), and on the evolution of cue perception more broadly.

The tradeoff hypothesis for virulence evolution stipulates that pathogen transmission inevitably harms the host [76]. While the generality of this framework has been contested (see [77] and references within), empirical data from *Plasmodium* spp. have generally supported a positive relationship between virulence and transmissibility [78–81]. Here, we demonstrate that if parasites are limited to sensing a single cue for setting transmission investment, then the strategies that result in the most onward transmission actually result in less host exploitation. Allowing parasites to sense two cues reverses this relationship and produces the expected pattern of the most onward transmission being achieved by the strategies that also produce the most virulent infections. Importantly, previous work has shown that host heterogeneity can constrain parasite virulence evolution [82–84] and our results suggest another mechanism by which this may be true: responding to two cues—while allowing for state differentiation in one host environment—can dramatically reduce fitness in the face of heterogeneity, potentially favouring single cue (hence, lower virulence) strategies.

Mathematical modeling has enabled us to explore the fitness consequences of cue perception in *P. chabaudi*, and our results suggest where greater model complexity would be most or least informative. For example, incorporating variation in gametocyte circulation time [26] would undercut the reliability of gametocytes as a cue, further supporting our results that it is unlikely to represent the optimal cue. The human malaria parasite *P. falciparum* has a longer sexual development period of 9–12 days [85] and this longer lag between cue perception and phenotype expression will reduce cue reliability [86]. Thus, we anticipate the value of sensing gametocytes to decrease even further in this species. While we have ignored parasite developmental processes that

occur in the vector, recent theory suggests they have little qualitative impact on optimal transmission investment strategies [87]. In contrast, the impact of previous exposure on host immunity represents one area where more complexity could qualitatively alter predictions. Our models depict immunologically naive hosts, but where malaria is endemic, parasites will often find themselves in a host with a prior history of infection. Understanding how immune memory influences the quality and reliability of potential within-host cues represents an important area for future research. Our selection of cues assumes that parasites sense only the magnitude of a cue, without detecting temporal changes in that magnitude. Schneider *et al.* [13], in their empirical demonstration of *P. chabaudi* adaptively altering transmission investment, found that replication rate, i.e., the proportional change in parasite densities, had better explanatory power than absolute densities. Broadly speaking, they found transmission investment increased with replication rate, a pattern that is not found with our model since replication rate is highest at the start of infections, before resource or immune limitation kicks in, and not investing in transmission at this point is best. We note that a direct comparison of results is tricky since estimates of transmission investment underlying their observed pattern come from day nine of infections or later. Nonetheless, we identified sensitivity to changes at low population density as a key feature of a “useful” cue and the proportional changes captured by replication rate will similarly share this feature. Another important model assumption is that all parasites perceive the same cue at the same strength, but cues may not be spatially uniform. Variation in signal strength and imperfect cue sensing could lead to heterogeneity in transmission investment in a genotypically uniform parasite population, similar to the dynamics observed in bacterial biofilms [88]. Extrapolating from our results, we hypothesize that if the variation in signal strength is large enough, selection may favour single cue sensing due to the relative robustness to stochastic variation. We note that the variation we incorporated in our semi-stochastic simulations likely underestimates that which would be observed in natural infections, since the dataset fitted by Kamiya *et al.* [41] includes only a single mouse host genetic background studied in a highly controlled experimental system [29]. Thus, environment-phenotype mismatching could be an even larger problem in nature than we have estimated. Additionally, given the variation in burst size and developmental timing across *Plasmodium spp.*, there is a great need to quantify the extent of variation across infection parameters to better understand the role of environmental/developmental heterogeneity on parasite evolution.

Beyond malaria, our work presents a generalizable computational approach to study phenotypic plasticity. By defining plastic traits as an optimizable reaction norm, it is possible to explore the fitness consequences of sensing different cues and generate hypotheses regarding why organisms respond to certain cues and not others. Mechanistic models of plasticity that capture feedbacks between traits and the environments that shape them will allow for predicting the consequences of perturbations, which is relevant for understanding not only disease dynamics in the face of control measures, but also the fate of organisms whose environment is rapidly altered by climate change.

Materials and methods

Within-host model of *P. chabaudi* infection dynamics

We modified and extended a previously published mathematical model describing the within-host infection dynamics of the rodent malaria parasite, *P. chabaudi* [23] (Fig 1).

The original model consists of ordinary and delay differential equations representing the interactions between the parasite, host red blood cells (RBCs), and host immunity over a 20-day infection. There are three key differences in our model from that previous work. First, we include the activity of two forms of immunity to better capture host defense mechanisms: a targeted immune response that only kills infected RBCs (iRBCs) and an indiscriminate response that kills any RBC, as in [41]. Second, we revised the model to account for the observation that sexually-committed parasites do not engage in gametocytogenesis until the next infection cycle (Next Cycle Conversion; [39]). This necessitated separately tracking the production of sexual merozoites (M_g) and asexual merozoites (M) from infected cells and consequently introduced a delay between sexual commitment and the outcome of sexual commitment (i.e., production of an iRBC that will produce a gametocyte, I_g). Third, we modeled transmission investment as a function of one or two within-host cues, instead of time as in [23], to better capture the mechanism of phenotypic plasticity, since there is no evidence malaria parasites track time across multiple generations. We further describe the implications of these changes on the model below.

We depict an immunologically naive host given the availability of experimentally derived parameters. Throughout an infection, we assume indiscriminate immunity eliminates all RBCs (R , I , I_g) whereas targeted immunity eliminates only iRBC (I , I_g). As in [41], we define N and W as the proportion of cells cleared per day by indiscriminate and targeted responses, respectively. The magnitude of this activity is dependent on the total abundance of iRBC ($I + I_g$):

$$\frac{dN}{dt} = \underbrace{\psi_n \frac{I(t) + I_g(t)}{\iota} (1 - N(t))}_{\text{Immune activation}} - \underbrace{\frac{N(t)}{\phi_n}}_{\text{Immune decay}} \quad (1)$$

$$\frac{dW}{dt} = \underbrace{\psi_w \frac{I(t) + I_g(t)}{\iota} (1 - W(t))}_{\text{Immune activation}} - \underbrace{\frac{W(t)}{\phi_w}}_{\text{Immune decay}}, \quad (2)$$

where ψ_n and ψ_w denote the immune activation strengths, ϕ_n and ϕ_w represent the immune activity half-lives, and ι represents the theoretical maximum total iRBC density, which determines the threshold at which immunity saturates. Experimental data from mouse infections indicate that indiscriminate immunity is quickly activated but short-lived, whereas targeted immunity has a longer activation period but a longer half-life [41]. Because N and W have been defined as proportions of cells cleared, they can be rewritten to reflect daily rates of clearance. For example, $N = 1 - e^{-x_N}$, where x_N is equal to the rate of clearance on a given day, and is defined as $-\ln(1 - N)$, with some rearranging of terms. A similar expression provides the daily clearance rate for targeted killing, $-\ln(1 - W)$.

Mimicking an experimental *P. chabaudi* infection, the host receives an initial dose of asexually-committed iRBC (I_0) to start the infection. We assume no parasites in this initial dose commit to gametocytogenesis and all produce asexual merozoites (M). Therefore, before the end of the first asexual development period ($t \leq \alpha$, where α is the development period of one day for this species), the dynamics of asexual iRBC (I) in the host are determined solely by the invasion of RBC by asexual merozoites, natural RBC death, maturation of the initial cohort of infected cells, and immune-mediated

killing such that

$$\frac{dI}{dt} = \underbrace{\epsilon R(t)M(t)}_{\text{RBC invasion}} - \underbrace{\mu I(t)}_{\text{Natural death}} - \underbrace{I_0 \text{Beta}(sp, sp)S}_{\text{iRBC maturation}} - \underbrace{[-\ln(1 - N(t)) - \ln(1 - W(t))]I(t)}_{\text{Indiscriminate and targeted immunity}}. \quad (3)$$

The first term, $\epsilon R(t)M(t)$, represents the production of iRBC through asexual merozoite invasion of RBCs, which occurs at a rate of ϵ per day, given contact. Infected RBCs die at a background rate of μ , while $-\ln(1 - N(t))$ and $-\ln(1 - W(t))$ describe their rate of death due to indiscriminate and targeted immunity, respectively. The infection-age structure of the initial cohort of asexual iRBCs, denoted as $\text{Beta}(sp, sp)$, is modeled using a Beta distribution with a shape parameter $sp = 1$, allowing iRBC maturation to occur with a uniform probability throughout the development period [89]. The uniform probability of maturation for the initial cohort is computationally efficient and provided extremely similar results to more narrow age distributions in a previous modeling study [23]. The rate of asexual iRBC maturation depends on an exponential survival function S , which describes the proportion of iRBC that survives until maturation. S is defined by a cumulative hazard function that is calculated by integrating the total rate of death across the development period:

$$S = \exp\left[-\int_{t-\alpha}^t \underbrace{\mu}_{\text{Natural death}} + \underbrace{-\ln(1 - N(\omega)) - \ln(1 - W(\omega))}_{\text{Indiscriminate and targeted immunity}} d\omega\right]. \quad (4)$$

During the period $t \leq \alpha$, asexual merozoite production is described by

$$\frac{dM}{dt} = \underbrace{\beta I_0 \text{Beta}(sp, sp)S}_{\text{iRBC maturation}} - \underbrace{\mu_m M(t)}_{\text{Natural death}} - \underbrace{\epsilon R(t)M(t)}_{\text{RBC invasion}}, \quad (5)$$

where the first term represents the production of asexually-committed merozoites by asexual iRBC that have survived the duration of the developmental cycle and burst to produce β merozoites each. Asexual merozoites are lost by natural death at a rate of μ_m and are removed by their invasion of susceptible RBC.

After all the initially injected iRBC have matured or died ($t > \alpha$), the next cohort of asexual iRBC starts to reach maturity. Transmission investment is defined as the proportion, c , of this new generation that commits to gametocytogenesis, resulting in the production of β sexually-committed merozoites (M_g) per cell. These sexual merozoites invade susceptible RBCs to produce sexual iRBCs (I_g). We assume that sexually-committed merozoites have the same death rate (μ_m) and rate of RBC invasion (ϵ) as asexual merozoites. The remaining proportion $(1 - c)$ of iRBC produces asexual merozoites and continues the asexual replication cycle. Altogether, when $t > \alpha$, the dynamics of iRBC and merozoites are as follows:

$$\begin{aligned} \frac{dI}{dt} = & \underbrace{\epsilon R(t)M(t)}_{\text{RBC invasion}} - \underbrace{\mu I(t)}_{\text{Natural death}} - \underbrace{[-\ln(1 - N(t)) - \ln(1 - W(t))]I(t)}_{\text{Indiscriminate and targeted immunity}} \\ & - \underbrace{\epsilon R(t - \alpha)M(t - \alpha)S}_{\text{iRBC maturation}} \end{aligned} \quad (6)$$

$$\begin{aligned} \frac{dM}{dt} = & \underbrace{\beta(1 - c)\epsilon R(t - \alpha)M(t - \alpha)S}_{\text{iRBC maturation (asexual)}} - \underbrace{\mu_m M(t)}_{\text{Natural death}} - \underbrace{\epsilon R(t)M(t)}_{\text{RBC invasion}} \end{aligned} \quad (7)$$

$$\frac{dM_g}{dt} = \underbrace{\beta c \epsilon R(t - \alpha) M(t - \alpha) S}_{\text{iRBC maturation (sexual)}} - \underbrace{\mu_m M_g(t)}_{\text{Natural death}} - \underbrace{\epsilon R(t) M_g(t)}_{\text{RBC invasion}}. \quad (8)$$

The parameter governing transmission investment (c) represents the proportion of iRBCs that produce sexually-committed merozoites and is further defined in the next section.

Before the first cohort of sexual iRBC matures ($\alpha < t \leq \alpha + \alpha_g$), the dynamics of sexual iRBC is governed solely by the rate of RBC invasion by sexually-committed merozoites, natural RBC death, and immunity:

$$\frac{dI_g}{dt} = \underbrace{\epsilon R(t) M_g(t)}_{\text{RBC invasion}} - \underbrace{\mu I_g(t)}_{\text{Natural death}} - \underbrace{[-\ln(1 - N(t)) - \ln(1 - W(t))] I_g(t)}_{\text{Indiscriminate and targeted immunity}}. \quad (9)$$

When the first cohort of sexual iRBC matures ($t > \alpha + \alpha_g$), each sexual iRBC produces a single gametocyte, G , such that

$$\begin{aligned} \frac{dI_g}{dt} = & \underbrace{\epsilon R(t) M_g(t)}_{\text{RBC invasion}} - \underbrace{\mu I_g(t)}_{\text{Natural death}} - \underbrace{[-\ln(1 - N(t)) - \ln(1 - W(t))] I_g(t)}_{\text{Indiscriminate and targeted immunity}} \\ & - \underbrace{\epsilon R(t - \alpha_g) M_g(t - \alpha_g) S_g}_{\text{iRBC maturation (sexual)}}, \end{aligned} \quad (10)$$

$$\frac{dG}{dt} = \underbrace{\epsilon R(t - \alpha_g) M_g(t - \alpha_g) S_g}_{\text{iRBC maturation (sexual)}} - \underbrace{\mu_g G(t)}_{\text{Natural death}}, \quad (11)$$

where μ_g is the background death rate of gametocytes. The survival probability of sexual iRBCs during their development cycle is defined by the exponential survival function S_g , which is similar to that of asexual iRBC (S), except the cumulative hazard function is integrated over the length of sexual iRBC development, α_g :

$$S_g = \exp\left[-\int_{t-\alpha_g}^t \underbrace{\mu}_{\text{Natural death}} + \underbrace{-\ln(1 - N(\omega)) - \ln(1 - W(\omega))}_{\text{Indiscriminate and targeted immunity}} d\omega\right]. \quad (12)$$

Throughout the infection, the density of uninfected RBCs, R , is affected by erythropoiesis, natural death, indiscriminate immunity, and merozoite invasion, such that:

$$\begin{aligned} \frac{dR}{dt} = & \underbrace{\mu R_0}_{\text{Erythropoiesis}} + \underbrace{\rho(R_0 - R(t))}_{\text{Erythropoiesis upregulation}} - \underbrace{\mu R(t)}_{\text{Natural death}} \\ & - \underbrace{[-\ln(1 - N(t)) R(t)]}_{\text{Indiscriminate immunity}} - \underbrace{\epsilon R(t) [M(t) + M_g(t)]}_{\text{Merozoite invasion}}. \end{aligned} \quad (13)$$

In the absence of parasites, RBCs are replenished at a baseline rate, μ , that maintains a homeostatic equilibrium RBC concentration (R_0). During infection, the rate of

erythropoiesis increases, and the extent of this upregulation is proportional to the deviation of RBC concentration from equilibrium [90], where ρ represents the maximum proportion of the RBC deficit recovered per day. RBCs are eliminated at a natural death rate of μ (which balances new production in the absence of infection), removed through indiscriminate immunity ($-\ln(1 - N(t))$), or converted to iRBC via invasion by merozoites.

We parameterized our model based on prior work on malaria intrahost dynamics [23, 41] (Table 2). Because parameter values in Kamiya *et al.* [41] depend on initial dosage (I_0), we estimated our starting dose using the same method, which corresponds to approximately 10^4 - 10^5 iRBCs. Preliminary investigations indicate that varying the initial iRBC dose did not lead to quantitative differences in parasite relative fitness. Consequently, we parameterized $\psi_n, \psi_w, \phi_n, \phi_w$ and ρ using the median posterior estimates from [41] assuming an initial dosage of 10^4 iRBCs.

Table 2. Parameter values used in the intrahost *P. chabaudi* model.

Parameter	Definition	Value	Unit	Source
R_0	Equilibrium RBC concentration	8.89×10^6	RBC/ μ L	[29]
μ	Background RBC production and RBC/iRBC mortality rate	0.025	day ⁻¹	[57]
p	Infection probability of RBC by merozoite invasion	8×10^{-6}	day ⁻¹	[91]
α	Asexual iRBC development period	1	day	[92]
α_g	Sexual iRBC development period	2	day	[93]
β	Burst size; number of merozoites produced per iRBC	5.721	merozoites	[94]
μ_m	Background merozoite mortality rate	48	day ⁻¹	[91]
μ_g	Background gametocyte mortality rate	4	day ⁻¹	[93]
I_0	Initial iRBC inoculum size	43.860	iRBC/ μ L	[23]
sp	Shape parameter dictating iRBC synchrony	1	NA	[89]
ψ_n	Activation strength for indiscriminate immunity	16.692	NA	[41]
ψ_w	Activation strength for targeted immunity	0.843	NA	[41]
ϕ_n	Half-life of indiscriminate immunity	0.035	day	[41]
ϕ_w	Half-life of targeted immunity	550.842	day	[41]
ι	Minimum iRBC concentration to induce maximum immunity	2.18×10^6	RBC/ μ L	[29]
ρ	Maximum proportion of RBC recovered per day	0.263	NA	[41]

Modeling transmission investment

In our model, transmission investment (c) is determined by a transformed cue variable, f_t , perceived α days ago by a parasite infecting a cell, to account for NCC ($f_t(t - \alpha)$). Following the approach taken by Greischar *et al.* [23], we modeled the transmission investment reaction norm as the complementary log-log basis-spline with three degrees of freedom (B_3). Briefly, this method models the reaction norm as a linear combination of basis function, generating a continuous, smooth cubic curve. The relative contribution of each basis function (also known as the model coefficient) defines the overall shape of the reaction norm (see Figure 1C). We further modify the basis-spline by applying a complementary log-log transformation to constrain the transmission investment value to be between zero and one. This method allows us to use numerical optimization to search for the set of model parameters (i.e., reaction norm shape) that maximizes parasite fitness. Because cubic splines are fundamentally smooth, the resultant reaction norms may have difficulty representing sharp transitions. For that reason, we used untransformed and log-transformed cue values to build our models and

capture greater variation in putative reaction norms.

For numerical optimization, the perceived cue must be within the bounds of the reaction norm. To achieve this, the perceived cue variable $f(t - \alpha)$ governing transmission investment is transformed via a linear combination of Heaviside step functions (H) from the R package *CRONE* [95] that constrains the cue value to be between zero and the maximum value (max) defined in Table 3. In summary, the transmission investment function is defined as

$$c[f_t(t - \alpha)] = \exp(-\exp(B_3[f_t(t - \alpha)])), \quad (14)$$

where the Heaviside-transformed environmental cue $f_t(t - \alpha)$ is given by

$$f_t(t - \alpha) = H[f(t - \alpha)]f(t - \alpha) + H[f(t - \alpha) - \max][\max - f(t - \alpha)]. \quad (15)$$

For the best time-varying reaction norm, we defined the perceived cue as the time since infection, which has a range of 0-20 days and a step size of 0.001 (same as the simulation time step). In this case,

$$c(t) = \exp(-\exp(B_3(t))). \quad (16)$$

To incorporate dual cue perception, we used the *mgcv* package [96] to model transmission investment as the response variable of a generalized additive model. This model incorporates a full tensor product smooth of two environmental cues with nine degrees of freedom, represented as follows:

$$c[f_{t1}(t - \alpha), f_{t2}(t - \alpha)] = \exp(-\exp(\sum_{i=1}^3 \sum_{j=1}^3 \zeta_{ij} B_{3,1}[f_{t1}(t - \alpha)] B_{3,2}[f_{t2}(t - \alpha)])), \quad (17)$$

where transmission investment is governed by two Heaviside-transformed cues, f_{t1} and f_{t2} , perceived α days ago. The transmission investment surface (i.e., 2-dimensional reaction norm) is generated by taking the tensor product of the basis-splines of two cues ($B_{3,1}$ and $B_{3,2}$). The shape of this spline surface is defined by the model coefficients ζ_{ij} .

Quantifying parasite fitness

To quantify parasite fitness, we calculated the cumulative transmission potential of different transmission investment strategies, following the method described by Greischar *et al.* [23]. For a given time step, transmission potential, $\tau(t)$, is defined as the probability that a mosquito vector becomes infected after biting an infected host and has been shown empirically to be a sigmoid function of gametocyte density in this parasite species [10],

$$\tau(t) = \frac{3.08 \times 10^{-6} G(t)^{1.56}}{1 + 3.08 \times 10^{-6} G(t)^{1.56}}. \quad (18)$$

The cumulative transmission potential, or overall fitness of the parasite, is calculated by integrating transmission potential over the entire infection duration of δ days:

$$\text{fitness} = \int_0^\delta \tau(\omega) d\omega. \quad (19)$$

This fitness metric assumes that the mosquito population is at a constant abundance and continuously biting, for the duration of the infection. Although this is a

simplification of the evolutionary pressures faced by malaria parasites [35], it enables us to focus our investigation at the within-host level, where transmission investment “decisions” are being made and environmental cues are being perceived by parasites. In our numerical simulations, we estimated fitness with a step size of 0.001 days unless stated otherwise.

Cue choice and perception

Transmission investment has been shown to vary in response to within-host competition [11, 12] and RBC age structure and availability [13, 17–19, 24], lysophosphatidylcholine level (a key component of the RBC plasma membrane) and exosome-like particles [97, 98]. Whether parasites are responding to these cues directly, or if they are instead a proxy for changes in some other within-host environmental factor is unclear. To capture some of the diversity of putative cues, we defined transmission investment as a function of one of ten different host and parasite-centric measures (Table 3). Additionally, we implemented dual cue models, allowing parasites to simultaneously perceive two different cues. As transmission investment in *P. falciparum* varies non-linearly with cue strength (e.g., *P. falciparum* in response to lysophosphatidylcholine concentration [97]), and within-host dynamics are known to vary only when infectious dose changes by orders of magnitude [29], we introduced non-linear cue perception, enabling parasites to perceive cues either unaltered or $\log_{10}$ -transformed. We excluded merozoite density as a cue due to its short half-life and the rapid pace of RBC invasion [99]. We also did not include measurements of immunity as cues, given that our model does not track individual components of immunity (rather, it tracks immune killing, phenomenologically), nor do we have sufficient empirical evidence detailing the relationship between different immunity components and transmission investment.

Table 3. Within-host environmental cues tested and their range.

Cue (density per μL)	Abbreviation	Min. value	Max. value
Asexual iRBC	I	0	6×10^6
Asexual iRBC $\log_{10}$	I log	0	6.78
Sexual iRBC	Ig	0	6×10^6
Sexual iRBC $\log_{10}$	Ig log	0	6.78
Asexual iRBC + sexual iRBC	Total I	0	6×10^6
Asexual iRBC $\log_{10}$ + sexual iRBC $\log_{10}$	Total I log	0	6.78
Gametocyte	G	0	6×10^4
Gametocyte $\log_{10}$	G log	0	4.78
RBC	R	10^6	10^7
RBC $\log_{10}$	R log	6	7

For all cues, we adopt a standardized range that spans the values typically observed when infecting immunologically naive mice with *P. chabaudi* [11], given by the minimum and maximum values in Table 3. All values that exceed the maximum bounds are “perceived” by the parasite as the maximum bound via the modified Heaviside transformation function.

Fitness optimization

Acute *P. chabaudi* infection tends to resolve within 20 days [100]. For all models, we therefore confined our simulations and optimizations to a 20-day infection period using

the parameters detailed in Table 2 and using the *deSolve* package in R [101]. To ensure the convergence and continuity of output dynamics, we simulated the models with a small time step size of 0.001 days using the vode solver [102]. For the dual cue model, we used a larger time step of 0.01 days (which does not greatly alter parasite dynamics but improves computational speed immensely).

To investigate how transmission investment strategies affect parasite fitness, we performed numerical optimization of the basis-spline (single cue) or generalized additive model coefficients (dual cue) to maximize the cumulative transmission potential. We optimized the models using two methods to increase the chances of obtaining the global fitness optima. First, we performed local optimization following the methodology outlined by Greischar *et al.* [23] with the same arbitrary initial values of 0.5 for all spline coefficients, but using the more efficient parallelized L-BFGS-B algorithm (Limited-Memory Broyden–Fletcher–Goldfarb–Shanno algorithm-B) implemented in the *OptimParallel* package [103]. Second, we optimized each model using a hybrid approach to avoid convergence to a local fitness maximum. This technique involved initial global optimization using the Differential Evolution (DE) algorithm from the *DEoptim* package [104] (maximum iterations = 500, step tolerance = 50, differential weighting factor = 0.8, cross-over constant = 0.5 (single cue) or 0.9 (dual cue), population size = 40 (single cue) or 90 (dual cue), time step = 0.01 (single cue) or 0.05 (dual cue)). DE is an efficient and robust algorithm with comparable/better performance relative to other stochastic optimization algorithms [105]. Following optimization by the Differential Evolution algorithm, we further optimized the objective function using the L-BFGS-B algorithm to pinpoint a probable fitness maximum. For the differential Evolution algorithm, we used a parameter bound of $(-5, -500, -500, -500)$ and $(5, 500, 5000, 500)$ for the single cue model and $(-10, -500, -1000, -1000, -250, -500, -1000, -500, -250)$ and $(10, 500, 1000, 1000, 250, 500, 1000, 500, 250)$ for the dual cue model. These bounds effectively encompass the majority of basis-spline flexibility. From the two optimization results, we selected the parameter sets that resulted in the highest fitness.

For dual cue models, we implemented an additional optimization method that relies on the single cue models to inform the initial starting point. For dual cue models, we reasoned that the best single cue strategy out of the two provides a theoretical baseline fitness. In other words, the worst a dual-cue strategy should do is as good as the best single cue (out of the two cues), while being unresponsive to the second cue. As such, we applied the L-BFGS-B algorithm to derive a dual cue strategy that “resembles” the best single cue strategy, specifically by minimizing an objective function with two terms: (i) the summed absolute difference between the dual and single cue reaction norms, and (ii) the total standard deviation in transmission investment across the worse cue. Inclusion of the second term ensures that transmission investment is not affected by variations in the worse cue. Together, the objective function is represented by:

$$\sum_{i=1}^{500} |\bar{c}_{i \text{ dual}} - c_{i \text{ single}}| + \sum_{i=1}^{500} \sum_{j=1}^{500} \sigma_{c_{ij \text{ dual}}} \quad (20)$$

We first discretized the cue ranges into 500 evenly spaced values ($i = 1 \dots 500$, better performing cue; $j = 1 \dots 500$, worse performing cue). For each cue value with the index i , we calculate the deviation of the dual cue model using the absolute difference between $\bar{c}_{i \text{ dual}}$, the dual cue transmission investment averaged over the worse cue, and $c_{i \text{ single}}$, the corresponding transmission investment in the single cue model. The objective function also quantified the extent to which the worse cue affects transmission investment, specifically by summing the standard deviation across the worse cue value

$\sum_{j=1}^{500} \sigma_{c_{ij} \text{ dual}}$ for each cue value indexed i .

Following this initial optimization, the resultant parameter is set as the starting point for an additional round of local optimization using the L-BFGS-B algorithm to maximize parasite fitness (same parameters as mentioned above).

As a further validation step, we compared the fitness of each putative single cue optimal strategy to the fitness obtained by 1000 randomly generated strategies. These random strategies were generated within the predefined bounds of $(-5, -500, -500, -500)$ and $(5, 500, 5000, 500)$, established previously to encompass much of the spline space. In no case was the putative best strategy outperformed by a random one (S1 Fig).

Semi-stochastic infection model

In order to explore the impact of individual infection-level variation on putative optimal transmission investment strategies, we incorporated the following random variables in the within-host model: maximum proportion of RBC deficit recovered per day (ρ), parasite burst size (β), immunity activation strengths (ψ_n and ψ_w), and immunity half-lives (ϕ_n and ϕ_w). For each iteration, random variables were drawn from 167 Markov chains (selected at regular intervals from a pool of 16,000 chains) used to estimate the posterior distribution in [41]. For the maximum proportion of RBC recovered per day and parasite burst size, the posterior distributions (originally centered at zero) are re-centered at the deterministic model values of 0.263 and 5.721, respectively. Since the paper from which we draw parameters explored the impact of dose [41], we only considered estimates pertaining to hosts infected with an initial dose of 10^4 parasites (constituting around 1000 parameter values), which corresponds to our choice for this model parameter (Table 2).

For each optimal transmission investment strategy (i.e., the best strategy for each cue or dual cue combination), we performed 20-day simulations with (i) all parameters allowed to vary from their deterministic default and (ii) only a single parameter varying. Here, we used a time step of 0.01 days, which does not change fitness substantially relative to performing the simulation with a time step of 0.001 days.

Disease mapping of laboratory mice infections


To explore the dynamics of host and parasite-centric cues in an *in vivo* setting, we curated data from publicly available sources of *P. chabaudi* infection dynamics in mice [11, 42–45, 47]. The curated dataset includes experiments that met the following criteria: a) single parasite strain infection, b) no administration of anti-malarial drugs, and c) availability of data on iRBC, RBC, and gametocyte densities. We then visualized two-dimensional disease maps (e.g., iRBC vs. RBC path plots) using the R package *ggplot2* [106].

Quantifying parasite virulence

We used the area encompassed within a disease map (where the x-axis displays total iRBC density and the y-axis displays uninfected RBC density, as in [46]) as a proxy for parasite virulence. To calculate this area, we first used the *chull* function [107] to define the convex hull, the smallest set of coordinates that encloses the points within the infection trajectory. We then used the *areapl* function from the *splanes* package [108] to quantify the area of the convex hull.

Supporting information

763



764

S1 Fig. Validation of transmission investment optimization. We conducted 1000 infection simulations for parasites sensing different cues, with each simulation using randomly generated transmission investment strategies. The fitness values of these simulations were compared to the fitness of parasites adopting the optimal transmission investment strategy. Each histogram represents the difference between the optimized fitness and the fitness obtained from the randomly generated strategy. The fitness of the random strategies is always lower.

765

766

767


768

769

770

771

772



773

S2 Fig. Transmission investment dynamics produced by parasites sensing different dual cue combinations. Transmission investment is plotted for all possible dual cue combinations, with heatmaps ranked in descending order (from top to bottom) based on their associated cumulative transmission potential.


774

775


776

777


778




S3 Fig. Terminal investment increases the fitness of dual cue-perceiving parasites. The cumulative transmission potential of parasites sensing time, the best performing dual cue combination (R log & I log), or the individual single cues (I log, R log) are indicated with a black cross, blue square, orange circle, and yellow triangle, respectively. The cumulative transmission potential is normalized with respect to the fitness of the best time-varying strategy. Note that time, the dual cue combination, and I log all permit an initial delay in transmission investment, while R log does not; the fitness advantage of this delayed strategy can be clearly viewed on these plots as fitness accrues faster. Only time and the dual cue combination lead to terminal investment, leading to additional fitness gains from day 18 onward.



S5 Fig. RBC/logged RBC and logged iRBC form a looping trajectory *in vivo*. To identify combinations of within-host variables that exhibit a looping relationship, we plotted all possible permutations of experimentally derived *P. chabaudi* variables. Each line represents the dynamics of a single strain of infecting parasite. The line colour indicates the progression of time.



S6 Fig. Posterior distribution of parameter values used for the semi-stochastic model. The histograms display the distribution of 1000 parameter values drawn from 167 Markov chains derived by Kamiya *et al* [41]. The parameter values used in the deterministic model are indicated by the dotted line.



S7 Fig. Perceiving the optimal dual cue combinations increases parasite susceptibility to environmental and developmental stochasticity. We conducted 1000 simulations, each lasting 20 days, in which parasites adopted the optimal transmission investment strategy (from the deterministic model) using a semi-stochastic model where only one parameter was randomly drawn from the posterior distributions (see S6 Fig). For each parameter that is varied (indicated in the facet panels), we plot the geometric mean of the fitness on the y-axis and the fitness from the deterministic model on the x-axis. All fitness values are normalized with respect to the best time-varying strategy. The dashed line represents a 1:1 relationship between the x-axis and the y-axis; the line is not visible on the plot for burst size, since variation in this parameter has the greatest negative influence on fitness.

Acknowledgments

We thank members of the Mideo Lab for useful feedback on the project.

References

1. Jones OR, Scheuerlein A, Salguero-Gómez R, Camarda CG, Schaible R, Casper BB, et al. Diversity of ageing across the tree of life. *Nature*. 2014;505(7482):169–173. doi:10.1038/nature12789.
2. Gadgil M, Bossert WH. Life Historical Consequences of Natural Selection. *The American Naturalist*. 1970;104(935):1–24. doi:10.1086/282637.
3. Clutton-Brock TH, Albon SD, Guinness FE. Maternal dominance, breeding success and birth sex ratios in red deer. 1984;308(5957):358–360. doi:https://doi.org/10.1038/308358a0.
4. McNamara JM, Houston AI. State-dependent life histories. *Nature*. 1996;380(6571):215–221. doi:10.1038/380215a0.
5. Cotter SC, Ward RJS, Kilner RM. Age-specific reproductive investment in female burying beetles: independent effects of state and risk of death. *Functional Ecology*. 2011;25(3):652–660. doi:10.1111/j.1365-2435.2010.01819.x.

6. Velando A, Drummond H, Torres R. Senescent birds redouble reproductive effort when ill: confirmation of the terminal investment hypothesis. *Proceedings of the Royal Society B: Biological Sciences*. 2006;273(1593):1443–1448. doi:10.1098/rspb.2006.3480.
7. Carter LM, Kafsack BFC, Llinás M, Mideo N, Pollitt LC, Reece SE. Stress and sex in malaria parasites: Why does commitment vary? *Evolution, Medicine, and Public Health*. 2013;2013(1):135–147. doi:10.1093/EMPH/EOT011.
8. Churcher TS, Bousema T, Walker M, Drakeley C, Schneider P, Ouédraogo AL, et al. Predicting mosquito infection from *Plasmodium falciparum* gametocyte density and estimating the reservoir of infection. *eLife*. 2013;2:e00626. doi:10.7554/eLife.00626.
9. Bell AS, Huijben S, Paaijmans KP, Sim DG, Chan BHK, Nelson WA, et al. Enhanced Transmission of Drug-Resistant Parasites to Mosquitoes following Drug Treatment in Rodent Malaria. *PLOS ONE*. 2012;7(6):e37172. doi:10.1371/journal.pone.0037172.
10. Huijben S, Nelson WA, Wargo AR, Sim DG, Drew DR, Read AF. Chemotherapy, within-host ecology and the fitness of drug-resistant malaria parasites. *Evolution*. 2010;64(10):2952–2968. doi:10.1111/J.1558-5646.2010.01068.X.
11. Pollitt LC, Mideo N, Drew DR, Schneider P, Colegrave N, Reece SE. Competition and the Evolution of Reproductive Restraint in Malaria Parasites. *The American Naturalist*. 2011;177(3):358–367. doi:10.1086/658175.
12. Cameron A, Reece SE, Drew DR, Haydon DT, Yates AJ. Plasticity in transmission strategies of the malaria parasite, *Plasmodium chabaudi*: environmental and genetic effects. *Evolutionary Applications*. 2013;6(2):365. doi:10.1111/EVA.12005.
13. Schneider P, Greischar MA, Birget PLG, Repton C, Mideo N, Reece SE. Adaptive plasticity in the gametocyte conversion rate of malaria parasites. *PLOS Pathogens*. 2018;14(11):e1007371. doi:10.1371/JOURNAL.PPAT.1007371.
14. Williams JL. Stimulation of *Plasmodium falciparum* gametocytogenesis by conditioned medium from parasite cultures. *The American Journal of Tropical Medicine and Hygiene*. 1999;60(1):7–13. doi:10.4269/ajtmh.1999.60.7.
15. Dyer M, Day Karen P . Regulation of the rate of asexual growth and commitment to sexual development by diffusible factors from in vitro cultures of *Plasmodium falciparum*. *The American Journal of Tropical Medicine and Hygiene*. 2003;68(4):403–409. doi:10.4269/AJTMH.2003.68.403.
16. Mantel PY, Hoang A, Goldowitz I, Potashnikova D, Hamza B, Vorobjev I, et al. Malaria-Infected Erythrocyte-Derived Microvesicles Mediate Cellular Communication within the Parasite Population and with the Host Immune System. *Cell Host & Microbe*. 2013;13(5):521–534. doi:10.1016/J.CHOM.2013.04.009.
17. Trager W, Gill GS. Enhanced Gametocyte Formation in Young Erythrocytes by *Plasmodium falciparum* In Vitro. *The Journal of Protozoology*. 1992;39(3):429–432. doi:10.1111/j.1550-7408.1992.tb01476.x.

18. Trager W, Gill GS, Lawrence C, Nagel RL. *Plasmodium falciparum*: Enhanced Gametocyte Formation *in Vitro* in Reticulocyte-Rich Blood. *Experimental Parasitology*. 1999;91(2):115–118. doi:10.1006/expr.1998.4347.
19. Reece SE, Duncan AB, West SA, Read AF. Host cell preference and variable transmission strategies in malaria parasites. *Proceedings of the Royal Society B: Biological Sciences*. 2005;272(1562):511–517. doi:10.1098/rspb.2004.2972.
20. Buckling A, Crooks L, Read A. *Plasmodium chabaudi*: Effect of Antimalarial Drugs on Gametocytogenesis. *Experimental Parasitology*. 1999;93(1):45–54. doi:10.1006/expr.1999.4429.
21. Koella JC, Antia R. Optimal Pattern of Replication and Transmission for Parasites with Two Stages in Their Life Cycle. *Theoretical Population Biology*. 1995;47(3):277–291. doi:10.1006/tpbi.1995.1012.
22. Reece SE, Ramiro RS, Nussey DH. Synthesis: Plastic parasites: sophisticated strategies for survival and reproduction? *Evolutionary Applications*. 2009;2(1):11–23. doi:10.1111/j.1752-4571.2008.00060.x.
23. Greischar MA, Mideo N, Read AF, Bjørnstad ON. Predicting optimal transmission investment in malaria parasites. *Evolution*. 2016;70(7):1542–1558. doi:10.1111/EVO.12969.
24. Birget PLG, Repton C, O'Donnell AJ, Schneider P, Reece SE. Phenotypic plasticity in reproductive effort: malaria parasites respond to resource availability. *Proceedings of the Royal Society B: Biological Sciences*. 2017;284(1860). doi:10.1098/RSPB.2017.1229.
25. Clutton-Brock TH. Reproductive Effort and Terminal Investment in Iteroparous Animals. <https://doi.org/10.1086/284198>. 2015;123(2):212–229. doi:10.1086/284198.
26. Eichner M, Diebner HH, Molineaux L, Collins WE, Jeffery GM, Dietz K. Genesis, sequestration and survival of *Plasmodium falciparum* gametocytes: parameter estimates from fitting a model to malariatherapy data. *Transactions of The Royal Society of Tropical Medicine and Hygiene*. 2001;95(5):497–501. doi:10.1016/S0035-9203(01)90016-1.
27. Greischar MA, Mideo N, Read AF, Bjørnstad ON. Quantifying Transmission Investment in Malaria Parasites. *PLOS Computational Biology*. 2016;12(2):e1004718. doi:10.1371/JOURNAL.PCBI.1004718.
28. Pianka ER, Parker WS. Age-Specific Reproductive Tactics. *The American Naturalist*. 1975;109(968):453–464. doi:10.1086/283013.
29. Timms R, Colegrave N, Chan BHK, Read AF. The effect of parasite dose on disease severity in the rodent malaria *Plasmodium chabaudi*. *Parasitology*. 2001;123(Pt 1):1–11. doi:10.1017/S0031182001008083.
30. Metcalf CJE, Graham AL, Huijben S, Barclay VC, Long GH, Grenfell BT, et al. Partitioning regulatory mechanisms of within-host malaria dynamics using the effective propagation number. *Science*. 2011;333(6045):984–988. doi:10.1126/SCIENCE.1204588.

31. Barclay VC, Råberg L, Chan BHK, Brown S, Gray D, Read AF. CD4+T cells do not mediate within-host competition between genetically diverse malaria parasites. *Proceedings of the Royal Society B: Biological Sciences*. 2008;275(1639):1171–1179. doi:10.1098/rspb.2007.1713.
32. McKenzie FE, Bossert WH. An integrated model of *Plasmodium falciparum* dynamics. *Journal of Theoretical Biology*. 2005;232(3):411–426. doi:10.1016/j.jtbi.2004.08.021.
33. Bousema T, Drakeley C. Epidemiology and infectivity of *Plasmodium falciparum* and *Plasmodium vivax* gametocytes in relation to malaria control and elimination. *Clinical Microbiology Reviews*. 2011;24(2):377–410. doi:10.1128/CMR.00051-10.
34. Cao P, Collins KA, Zaloumis S, Wattanakul T, Tarning J, Simpson JA, et al. Modeling the dynamics of *Plasmodium falciparum* gametocytes in humans during malaria infection. *eLife*. 2019;8:e49058. doi:10.7554/eLife.49058.
35. Greischar MA, Beck-Johnson LM, Mideo N. Partitioning the influence of ecology across scales on parasite evolution. *Evolution*. 2019;73(11):2175–2188. doi:10.1111/EVO.13840.
36. Early AM, Camponovo F, Pelleau S, Cerqueira GC, Lazrek Y, Volney B, et al. Declines in prevalence alter the optimal level of sexual investment for the malaria parasite *Plasmodium falciparum*. *Proceedings of the National Academy of Sciences*. 2022;119(30):e2122165119. doi:10.1073/pnas.2122165119.
37. Lensen A, Bril A, van de Vegte M, van Gemert GJ, Eling W, Sauerwein R. *Plasmodium falciparum*: infectivity of cultured, synchronized gametocytes to mosquitoes. *Experimental Parasitology*. 1999;91(1):101–103. doi:10.1006/expr.1998.4354.
38. de Boor C. A Practical Guide to Splines. 1st ed. No. 27 in Applied Mathematical Sciences. Springer New York; 2001.
39. Dixon MWA, Thompson J, Gardiner DL, Trenholme KR. Sex in *Plasmodium*: a sign of commitment. *Trends in Parasitology*. 2008;24(4):168–175. doi:10.1016/J.PT.2008.01.004.
40. Birget PLG, Greischar MA, Reece SE, Mideo N. Altered life history strategies protect malaria parasites against drugs. *Evolutionary Applications*. 2018;11(4). doi:10.1111/eva.12516.
41. Kamiya T, Greischar MA, Schneider DS, Mideo N. Uncovering drivers of dose-dependence and individual variation in malaria infection outcomes. *PLOS Computational Biology*. 2020;16(10):e1008211. doi:10.1371/JOURNAL.PCBI.1008211.
42. Huijben S, Sim DG, Nelson WA, Read AF. Data from: The fitness of drug-resistant malaria parasites in a rodent model: multiplicity of infection. Dryad. 2011;.
43. Huijben S, Nelson WA, Wargo AR, Sim DG, Drew DR, Read AF. Data from: Chemotherapy, within-host ecology and the fitness of drug-resistant malaria parasites. Dryad. 2013;.


44. Huijben S, Bell AS, Sim DG, Tomasello D, Mideo N, Day T, et al. Data from: Aggressive chemotherapy and the selection of drug resistant pathogens. Dryad. 2014;.
45. Huijben S, Chan BHK, Nelson WA, Read AF. Data from: The impact of within-host ecology on the fitness of a drug-resistant parasite. Dryad. 2018;.
46. Torres BY, Oliveira JHM, Thomas Tate A, Rath P, Cumnock K, Schneider DS. Tracking Resilience to Infections by Mapping Disease Space. PLOS Biology. 2016;14(4):e1002436. doi:10.1371/JOURNAL.PBIO.1002436.
47. Reece SE, Drew DR, Gardner A. Sex ratio adjustment and kin discrimination in malaria parasites. Nature. 2008;453(7195). doi:10.1038/nature06954.
48. Moran NA. The Evolutionary Maintenance of Alternative Phenotypes. The American Naturalist. 1992;139(5):971–989. doi:10.1086/285369.
49. Scheiner SM. The genetics of phenotypic plasticity. VII. Evolution in a spatially-structured environment. Journal of Evolutionary Biology. 1998;11(3):303–320. doi:10.1046/j.1420-9101.1998.11030303.x.
50. De Jong G. Unpredictable selection in a structured population leads to local genetic differentiation in evolved reaction norms. Journal of Evolutionary Biology. 1999;12(5). doi:10.1046/j.1420-9101.1999.00118.x.
51. Schlaepfer MA, Runge MC, Sherman PW. Ecological and evolutionary traps. Trends in Ecology & Evolution. 2002;17(10):474–480. doi:10.1016/S0169-5347(02)02580-6.
52. Reed TE, Robin SW, Schindler DE, Hard JJ, Kinnison MT. Phenotypic plasticity and population viability: The importance of environmental predictability. Proceedings of the Royal Society B: Biological Sciences. 2010;277(1699). doi:10.1098/rspb.2010.0771.
53. Tufto J. The evolution of plasticity and nonplastic spatial and temporal adaptations in the presence of imperfect environmental cues. American Naturalist. 2000;156(2). doi:10.1086/303381.
54. Antia R, Yates A, de Roode JC. The dynamics of acute malaria infections. I. Effect of the parasite's red blood cell preference. Proceedings of the Royal Society B: Biological Sciences. 2008;275(1641):1449–1458. doi:10.1098/rspb.2008.0198.
55. Birget PLG, Kimberley F Prior, Savill NJ, Steer L, Reece SE. Plasticity and genetic variation in traits underpinning asexual replication of the rodent malaria parasite, *Plasmodium chabaudi*. Malaria Journal. 2019;18(1):222. doi:10.1186/s12936-019-2857-0.
56. Haydon DT, Matthews L, Timms R, Colegrave N. Top-down or bottom-up regulation of intra-host blood-stage malaria: do malaria parasites most resemble the dynamics of prey or predator? Proceedings of the Royal Society B: Biological Sciences. 2003;270(1512):289–298. doi:10.1098/rspb.2002.2203.
57. Miller MR, Råberg L, Read AF, Savill NJ. Quantitative Analysis of Immune Response and Erythropoiesis during Rodent Malarial Infection. PLOS Computational Biology. 2010;6(9):e1000946. doi:10.1371/JOURNAL.PCBI.1000946.


58. Kamiya T, Davis NM, Greischar MA, Schneider D, Mideo N. Linking functional and molecular mechanisms of host resilience to malaria infection. *eLife*. 2021;10. doi:10.7554/ELIFE.65846.
59. Lewontin RC, Cohen D. On population growth in a randomly varying environment. *Proceedings of the National Academy of Sciences of the United States of America*. 1969;62(4):1056–1060. doi:10.1073/pnas.62.4.1056.
60. Gillespie J. Polymorphism in random environments. *Theoretical Population Biology*. 1973;4(2):193–195. doi:10.1016/0040-5809(73)90028-2.
61. Pak D, Kamiya T, Greischar MA. Proliferation in malaria parasites: How resource limitation can prevent evolution of greater virulence. *Evolution*. 2024;78(7):1287–1301. doi:10.1093/evolut/qpae057.
62. Schneider P, Reece SE. The private life of malaria parasites: Strategies for sexual reproduction. *Molecular and Biochemical Parasitology*. 2021;244:111375. doi:10.1016/J.MOLBIOPARA.2021.111375.
63. Pigliucci M. Evolution of phenotypic plasticity: where are we going now? *Trends in Ecology & Evolution*. 2005;20(9):481–486. doi:10.1016/j.tree.2005.06.001.
64. Kochin BF, Bull JJ, Antia R. Parasite Evolution and Life History Theory. *PLOS Biology*. 2010;8(10):e1000524. doi:10.1371/journal.pbio.1000524.
65. McKenzie FE, Bossert WH. The Optimal Production of Gametocytes by *Plasmodium falciparum*. *Journal of Theoretical Biology*. 1998;193(3):419–428. doi:10.1006/jtbi.1998.0710.
66. Mideo N, Day T. On the evolution of reproductive restraint in malaria. *Proceedings of the Royal Society B: Biological Sciences*. 2008;275(1639):1217–1224. doi:10.1098/RSPB.2007.1545.
67. Charvin G, Oikonomou C, Siggia ED, Cross FR. Origin of Irreversibility of Cell Cycle Start in Budding Yeast. *PLOS Biology*. 2010;8(1):e1000284. doi:10.1371/journal.pbio.1000284.
68. Melen GJ, Levy S, Barkai N, Shilo B. Threshold responses to morphogen gradients by zero-order ultrasensitivity. *Molecular Systems Biology*. 2005;1(1):2005.0028. doi:10.1038/msb4100036.
69. Schneider DS. Tracing Personalized Health Curves during Infections. *PLOS Biology*. 2011;9(9):e1001158. doi:10.1371/journal.pbio.1001158.
70. Kasumovic MM. The multidimensional consequences of the juvenile environment: towards an integrative view of the adult phenotype. *Animal Behaviour*. 2013;85(5):1049–1059. doi:10.1016/j.anbehav.2013.02.009.
71. Dore AA, McDowall L, Rouse J, Bretman A, Gage MJG, Chapman T. The role of complex cues in social and reproductive plasticity. *Behavioral Ecology and Sociobiology*. 2018;72(8):124. doi:10.1007/s00265-018-2539-x.
72. Westneat DF, Potts LJ, Sasser KL, Shaffer JD. Causes and Consequences of Phenotypic Plasticity in Complex Environments. *Trends in Ecology & Evolution*. 2019;34(6):555–568. doi:10.1016/j.tree.2019.02.010.
73. Moller AP, Pomiankowski A. Why have birds got multiple sexual ornaments? *Behavioral Ecology and Sociobiology*. 1993;32(3):167–176. doi:10.1007/BF00173774.


74. Getty T. The maintenance of phenotypic plasticity as a signal detection problem. *The American Naturalist*. 1996;148(2):378–385.
75. DeWitt TJ. Costs and limits of phenotypic plasticity: Tests with predator-induced morphology and life history in a freshwater snail. *Journal of Evolutionary Biology*. 1998;11(4). doi:10.1007/s000360050100.
76. Anderson RM, May RM. Coevolution of hosts and parasites. *Parasitology*. 1982;85(2):411–426. doi:10.1017/S0031182000055360.
77. Alizon S, Hurford A, Mideo N, Van Baalen M. Virulence evolution and the trade-off hypothesis: history, current state of affairs and the future. *Journal of Evolutionary Biology*. 2009;22(2):245–259. doi:10.1111/j.1420-9101.2008.01658.x.
78. Mackinnon MJ, Read AF. Genetic Relationships between Parasite Virulence and Transmission in the Rodent Malaria *Plasmodium chabaudi*. *Evolution*. 1999;53(3):689–703. doi:10.1111/j.1558-5646.1999.tb05364.x.
79. Mackinnon MJ, Gaffney DJ, Read AF. Virulence in rodent malaria: host genotype by parasite genotype interactions. *Infection, Genetics and Evolution*. 2002;1(4):287–296. doi:10.1016/S1567-1348(02)00039-4.
80. Mackinnon MJ, Read AF. The effects of host immunity on virulence–transmissibility relationships in the rodent malaria parasite *Plasmodium chabaudi*. *Parasitology*. 2003;126(2):103–112. doi:10.1017/S003118200200272X.
81. Paul R, Lafond T, Müller-Graf C, Nithiuthai S, Brey P, Koella J. Experimental evaluation of the relationship between lethal or non-lethal virulence and transmission success in malaria parasite infections. *BMC Evolutionary Biology*. 2004;4(1):30. doi:10.1186/1471-2148-4-30.
82. Regoes RR, Nowak MA, Bonhoeffer S. Evolution of Virulence in a Heterogeneous Host Population. *Evolution*. 2000;54(1):64–71. doi:10.1111/j.0014-3820.2000.tb00008.x.
83. Williams PD. New Insights into Virulence Evolution in Multigroup Hosts. *The American Naturalist*. 2012;179(2):228–239. doi:10.1086/663690.
84. White PS, Choi A, Pandey R, Menezes A, Penley M, Gibson AK, et al. Host heterogeneity mitigates virulence evolution. *Biology Letters*. 2020;16(1):20190744. doi:10.1098/rsbl.2019.0744.
85. Hawking F, Wilson ME, Gammage K. Evidence for cyclic development and short-lived maturity in the gametocytes of *Plasmodium falciparum*. *Transactions of The Royal Society of Tropical Medicine and Hygiene*. 1971;65(5):549–559. doi:10.1016/0035-9203(71)90036-8.
86. Padilla DK, Adolph SC. Plastic inducible morphologies are not always adaptive: The importance of time delays in a stochastic environment. *Evolutionary Ecology*. 1996;10(1):105–117. doi:10.1007/BF01239351.
87. Hoi AG, Greischar MA, Mideo N. Limited impact of within-vector ecology on the evolution of malaria parasite transmission investment. *Frontiers in Malaria*. 2024;2. doi:10.3389/fmala.2024.1392060.


88. Mukherjee S, Bassler BL. Bacterial quorum sensing in complex and dynamically changing environments. *Nature reviews Microbiology*. 2019;17(6):371–382. doi:10.1038/s41579-019-0186-5.
89. Greischar MA, Read AF, Bjørnstad ON. Synchrony in malaria infections: How intensifying within-host competition can be adaptive. *American Naturalist*. 2014;183(2). doi:10.1086/674357.
90. Chang KH, Tam M, Stevenson MM. Modulation of the Course and Outcome of Blood-Stage Malaria by Erythropoietin-Induced Reticulocytosis. *The Journal of Infectious Diseases*. 2004;189(4):735–743. doi:10.1086/381458.
91. Mideo N, Barclay VC, Chan BHK, Savill NJ, Read AF, Day T. Understanding and predicting strain-specific patterns of pathogenesis in the rodent malaria *Plasmodium chabaudi*. *American Naturalist*. 2008;172(5):214–238. doi:doi/10.1086/591684.
92. Landau I, Boulard Y. Life Cycles and Morphology. In: Killick-Kendrick R, Peters W, editors. *Rodent Malaria*. Academic Press; 1978. p. 53–84.
93. Gautret P, Miltgen F, Gantier JC, Chabaud AG, Landau I. Enhanced gametocyte formation by *Plasmodium chabaudi* in immature erythrocytes: Pattern of production, sequestration, and infectivity to mosquitoes. *Journal of Parasitology*. 1996;82(6). doi:10.2307/3284196.
94. Mideo N, Savill NJ, Chadwick W, Schneider P, Read AF, Day T, et al. Causes of variation in malaria infection dynamics: Insights from theory and data. *American Naturalist*. 2011;178(6). doi:10.1086/662670.
95. Smith E, Evans G, Foadi J. An effective introduction to structural crystallography using 1D Gaussian atoms. *European Journal of Physics*. 2017;38(6):065501. doi:10.1088/1361-6404/AA8188.
96. Wood SN. mgcv: GAMs and generalized ridge regression for R. *R News*. 2001;1.
97. Brancucci NMB, Gerdt JP, Wang C, Niz MD, Philip N, Adapa SR, et al. Lysophosphatidylcholine Regulates Sexual Stage Differentiation in the Human Malaria Parasite *Plasmodium falciparum*. *Cell*. 2017;171(7):1532–1544.e15. doi:10.1016/J.CELL.2017.10.020.
98. Regev-Rudzki N, Wilson DW, Carvalho TG, Sisqueira X, Coleman BM, Rug M, et al. Cell-Cell Communication between Malaria-Infected Red Blood Cells via Exosome-like Vesicles. *Cell*. 2013;153(5):1120–1133. doi:10.1016/J.CELL.2013.04.029.
99. Boyle MJ, Wilson DW, Richards JS, Riglar DT, Tetteh KKA, Conway DJ, et al. Isolation of viable *Plasmodium falciparum* merozoites to define erythrocyte invasion events and advance vaccine and drug development. *Proceedings of the National Academy of Sciences of the United States of America*. 2010;107(32). doi:10.1073/pnas.1009198107.
100. Stephens R, Culleton RL, Lamb TJ. The contribution of *Plasmodium chabaudi* to our understanding of malaria. *Trends in Parasitology*. 2012;28(2). doi:10.1016/j.pt.2011.10.006.
101. Soetaert K, Petzoldt T, Setzer RW. Solving Differential Equations in R: Package deSolve. *Journal of Statistical Software*. 2020;33(9):1–25. doi:10.18637/jss.v033.i09.


102. Brown PN, Byrne GD, Hindmarsh AC. VODE: A Variable-Coefficient ODE Solver. <http://dxdoiorg/101137/0910062>. 2006;10(5):1038–1051. doi:10.1137/0910062.
103. Gerber F, Furrer R. optimParallel: An R Package Providing a Parallel Version of the L-BFGS-B Optimization Method. *The R Journal*. 2019;11(1):352–358. doi:10.32614/RJ-2019-030.
104. Mullen KM, Ardia D, Gil DL, Windover D, Cline J. DEoptim: An R package for global optimization by differential evolution. *Journal of Statistical Software*. 2011;40(6). doi:10.18637/jss.v040.i06.
105. Lampinen J, Storn R. Differential Evolution. In: Onwubolu GC, Babu BV, editors. *New Optimization Techniques in Engineering*. Studies in Fuzziness and Soft Computing. Berlin, Heidelberg: Springer; 2004. p. 123–166. Available from: https://doi.org/10.1007/978-3-540-39930-8_6.
106. Wickham H. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York; 2016. Available from: <https://ggplot2.tidyverse.org>.
107. R Core Team. *R: A language and environment for statistical computing*; 2021. Available from: <https://www.r-project.org/>.
108. Bivand R, Rowlingson B, Diggle P, Petris G, Eglen S. *splancs: Spatial and Space-Time Point Pattern Analysis*; 2024. Available from: <https://CRAN.R-project.org/package=splancs>.

















[Click here to access/download](#)
Supporting Information
S1_Fig.tif



Click here to access/download
Supporting Information
S2_Fig.tif



[Click here to access/download](#)
Supporting Information
S3_Fig.tif





[Click here to access/download](#)
Supporting Information
S4_Fig.tif



Click here to access/download
Supporting Information
S5_Fig.tif



[Click here to access/download](#)
Supporting Information
S6_Fig.tif



Click here to access/download
Supporting Information
S7_Fig.tif



Click here to access/download

Other

2025_05_07_plos_project_plasmodium_draft8_clean_no
_figures.tex

Optimal cues for transmission investment in malaria parasites

Avril L. Wang^{1□*}, Megan A. Greischar², Nicole Mideo¹

1 Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, Ontario, Canada

2 Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, New York, United States

□Current Address: Department of Biology, Massachusetts Institute of Technology, Cambridge, Massachusetts, United States

* avril137@mit.edu

Abstract

The timing of investment into reproduction is a key determinant of lifetime reproductive success (fitness). Many organisms exhibit plastic, i.e., environmentally-responsive, investment strategies, raising the questions of what environmental cues trigger responses and why organisms have evolved to respond to those particular cues. For malaria parasites (*Plasmodium spp.*), investment into the production of specialized transmission stages (versus stages that replicate asexually within the host) is synonymous with reproductive investment and also plastic, responding to host- and parasite-derived factors. Previous theory has identified optimal plastic transmission investment strategies for the rodent malaria parasite, *Plasmodium chabaudi*, as a function of the time since infection, implicitly assuming that parasites have perfect information about the within-host environment and how it is changing. We extend that theory to ask which cue(s) *should parasites use? Put another way, which cue(s) [snip]* maximize parasite fitness, quantified as host infectiousness during acute infection? Our results show that sensing a parasite-associated cue, e.g., the abundance of infected red blood cells or transmission stages, allows parasites to achieve fitness approaching that of the optimal time-varying strategy, but only when parasites perceive the cue non-linearly, responding more sensitively to changes at low densities. [snip] However, no single cue can recreate the best time-varying strategy or allow parasites to adopt terminal investment as the infection ends, a classic expectation for reproductive investment. Sensing two cues—log-transformed infected and uninfected red blood cell abundance—enables parasites to accurately track the progression of the infection, permits terminal investment, and recovers the fitness of the optimal time-varying investment strategy. Importantly, parasites that detect two cues more efficiently exploit hosts, resulting in higher virulence compared with those sensing only one cue. Finally, our results suggest a potential tradeoff between achieving an optimal transmission investment strategy *in a given host environment [snip]* and robustness in the face of environmental or developmental fluctuations.

Introduction

A key goal of life history theory is to understand how organisms should invest in reproduction over the course of their lives and explaining the substantial variation in patterns observed (e.g., [1]). Reproductive investment entails a resource allocation

tradeoff: resources allocated to growth cannot, generally, also be allocated to reproduction. Classic work predicts that reproductive investment will increase with age [2,3]. Additionally, reproductive investment is expected to depend on the state of an organism, a product of intrinsic, physiological factors and external, environmental ones [4]. Experimental data have shown that investment can be plastic, [snip]with age and state [snip] both shaping[snip] an organism’s reproductive efforts to maximize lifetime fitness (e.g., [5,6]). How organisms estimate the time remaining for reproduction—that is, what cues they utilize from their own physiology and the environment—remains an open question for many systems.

For sexually-reproducing parasites, patterns of reproductive investment can have clinical and epidemiological consequences. Protozoan parasites of *Plasmodium* spp.—the etiological agents of malaria—infect red blood cells of their host, and a given infected cell will either produce more asexual, cell-infecting stages (“growth” of an infection) or produce a single gametocyte, a sexually-differentiated stage that can be transmitted to mosquito vectors (potentially “reproducing” an infection). The proportion of infected cells that produce transmissible gametocytes, known in the malaria literature as the “conversion rate,” is a measure of investment in reproduction/transmission [7]. This trait has a direct link to infectiousness, since more gametocytes mean greater chance of onward transmission to a biting mosquito [8–10] and can influence the virulence or severity of infection, since all else equal, lower transmission investment means more host-damaging asexual parasites are produced [7].

Importantly, transmission investment in *Plasmodium* spp. is plastic, changing in response to conspecific density [11,12], replication rate [13], parasite-conditioned media [14,15] (which could contain infected red blood cell (iRBC)-derived vesicles [16]), RBC age structure/availability [13,17–19], and antimalarial exposure (e.g., [13,20]). Understanding why transmission investment is plastic and what the optimal plastic strategy is has been the subject of several prior studies [7,13,21–24]. In one example, using the experimentally tractable rodent malaria system, *Plasmodium chabaudi*, and different doses of antimalarial drugs, Schneider *et al.* [13] showed that intermediate reductions in parasite density led to reduced investment in transmission while strong reductions had the opposite effect. These results are in line with standard predictions from life history theory for malaria parasites, where mild stressors should promote reproductive restraint and strong stressors that signal imminent clearance of an infection should lead to complete investment in transmission [7], i.e., terminal investment [25]. But “stress” will vary over the course of infection—even in the absence of drug treatment—as parasites deplete host resources and hosts respond via immunity. These dynamics are thought to [snip] explain why transmission investment in malaria parasites is variable over the lifespan of infections (i.e., plastic) [11–13,22,26,27].

Mathematical modelling provides another route to understanding and predicting adaptive plasticity in parasite traits. Models assessing the fitness consequences of different patterns of transmission investment demonstrate that the best strategy involves initial reproductive restraint that facilitates within-host parasite population growth (rather than early investment in gametocytes that would have very low likelihood of onward transmission due to low densities), followed by an increase in investment once parasite densities have risen, culminating in terminal investment as the infection is ending [21,23]. This work derived the optimal transmission investment strategy by assuming that investment is a function of infection age, about which parasites have perfect knowledge. In addition to assuming that parasites can, in effect, “tell time,” this work also assumes [snip] all infections proceed identically, a limitation[snip] common to

models of age-specific reproductive investment [28]. In reality, time would be a poor proxy for environmental conditions since the pace of infection changes with parasite dose [29,30], host immunity [30,31], and myriad other factors. Therefore, understanding the links between transmission investment and within-host environmental cues will result in more biologically relevant models that can inform empirical observation of *Plasmodium* spp. transmission investment. Currently, it is unclear whether observed deviations from the theoretical optimal strategy are due to faulty theories or limitations in the cues that parasites perceive. Understanding the fitness consequences of sensing different cues can delineate those possibilities. Additionally, understanding what cues parasites should sense will enable better predictions on [snip] the influence of novel perturbations, like vaccination, on [snip] infection dynamics and outcomes.

In this study, we extend a previously-published model of within-host rodent malaria infection dynamics [23] to include transmission investment as a function of environmental cues, i.e., components of state. [snip]. In our model, different cues may give rise to different early patterns of transmission investment, which will alter the trajectory of infections and feedback to influence the subsequent expression of the trait. We use numerical optimization to investigate which within-host environmental cue or cues malaria parasites “should” sense for determining transmission investment (i.e., which cues give rise to the highest parasite fitness) and the optimal shape of the relationship between each cue and transmission investment (i.e., the optimal reaction norms). We assess the consequences of perceiving different cues on infection dynamics and disease severity and quantify the robustness of different transmission investment strategies in the face of environmental and developmental stochasticity. [snip] Collectively, our study further characterizes plasticity in *P. chabaudi* life history, identifies the cues we expect parasites to be using, and introduces [snip] a mathematical framework that explicitly incorporates cue perception that could be used to generate testable hypotheses and broadly applied to other organisms.

Results

[snip] Our mathematical model tracks within-host dynamics of infection with the rodent malaria parasite *P. chabaudi* (Fig 1A). Asexual parasites, called merozoites (M), invade uninfected red blood cells (RBCs; R). The resulting infected RBC (iRBC; I) can go on to produce more merozoites that are either asexually-committed (M ; producing more merozoites if they infect an RBC) or sexually-committed (M_g ; producing a single gametocyte, G , if they infect an RBC). RBCs infected with these different types of merozoites are denoted as I and I_g , respectively. Importantly, the proportion of iRBCs that follow these different developmental routes (i.e., producing [snip] M_g) defines transmission investment. While previous approaches have treated transmission investment [snip] as either fixed or time-dependent (e.g., [21,23,32–36]), here we explicitly model it as a function of within-host environmental cues (see Materials and Methods). We select as plausible cues various measures of the density of parasites and host resources for which there is some experimental evidence to impact transmission investment in *Plasmodium* spp. (Table 1). We quantify the fitness of parasites sensing various cues using the cumulative transmission potential, which measures the summed probability of infecting mosquito vectors across a 20-day infection period, assuming a constant biting rate and mosquito abundance [23].

In our model, cue perception occurs at the beginning of asexual iRBC development (Fig 1B), determining the proportion of asexual iRBCs that produces sexually-committed merozoites. Note that this modeling choice introduces a time lag

Fig 1. Schematic of *P. chabaudi* within-host infection dynamics and fitness optimization. (A) Flow diagram of model tracking the dynamics of red blood cells (R ; RBCs), RBCs infected with asexual parasites (I), RBCs infected with sexual—but not yet mature—parasites (I_g), asexually-committed merozoites (M), sexually-committed merozoites (M_g), and gametocytes (G). (B) Temporal dynamics of cue perception. Cue perception occurs one day before the production of asexual/sexual merozoites, thereby introducing a delay between cue perception and the consequences of cue perception (i.e., production of more asexual infected RBCs or gametocytes). A further delay occurs in the sexual cycle since gametocytes of *P. chabaudi* take two days to mature [37]. Dashed circles indicate the bursting/maturation of a particular cohort of iRBCs. (C) Modeling reaction norm evolution. We model the relationship between the perceived cue and transmission investment (i.e., the reaction norm) as a flexible spline (black lines). Specifically, a cubic spline can specify any third order polynomial at a given x value as a linear combination of the columns in its basis matrix [38] (exemplified by the red, green, and blue dashed lines; combined they produce the solid, black line). For local optimization, an arbitrary starting spline is picked (left panel) and an optimization algorithm is used to adjust the relative weights of the basis functions until a fitness maximum is achieved (going from left to right). The small coloured arrows depict how the weight of a given basis function has changed from the previous panel.

between sexual commitment (as a result of cue perception) and gametocyte production, which is a departure from previous modeling work [23], but reflects empirical data on the canonical timeline of sexual commitment from the best studied system (*Plasmodium falciparum* [39]). For each putative cue, we define its associated reaction norm using flexible splines (Fig 1C). Specifically, the shape of the reaction norm is determined by a linear combination of three basis functions comprising a cubic spline. Adjusting the spline coefficients (i.e., the weights associated with each column of the spline basis matrix) changes the shape of the reaction norm. We used a combination of local and global numerical optimization algorithms (see Materials and Methods) to adjust the spline coefficients and find the reaction norm that maximizes parasite fitness over a 20-day infection, as in previous work [23, 40] (Fig 1C). We test both raw densities and logged densities as cues to expand the range of reaction norm shapes explored and because changes in infection dynamics are observed in this system when doses change by orders of magnitude [29]. For parasites sensing time, we used the same optimization process, except setting the cue range to time since infection (day 0 to 20). Finally, following [41], our within-host model includes two forms of immunity, one targeting specifically infected cells (I and I_g) and one targeting all red blood cells (I , I_g and R). Thus, while resource limitation plays some role in driving infection dynamics in this model, immunity allows for the clearance of the infection.

Cue perception mediates parasite life history and fitness

Based on previous work modeling *P. chabaudi* transmission investment as a function of time [23], we expect the following as the best strategy over a 20-day infection: parasites should delay investment early on, display a rapid increase in investment when sufficient gametocytes can be produced to ensure transmission, subtly reduce investment in the face of declining resources, and ultimately rise to the highest levels of investment shortly before the end of infection (roughly, “terminal investment” [7, 23]). Although the model we employ here has a few key differences from that work (see Materials and Methods), we find a similar optimal transmission investment pattern as in [23] when we assume that investment is a function of time (Fig 2A). When parasites instead respond to their environment and are constrained to perceive a single within-host cue, these same

Table 1. Putative within-host environmental cues

Cue	Abbreviation	Experimental Evidence
Infected RBC (iRBC) density		[12, 13, 16, 24]
Asexually-committed	I	
Asexually-committed, $\log_{10}$	I log	
Sexually-committed	Ig	
Sexually-committed, $\log_{10}$	Ig log	
Total (I+Ig)	Total I	
Total (I+Ig) , $\log_{10}$	Total I log	
Gametocyte density		[12, 16, 24]
Gametocytes	G	
Gametocytes, $\log_{10}$	G log	
Resource density		[12, 13, 24]
Uninfected RBC	R	
Uninfected RBC, $\log_{10}$	R log	

patterns of transmission investment cannot be fully recapitulated. Consequently, parasite fitness is lower in every case than when the cue is time (Fig 2A, left panel).

The environmental cues that result in the highest parasite fitness include logged parasite-centric cues (e.g., $\log_{10}$ densities of sexually-committed iRBCs). Perceiving these cues is sufficient to produce the initial delay in transmission investment, but the delay is longer than parasites **sensing time as the cue**, and there is no increase in investment near the end of infection (Fig 2A, right panel). Given that parasite densities rise non-linearly at the start of infections (both simulated and real [11, 42–45]), sensing logged rather than raw densities results in reaction norms that are more sensitive to those early changes (Fig 2B), enabling parasites to increase transmission investment after the densities have reached a rough threshold. In contrast, sensing raw parasite densities results in reaction norms that essentially ignore changes at the low parasite densities characteristic of early infections (Fig 2B). The density of host resources (RBCs; raw or logged) is also a poor cue for initial stages of infection, since they change very little early on and, therefore, do not allow parasites to engage in delayed transmission investment (Fig 2A). Overall, our results broadly suggest that early transmission delay necessitates **sensitivity [snip]** to changes at low parasite density.

Fig 2. Cue perception mediates parasite fitness and dynamics of transmission investment. (A) On the left, fitness values achieved by parasites adopting the optimal transmission investment strategy for each cue, normalized with respect to the time-varying strategy, are displayed in descending order. On the right, the dynamics of transmission investment, given the optimal reaction norms, are plotted. Investment varies from dark blue (low values) to yellow (high values). **The first four cues after time, which include measures of [snip]** parasite density on a $\log_{10}$ scale, enable parasites to obtain the highest fitness and delay in transmission investment. (B) Optimal reaction norms for different cues. Each facet displays the reaction norms for raw (black, dotted lines) or log-transformed (red, solid lines) cues. The ranges of cue values “experienced” by a parasite adopting the optimal reaction norm are displayed above (raw) or below (log-transformed) the reaction norms as rug plots (horizontal bars). Based on the shape of the reaction norms, we conclude that sensing log-transformed parasite-centric cues enables more sensitive detection of (and responses to) **the initial rise [snip]** in parasite abundance.

Fig 3. Dual cue perception increases parasite fitness and enables terminal investment. (A) RBC density uniquely marks the onset of terminal investment. In the left panel, we plot transmission investment of parasites sensing time (i.e., the ideal dynamics) and mark its key features: transmission delay (purple) and terminal investment (red). We then plot the dynamics of single cues on the right and mark the range of cue values reached during terminal investment with two red, dashed lines. The overlap of the red dashed lines with earlier cue values indicates that parasites may not be able to uniquely recognize the end of the infection. (B) The fitness of parasites (normalized with respect to the time-varying strategy) adopting the optimal strategy for each dual cue combination is indicated with a blue triangle, while the fitness obtained when perceiving the best performing single cue (of the combination) is represented with an orange circle. To obtain the fitness of a fair comparator time-varying strategy, we optimized transmission investment as a function of time using a spline with nine degrees of freedom (the same degrees of freedom used in the dual cue model), rather than the three degrees of freedom used above and previously [23]. (Note that the marginal fitness gains of using more flexible splines are small [23], but tracking two environmental cues necessitates more degrees of freedom.) (C) The pattern of transmission investment over time for parasites sensing the best dual cue combination (R log & I log, blue square) along with individual components of the cue (I log, orange circle; R log, yellow triangle; each optimized with nine degrees of freedom) and the ideal dynamic (Time, black cross) are plotted together. Parasites sensing the combination of logged asexual iRBC and logged RBC best approximate the time-varying strategy. (D) Transmission investment (colour of heatmap) is plotted as a two-dimensional reaction norm, with logged RBC and logged asexual iRBC as cues. The white line indicates the trajectory of infection, starting at the top left, for parasites adopting this optimal reaction norm, with each point representing a single day. The arrow on the white line indicates the passage of time. For (C-D), the four stages outlined in the results section are annotated with I, II, III, and IV.

Dual cue perception is necessary for terminal investment

We found that parasites sensing a single cue cannot reproduce the expected optimal transmission investment strategy, particularly the rapid increase in transmission investment during late infection (i.e., terminal investment). Achieving terminal investment presumably requires sensing a cue that provides a unique signal at the later stages of infection. Inspecting infection dynamics produced by parasites that sense time revealed that only RBC density provides such a distinctive late-stage signal (Fig 3A). Given the poor performance of this cue in early stages of infection, these results suggest that simultaneous perception of RBC and a parasite-centric cue could provide infection stage-specific information that enables both early reproductive restraint and terminal investment. Consistent with this expectation, in a rodent malaria experiment, including variables that reflect parasite density and host resources (specifically, replication rate and changes in RBC density) in statistical models explained substantially more variation in transmission investment than the parasite-centric variable alone [13].

To investigate the fitness consequences of sensing two cues, we optimized the transmission investment strategy of parasites sensing all pairwise combinations of log-transformed cues, using two-dimensional spline surfaces and focusing on combinations of log-transformed cues given their higher performance in the single cue optimizations. Our results suggest that sensing [snip] dual cue combinations improves parasite fitness relative to sensing a single cue (Fig 3B), with parasites sensing logged RBC and logged iRBC (asexual or total) exhibiting notably higher fitness (Fig 3B).

This higher fitness is conferred by transmission investment patterns that are close to the time-varying ideal, with an initial delay in transmission investment and a late infection (nearly) terminal investment before returning to a low transmission investment (Fig 3C, S2 Fig). Increasing the spline flexibility of the single cue models (to match the degrees of freedom in dual cue models) could not reproduce this same pattern (Fig 3C), confirming that dual cue perception specifically enables terminal investment and its associated fitness benefits (S3 Fig).

To better understand the optimal patterns of transmission investment, we next analyzed the reaction norm of the dual cue combination that conferred the highest fitness: logged asexual iRBC and logged RBC. The pattern of transmission investment is defined by a looping journey through the reaction norm surface (Fig 3D), confirming that the cue combination traces a unique path over the course of an infection. We categorized the reaction norm into four distinct sections/stages (Fig 3D). Stage I corresponds to the initial phase of infection, characterized by low transmission investment, low parasite density, and high RBC density (Fig 3C-D). As the infection progresses, the reaction norm enters stage II, characterized by asexual iRBC production and RBC decline. During this stage, the optimal transmission investment increases considerably, followed by subtle declines as the iRBC population continues to increase (Fig 3C-D). The transition from a low to high transmission investment observed as stage I transitions into stage II constitutes delayed transmission investment. Following peak parasite density, the reaction norm enters stage III, which is marked by decreasing iRBC population and gradual replenishment of RBCs, signaling imminent infection clearance. At this stage, transmission investment increases to peak levels (i.e., terminal investment). Finally, as the infection nears clearance and the reaction norm enters stage IV, transmission investment sharply drops as the host heads towards recovery (Fig 3C-D). We note that this drop in predicted transmission investment is likely a consequence of the constraints of the spline surface, and different strategies during this stage would have negligible impact on fitness [23]. Overall, our results suggest that each cue in this combination plays a crucial role in defining progression through infection stages I to III, with asexual iRBC density exhibiting the largest variation in early and late infection (stages I and III), and logged RBC density exhibiting the largest variation during mid-infection (Stage II; Fig 3D). This temporal staggering of iRBC and RBC dynamics (i.e., their hysteretic relationship) ensures that these infection stages can be tracked. As a result, dual cue perception provides parasites with the ability to “state differentiate,” to accurately monitor the progress of the infection and adjust transmission investment accordingly.

Intriguingly, the dual cue combinations that are most informative for parasites making life history decisions (top two in Fig 3B) are also the most informative for predicting patient outcomes in clinical malaria infections [46]. In both cases, two slightly out-of-phase within-host measures can uniquely define a patient’s (or simulated host’s) position on the path from health through sickness to recovery. Plotting this trajectory through disease maps (i.e., the two measures plotted against each other, as in Fig 3D) produces a looping, or hysteretic, pattern, with each infection stage occupying a unique coordinate in this two-dimensional space [46]. To further validate that forming the characteristic looping pattern is a unique feature of our identified best cue combinations, we constructed disease maps using experimental data from laboratory mice infected by a single strain of *P. chabaudi* [11, 42–45, 47] and simulated data from our models where parasite senses time. Both experimental and simulated data indicate that raw or logged RBC with logged iRBC (asexual or total in the model; only total is available in the experimental data) are the only cue combinations that generate fully

separated loops (e.g., Fig 4A), while other cue combinations do not (e.g., Fig 4B and C; full data shown in S4 Fig and S5 Fig). When early and late infection stages are “condensed” in space, simulated parasites fail to state differentiate and do not delay transmission investment or terminally invest (S2 Fig).[snip] In summary, our results demonstrate that optimal transmission investment requires parasites to unambiguously state differentiate, track infection stages, and dynamically adjust investment in response to changing conditions. Our data suggest that perceiving hysteretic cues — of which a combination of RBC and logged iRBC provides a clean delineation of different infection stages — is a viable method of attaining state differentiation.

Fig 4. Trajectories of log-transformed RBC and iRBC densities form a looping pattern *in silico* and *in vivo*. Simulated (left) and experimental (right) trajectories of cues that (A) enable transmission delay and terminal investment or (B-C) fail to achieve these patterns [snip]. For each panel, the left graph displays the results from our simulations where parasite senses time with nine degrees of freedom, with the distance between each circle representing a single day and the colour of the line representing the observed transmission investment. The right graph displays the infection trajectory of laboratory *P. chabaudi* infections in mice, with the colour of the lines indicating the day post-infection. Note that asexually- and sexually-committed iRBCs were not distinguished in the experimental data. For the simulated trajectories, the first 3 days correspond to the period of delayed transmission investment and the last 5 days correspond to when terminal investment is expected. In (B), early changes in asexual iRBC density cannot be clearly separated (see the red, dotted box), therefore preventing a delayed transmission investment strategy. Similarly, in (C), late state infection stages cannot be separated from earlier time points, preventing parasites from terminally investing.

Dual cue perception exacerbates the fitness loss incurred by environmental and developmental stochasticity

The benefits of plasticity depend on the quality of information organisms obtain from environmental cues. Adaptive phenotypic plasticity is contingent on proper phenotype–environment matching, where the perceived cue is a reliable predictor of the optimal phenotype [48–52]. Theory suggests that a weak correlation between cue(s) and the optimal phenotype (i.e., low cue reliability) could disfavour plasticity relative to other strategies (i.e., local adaptation [49,50,53], canalization [52], bet-hedging [53]). For malaria parasites, there may be unpredictable variation in the environment (e.g., RBC availability [41,54,55], immunity [41,56–58]) or in traits governing parasite development within the host [41]. Such environmental and developmental stochasticity could preclude evolution towards the strategies we predict above. Although our model allows parasites to sense components of their state, and their resulting level of transmission investment will feedback to influence that state, the results above assume that all else is equal between hosts. Specifically, [snip] the optimal reaction norms identified above are associated with a particular [snip] set of model parameters (the median estimates from posterior distributions in [41]; we refer to this as the deterministic model). This implicitly assumes that hosts are all the same or that parasites are moving between hosts sufficiently quickly that they adapt to an “average” environment.

[snip]

To investigate how biologically relevant variation affects cue performance, we built a semi-stochastic model based on the work of Kamiya *et al.* [41], which employed a hierarchical Bayesian method to quantify inter-host variation in parasite burst size,

erythropoiesis (RBC replenishment), and immune strengths. [snip] For all log-transformed single cues and [snip] dual cues, we performed 1000 20-day simulations, holding the reaction norm at the optimum from the relevant deterministic simulation, and simultaneously varying the RBC replenishment rate (ρ), parasite burst size (β), immune activation strengths (ψ_n and ψ_w) and immunity half-lives (ϕ_n and ϕ_w), sampling parameter values from the reported posterior distributions [41][snip](S6 Fig). Each choice of parameter set will influence the within-host infection dynamics, potentially altering the trajectory of the perceived cue(s), resulting in different, expressed patterns of transmission investment and, thus, fitness. Here we are asking, if parasites are “optimized” to the average host, does environmental and developmental variation change inferences about which cues are best? We quantified the effect of variation on cue performance using the geometric mean fitness, commonly used to assess fitness in fluctuating environments [59,60].

We find that the geometric mean fitness of parasites responding to a given cue, in the face of this variation, is largely uncoupled from fitness in the deterministic model (i.e., the environment in which the strategy was optimized; Fig 5A), emphasizing that the “best” reaction norm for a certain environment does not always confer the best phenotype in another. Interestingly, all dual cues have a lower geometric mean fitness than the best-performing single cue (I log, Fig 5A). Furthermore, this result emphasizes the challenge of using time as an environmental proxy, since its geometric mean fitness in the semi-stochastic model is among the bottom half of cues. Allowing only one parameter to vary similarly tends to reduce the fitness of the best dual cue strategies disproportionately (S7 Fig). These results indicate a potential tradeoff between fine-tuning fitness in a given environment and susceptibility towards environmental and developmental stochasticity.

While sensing two cues provides more information about “where” an infection is, it could also increase the amount of perceived noise and exacerbate the negative effects of environmental/developmental stochasticity. To test whether increased noise perception reduces the fitness of dual cue-sensing parasites, we compared the trajectories of the most “robust” single cue (I log; Fig 5B left panel) and the dual cue combination with the highest deterministic model fitness (R log & I log; Fig 5B right panel). As expected, the variation in asexual iRBC density across our simulations is much lower than that of both asexual iRBC and RBC (Fig 5B top panels). While parasites sensing both types of cues delay transmission, those sensing logged RBC and iRBC display highly variable transmission investment dynamics (Fig 5B bottom right). In contrast, the transmission investment pattern of parasites sensing only logged asexual iRBC resembles that of the deterministic model (Fig 5B bottom left). These results suggest that sensing certain dual cue combinations can reduce parasite fitness by increasing noise perception that induces suboptimal transmission investment.

[snip]

Dual cue perception leads to higher virulence

Given that cue perception mediates infection dynamics, it could also influence the severity of infections, or virulence. We estimated virulence using the methods described in Torres *et al.* [46], where infection dynamics are mapped in disease space, with the y-axis quantifying host health (RBC density) and the x-axis quantifying parasite load (total iRBC density). A more virulent infection is expected to produce a higher parasite load while consuming more host resources, tracing a larger loop through disease space. To evaluate the effects of cue perception on virulence, we generated 30-day disease maps of infections, allowing sufficient time for the infection to fully resolve in our model. We

Fig 5. Optimal transmission investment strategies do not always translate effectively across different environmental and developmental conditions. (A) Fitness of parasite strategies in the deterministic model is largely uncoupled from the geometric fitness of those same strategies in the semi-stochastic model. Plotted on the y axis is the geometric mean fitness of 1000 simulations of parasites adopting the optimal transmission investment strategy (sensing time from the deterministic model) using a semi-stochastic model where six parameters are randomly drawn from posterior distributions derived by Kamiya *et al.* [41]. The fitness from the deterministic model is plotted on the x-axis and all values are normalized with respect to the time-varying strategy with nine degrees of freedom (i.e., the dual cue comparator shown in Fig 3C). Note we also plot the fitness of the time-varying strategy with three degrees of freedom. Overall, dual cue-sensing parasites tend to perform worse than single cue-sensing parasites in the face of variation. The cue with the highest geometric fitness (I log) is indicated by a dotted orange box, while the cue combination with the highest deterministic model fitness (R log & I log) is bound by a blue dotted box. [snip] (B) Sensing dual cues can increase noise perception and lead to highly variable transmission investment dynamics. The top two graphs display the cue trajectories for parasites sensing the best performing cue in the semi-stochastic simulations (logged asexual iRBC density, plotted against time; left) or dual cues (logged asexual iRBC and logged RBC density, plotted against each other; right). The red lines represent individual semi-stochastic simulation results, while the thicker white line represents the dynamics from the deterministic model. The background colour indicates the optimal reaction norm: on the right, this is the same as what is plotted in Fig 3D. The bottom two graphs display the transmission investment dynamics of the same parasites. The thicker orange/blue line represents the transmission investment trajectory from the deterministic model, while the thinner grey lines represent individual semi-stochastic simulation results. [snip]

use the area enclosed by each disease map as a proxy for virulence, with larger areas indicating higher virulence. We note that we are using “virulence” as synonymous with infection severity and not host mortality, thus we assume no link between virulence and the duration of infection. Recent modeling suggests that resource limitation imposes a more important constraint on the evolution of transmission investment in *P. chabaudi* than does host mortality due to infection [61].

Parasites perceiving only a single cue trace smaller disease maps than parasites that adopt the ideal pattern of transmission investment (by sensing time; Fig 6A). Additionally, the area of the disease map is negatively correlated with parasite fitness across all single cues (Fig 6B), suggesting that if parasites are constrained to perceive a single within-host cue, strategies that result in lower virulence would be favoured. Parasites sensing the top two dual cue combinations produce disease maps that are similar to the map traced by parasites adopting the time-varying strategy (Fig 6C) and result in higher virulence relative to parasites sensing a single cue (Fig 6D; compare x-axis values of points in B and D). Overall, these findings suggest that if parasites have limited information or have evolved to respond to only one cue (e.g., because of physiological constraints or high environmental/developmental variation), virulence will be lower, while if parasites are favoured to more finely track infection stages through dual cue perception, they will exploit host resources to a greater extent and infections will be more virulent.

Fig 6. Cue perception mediates infection severity (virulence). (A, C) display the 30-day disease maps of infections where parasites adopt the top two [snip] optimal transmission investment strategies when sensing (A) a single cue or (C) dual cues. We use the area of the map traced as a proxy for virulence [46]. The trajectory of parasites that adopt the ideal transmission investment strategy (sensing time) is shown in black. (B, D) Relationship between disease map area and fitness (normalized with respect to the time-varying strategy) for parasites that sense (B) only one cue or (D) dual cues. The top two [snip] cues for single and dual cue perceiving parasites are coloured, while the grey points represent the other (B) single or (D) dual cues. [snip] Overall, these results suggest that perceiving only a single cue generates selection for strategies that lead to lower virulence while perceiving two cues generates selection for strategies that lead to higher [snip] virulence [snip].

Discussion

Just as with any sexually-reproducing species, investing in reproduction by producing gametocytes is critical for the fitness of *Plasmodium* spp., as this is the only parasite stage capable of being transmitted to mosquito vectors. As for other species, this investment comes with tradeoffs for malaria parasites: a given infected host cell can either produce multiple asexual progeny that can infect new host cells and have the potential to grow or maintain a given infection, or a single, sexually-differentiated gametocyte with the potential to “reproduce” an infection [62]. Thus, maximizing parasite fitness requires balancing these competing demands, the relative importance of which is likely to shift over the course of infection, as the within-host environment changes in response to parasites exploiting the host. Indeed, data show that transmission investment varies throughout infections [11, 12, 26] and appears to change in response to drug treatment in ways predicted to be beneficial for parasites (e.g., [13, 40]). While adaptive plasticity is a well-known solution to the challenge of living in a changing environment (e.g., [63]), there has been skepticism about the fitness benefits of observed parasite plasticity [64], in part because knowledge of parasite environmental sensing mechanisms remains limited. Previous mathematical modeling has treated parasites as either insensitive to environmental change (i.e., constant transmission investment [21, 32–36, 65, 66]) or as perfectly informed (time-varying investment [21, 23, 24]). We instead asked what environmental cues malaria parasites *should* use to set transmission investment, demonstrating that no single cue allows parasites to achieve the best time-varying strategy [23]. Parasites must sense two cues to recapitulate the optimal strategy but that strategy enhances the risk of parasites being “tricked” by environmental and developmental stochasticity into expressing suboptimal phenotypes. We further show that if parasites have the capacity to sense more aspects of their within-host environment, then this has negative consequences for host health.

Previous theoretical work identifies two key features of optimal transmission investment strategies for *P. chabaudi* [23]. First, there should be no investment at the start of infections, when densities are too low to permit successful infection of vectors (i.e., transmission investment delay). Second, parasites should invest all resources into transmission just before an infection ends, a “terminal investment” strategy from life history theory [7]. Our results delineate the environmental cues that can recapitulate these features. Transmission delay requires a sharp, threshold-dependent transition from no investment to an intermediate investment level as the parasite population grows. Making clear-cut decisions in response to continuous cues is common in biological systems (e.g., cell cycle commitment [67] and developmental patterning [68]) and often relies on ultrasensitivity, characterized by a steep response curve. Our results

demonstrate that an ultrasensitive reaction norm where transmission investment rises sharply after a certain population size threshold— a pattern enabled by sensing parasite-centric cues on a logarithmic scale—is sufficient for transmission delay. Prior empirical studies have found that *Plasmodium* transmission investment varies in response to conspecific density [11, 12], parasite-conditioned media [14, 15], and infected red blood cell (iRBC)-derived vesicles [16], suggesting that parasites possess the ability to sense iRBC density. Our results support these responses being part of an adaptive strategy.

In addition to parasite density, *P. chabaudi* has been shown to modulate its transmission investment based on RBC availability and age structure [13, 17–19, 24]. Our results suggest that this is also part of an adaptive strategy, with the combination of iRBC and RBC cues allowing parasites to distinguish between early and late infection. Since most within-host variables go up and down, or vice versa, over time (e.g., iRBC densities rise and then fall due to resource limitation or immunity), one cue cannot provide enough detail to unambiguously establish “where” a parasite is in the course of infection. Combining cues solves this problem. In line with this logic, researchers seeking to define a host’s trajectory over time from health through sickness and on to recovery [69] found a combination of two within-host measures (e.g., parasite density and resource abundance) that are slightly out of phase with each other can uniquely determine a patient’s position on their path for malaria infection [46]. Remarkably, though perhaps predictably, we found that the information desired by medical professionals seeking to predict patient outcomes is also the most informative for parasite life history decisions.

There is a growing appreciation that, across diverse organisms, plasticity is often expressed in response to multiple cues that allow for more finely-tuned strategies [70–72]. Our work adds support to theories suggesting that the adaptive benefits of phenotypic plasticity are enhanced when cues provide non-redundant information [73] and enable organisms to sense distinct states that correspond to specific optimal phenotypes [74]. Yet it is also widely appreciated that integrating multiple cues presents challenges, including the additional metabolic burden of sensing another cue [75] and the multiplicative effect of noise when cues are non-additive [72]. Our complex (decidedly non-additive) dual cue reaction norms present a trade-off: they allow parasites to more accurately recreate optimal strategies in a given environment while rendering them more susceptible to stochastic environmental and developmental change, i.e., host-to-host variability. Such tradeoffs may represent important constraints on the evolution of plastic reproductive investment strategies, perhaps explaining why *P. chabaudi* rarely exhibits *true* terminal investment *in vivo* (displaying only modest increases in transmission investment near infection resolution; [13, 20]), and on the evolution of cue perception more broadly.

[snip]

The tradeoff hypothesis for virulence evolution stipulates that pathogen transmission inevitably harms the host [76]. While the generality of this framework has been contested (see [77] and references within), empirical data from *Plasmodium* spp. have generally supported a positive relationship between virulence and transmissibility [78–81]. Here, we demonstrate that if parasites are limited to sensing a single cue for setting transmission investment, then the strategies that result in the most onward transmission actually [snip] result in less host exploitation. Allowing parasites to sense two cues reverses this relationship and produces the expected pattern of the most onward transmission being achieved by the strategies that also produce the most virulent infections. Importantly, previous work has shown that host heterogeneity

can constrain parasite virulence evolution [82–84] and our results suggest another mechanism by which this may be true: [snip] responding to two cues—while allowing for state differentiation in one host environment—can dramatically reduce fitness [snip] in the face of heterogeneity, potentially favouring single cue (hence, lower virulence) strategies[snip].

Mathematical modeling has enabled us to explore the fitness consequences of cue perception in *P. chabaudi*, and our results suggest where greater model complexity would be most or least informative. For example, incorporating variation in gametocyte circulation time [26] would undercut the reliability of gametocytes as a cue, further supporting our results that it is unlikely to represent the optimal cue. The human malaria parasite *P. falciparum* has a longer sexual development period of 9–12 days [85] and this longer lag between cue perception and phenotype expression will reduce cue reliability [86]. Thus, we anticipate the value of sensing gametocytes to decrease even further in this species. While we have ignored parasite developmental processes that occur in the vector, recent theory suggests they have little qualitative impact on optimal transmission investment strategies [87]. In contrast, the impact of previous exposure on host immunity represents one area where more complexity could qualitatively alter predictions. Our models depict immunologically naive hosts, but where malaria is endemic, parasites will often find themselves in a host with a prior history of infection. Understanding how immune memory influences the quality and reliability of potential within-host cues represents an important area for future research. Our selection of cues assumes that parasites sense only the magnitude of a cue, without detecting temporal changes in that magnitude. Schneider *et al.* [13], in their empirical demonstration of *P. chabaudi* adaptively altering transmission investment, found that replication rate, i.e., the proportional change in parasite densities, had better explanatory power than absolute densities. Broadly speaking, they found transmission investment increased with replication rate, a pattern that is not found with our model since replication rate is highest at the start of infections, before resource or immune limitation kicks in, and not investing in transmission at this point is best. We note that a direct comparison of results is tricky since estimates of transmission investment underlying their observed pattern come from day nine of infections or later. Nonetheless, we identified sensitivity to changes at low population density as a key feature of a “useful” cue and the proportional changes captured by replication rate will similarly share this feature. Another important model assumption is that all parasites perceive the same cue at the same strength, but cues may not be spatially uniform. Variation in signal strength and imperfect cue sensing could lead to heterogeneity in transmission investment in a genotypically uniform parasite population, similar to the dynamics observed in bacterial biofilms [88]. Extrapolating from our results, we hypothesize that if the variation in signal strength is large enough, selection may favour single cue sensing due to the relative robustness to stochastic variation. We note that the variation we incorporated in our semi-stochastic simulations likely underestimates that which would be observed in natural infections, since the dataset fitted by Kamiya *et al.* [41] includes only a single mouse host genetic background studied in a highly controlled experimental system [29]. Thus, environment-phenotype mismatching could be an even larger problem in nature than we have estimated. Additionally, given the variation in burst size and developmental timing across *Plasmodium spp.*, there is a great need to quantify the extent of variation across infection parameters to better understand the role of environmental/developmental heterogeneity on parasite evolution.

Beyond malaria, our work presents a generalizable computational approach to study phenotypic plasticity. By defining plastic traits as an optimizable reaction norm, it is

possible to explore the fitness consequences of sensing different cues and generate hypotheses regarding why organisms respond to certain cues and not others. Mechanistic models of plasticity that capture feedbacks between traits and the environments that shape them will allow for predicting the consequences of perturbations, which is relevant for understanding not only disease dynamics in the face of control measures, but also the fate of organisms whose environment is rapidly altered by climate change.

[snip]
[snip]

Materials and methods

Within-host model of *P. chabaudi* infection dynamics

We modified and extended a previously published mathematical model describing the within-host infection dynamics of the rodent malaria parasite, *P. chabaudi* [23] (Fig 1). The original model consists of ordinary and delay differential equations representing the interactions between the parasite, host red blood cells (RBCs), and host immunity over a 20-day infection. There are three key differences in our model from that previous work. First, we include the activity of two forms of immunity to better capture host defense mechanisms: a targeted immune response that only kills infected RBCs (iRBCs) and an indiscriminate response that kills any RBC, as in [41]. Second, we revised the model to account for the observation that sexually-committed parasites do not engage in gametocytogenesis until the next infection cycle (Next Cycle Conversion; [39]). This necessitated separately tracking the production of sexual merozoites (M_g) and asexual merozoites (M) from infected cells and consequently introduced a delay between sexual commitment and the outcome of sexual commitment (i.e., production of an iRBC that will produce a gametocyte, I_g). Third, we modeled transmission investment as a function of one or two within-host cues, instead of time as in [23], to better capture the mechanism of phenotypic plasticity, since there is no evidence malaria parasites track time across multiple generations. We further describe the implications of these changes on the model below.

[snip]

We depict an immunologically naive host given the availability of experimentally derived parameters. Throughout an infection, we assume indiscriminate immunity eliminates all RBCs (R , I , I_g) whereas targeted immunity eliminates only iRBC (I , I_g). As in [41], we define N and W as the proportion of cells cleared per day by indiscriminate and targeted responses, respectively. The magnitude of this activity is dependent on the total abundance of iRBC ($I + I_g$):

$$\frac{dN}{dt} = \underbrace{\psi_n \frac{I(t) + I_g(t)}{\iota} (1 - N(t))}_{\text{Immune activation}} - \underbrace{\frac{N(t)}{\phi_n}}_{\text{Immune decay}} \quad (1)$$

$$\frac{dW}{dt} = \underbrace{\psi_w \frac{I(t) + I_g(t)}{\iota} (1 - W(t))}_{\text{Immune activation}} - \underbrace{\frac{W(t)}{\phi_w}}_{\text{Immune decay}}, \quad (2)$$

where ψ_n and ψ_w denote the immune activation strengths, ϕ_n and ϕ_w represent the immune activity half-lives, and ι represents the theoretical maximum total iRBC

density, which determines the threshold at which immunity saturates. Experimental data from mouse infections indicate that indiscriminate immunity is quickly activated but short-lived, whereas targeted immunity has a longer activation period but a longer half-life [41]. Because N and W have been defined as proportions of cells cleared, they can be rewritten to reflect daily rates of clearance. For example, $N = 1 - e^{-x_N}$, where x_N is equal to the rate of clearance on a given day, and is defined as $-\ln(1 - N)$, with some rearranging of terms. A similar expression provides the daily clearance rate for targeted killing, $-\ln(1 - W)$.

[snip] Mimicking an experimental *P. chabaudi* infection, the host receives an initial dose of asexually-committed iRBC (I_0) to start the infection. We assume no parasites in this initial dose commit to gametocytogenesis and all produce asexual merozoites (M). Therefore, before the end of the first asexual development period ($t \leq \alpha$, where α is the development period of one day for this species), the dynamics of asexual iRBC (I) in the host are determined solely by the invasion of RBC by asexual merozoites, natural RBC death, maturation of the initial cohort of infected cells, and immune-mediated killing such that

$$\frac{dI}{dt} = \underbrace{\epsilon R(t)M(t)}_{\text{RBC invasion}} - \underbrace{\mu I(t)}_{\text{Natural death}} - \underbrace{I_0 \text{Beta}(sp, sp)S}_{\text{iRBC maturation}} - \underbrace{[-\ln(1 - N(t)) - \ln(1 - W(t))]I(t)}_{\text{Indiscriminate and targeted immunity}}. \quad (3)$$

The first term, $\epsilon R(t)M(t)$, represents the production of iRBC through asexual merozoite invasion of RBCs, which occurs at a rate of ϵ per day, given contact. Infected RBCs die at a background rate of μ , while $-\ln(1 - N(t))$ and $-\ln(1 - W(t))$ describe their rate of death due to indiscriminate and targeted immunity, respectively. The infection-age structure of the initial cohort of asexual iRBCs, denoted as $\text{Beta}(sp, sp)$, is modeled using a Beta distribution with a shape parameter $sp = 1$, allowing iRBC maturation to occur with a uniform probability throughout the development period [89]. The uniform probability of maturation for the initial cohort is computationally efficient and provided extremely similar results to more narrow age distributions in a previous modeling study [23]. The rate of asexual iRBC maturation depends on an exponential survival function S , which describes the proportion of iRBC that survives until maturation. S is defined by a cumulative hazard function that is calculated by integrating the total rate of death across the development period:

$$S = \exp\left[-\int_{t-\alpha}^t \underbrace{\mu}_{\text{Natural death}} + \underbrace{-\ln(1 - N(\omega)) - \ln(1 - W(\omega))}_{\text{Indiscriminate and targeted immunity}} d\omega\right]. \quad (4)$$

During the period $t \leq \alpha$, asexual merozoite production is described by

$$\frac{dM}{dt} = \underbrace{\beta I_0 \text{Beta}(sp, sp)S}_{\text{iRBC maturation}} - \underbrace{\mu_m M(t)}_{\text{Natural death}} - \underbrace{\epsilon R(t)M(t)}_{\text{RBC invasion}}, \quad (5)$$

where the first term represents the production of asexually-committed merozoites by asexual iRBC that have survived the duration of the developmental cycle and burst to produce β merozoites each. Asexual merozoites are lost by natural death at a rate of μ_m and are removed by their invasion of susceptible RBC.

After all the initially injected iRBC have matured or died ($t > \alpha$), the next cohort of asexual iRBC starts to reach maturity. Transmission investment is defined as the proportion, c , of this new generation that commits to gametocytogenesis, resulting in the production of β sexually-committed merozoites (M_g) per cell. These sexual

merozoites invade susceptible RBCs to produce sexual iRBCs (I_g). We assume that sexually-committed merozoites have the same death rate (μ_m) and rate of RBC invasion (ϵ) as asexual merozoites. The remaining proportion ($1 - c$) of iRBC produces asexual merozoites and continues the asexual replication cycle. Altogether, when $t > \alpha$, the dynamics of iRBC and merozoites are as follows:

$$\frac{dI}{dt} = \underbrace{\epsilon R(t)M(t)}_{\text{RBC invasion}} - \underbrace{\mu I(t)}_{\text{Natural death}} - \underbrace{[-\ln(1 - N(t)) - \ln(1 - W(t))]I(t)}_{\text{Indiscriminate and targeted immunity}} - \underbrace{\epsilon R(t - \alpha)M(t - \alpha)S}_{\text{iRBC maturation}} \quad (6)$$

$$\frac{dM}{dt} = \underbrace{\beta(1 - c)\epsilon R(t - \alpha)M(t - \alpha)S}_{\text{iRBC maturation (asexual)}} - \underbrace{\mu_m M(t)}_{\text{Natural death}} - \underbrace{\epsilon R(t)M(t)}_{\text{RBC invasion}} \quad (7)$$

$$\frac{dM_g}{dt} = \underbrace{\beta c \epsilon R(t - \alpha)M(t - \alpha)S}_{\text{iRBC maturation (sexual)}} - \underbrace{\mu_m M_g(t)}_{\text{Natural death}} - \underbrace{\epsilon R(t)M_g(t)}_{\text{RBC invasion}}. \quad (8)$$

The parameter governing transmission investment (c) represents the proportion of iRBCs that produce sexually-committed merozoites [and is further defined in the next section](#).

[snip]

Before the first cohort of sexual iRBC matures ($\alpha < t \leq \alpha + \alpha_g$), the dynamics of sexual iRBC is governed solely by the rate of RBC invasion by sexually-committed merozoites, natural RBC death, and immunity:

$$\frac{dI_g}{dt} = \underbrace{\epsilon R(t)M_g(t)}_{\text{RBC invasion}} - \underbrace{\mu I_g(t)}_{\text{Natural death}} - \underbrace{[-\ln(1 - N(t)) - \ln(1 - W(t))]I_g(t)}_{\text{Indiscriminate and targeted immunity}}. \quad (9)$$

When the first cohort of sexual iRBC matures ($t > \alpha + \alpha_g$), each sexual iRBC produces a single gametocyte, G , such that

$$\frac{dI_g}{dt} = \underbrace{\epsilon R(t)M_g(t)}_{\text{RBC invasion}} - \underbrace{\mu I_g(t)}_{\text{Natural death}} - \underbrace{[-\ln(1 - N(t)) - \ln(1 - W(t))]I_g(t)}_{\text{Indiscriminate and targeted immunity}} - \underbrace{\epsilon R(t - \alpha_g)M_g(t - \alpha_g)S_g}_{\text{iRBC maturation (sexual)}}, \quad (10)$$

$$\frac{dG}{dt} = \underbrace{\epsilon R(t - \alpha_g)M_g(t - \alpha_g)S_g}_{\text{iRBC maturation (sexual)}} - \underbrace{\mu_g G(t)}_{\text{Natural death}}, \quad (11)$$

where μ_g is the background death rate of gametocytes. The survival probability of sexual iRBCs during their development cycle is defined by the exponential survival function S_g , which is similar to that of asexual iRBC (S), except the cumulative hazard function is integrated over the length of sexual iRBC development, α_g :

$$S_g = \exp\left[-\int_{t-\alpha_g}^t \underbrace{\mu}_{\text{Natural death}} \underbrace{-\ln(1-N(\omega)) - \ln(1-W(\omega))}_{\text{Indiscriminate and targeted immunity}} d\omega\right]. \quad (12)$$

[snip]

Throughout the infection, the density of uninfected RBCs, R , is affected by erythropoiesis, natural death, indiscriminate immunity, and merozoite invasion, such that:

$$\begin{aligned} \frac{dR}{dt} = & \underbrace{\mu R_0}_{\text{Erythropoiesis}} + \underbrace{\rho(R_0 - R(t))}_{\text{Erythropoiesis upregulation}} - \underbrace{\mu R(t)}_{\text{Natural death}} \\ & - \underbrace{[-\ln(1-N(t))R(t)]}_{\text{Indiscriminate immunity}} - \underbrace{\epsilon R(t)[M(t) + M_g(t)]}_{\text{Merozoite invasion}}. \end{aligned} \quad (13)$$

In the absence of parasites, RBCs are replenished at a baseline rate, μ , that maintains a homeostatic equilibrium RBC concentration (R_0). During infection, the rate of erythropoiesis increases, and the extent of this upregulation is proportional to the deviation of RBC concentration from equilibrium [90], where ρ represents the maximum proportion of the RBC deficit recovered per day. RBCs are eliminated at a natural death rate of μ (which balances new production in the absence of infection), removed through indiscriminate immunity ($-\ln(1-N(t))$), or converted to iRBC via invasion by merozoites.

We parameterized our model based on prior work on malaria intrahost dynamics [23, 41] (Table 2). Because parameter values in Kamiya *et al.* [41] depend on initial dosage (I_0), we estimated our starting dose using the same method, which corresponds to approximately 10^4 - 10^5 iRBCs. Preliminary investigations indicate that varying the initial iRBC dose did not lead to quantitative differences in parasite relative fitness. Consequently, we parameterized $\psi_n, \psi_w, \phi_n, \phi_w$ and ρ using the median posterior estimates from [41] assuming an initial dosage of 10^4 iRBCs.

Modeling transmission investment

In our model, transmission investment (c) is determined by a transformed cue variable, f_t , perceived α days ago by a parasite infecting a cell, to account for NCC ($f_t(t-\alpha)$). Following the approach taken by Greischar *et al.* [23], we modeled the transmission investment reaction norm as the complementary log-log basis-spline with three degrees of freedom (B_3). Briefly, this method models the reaction norm as a linear combination of basis function, generating a continuous, smooth cubic curve. The relative contribution of each basis function (also known as the model coefficient) defines the overall shape of the reaction norm (see Figure 1C). We further modify the basis-spline by applying a complementary log-log transformation to constrain the transmission investment value to be between zero and one. This method allows us to use numerical optimization to search for the set of model parameters (i.e., reaction norm shape) that maximizes parasite fitness. Because cubic splines are fundamentally smooth, the resultant reaction norms may have difficulty representing sharp transitions. For that reason, we used untransformed and log-transformed cue values to build our models and capture greater variation in putative reaction norms.

For numerical optimization, the perceived cue must be within the bounds of the reaction norm. To achieve this, the perceived cue variable $f(t-\alpha)$ governing

Table 2. Parameter values used in the intrahost *P. chabaudi* model.

Parameter	Definition	Value	Unit	Source
R_0	Equilibrium RBC concentration	8.89×10^6	RBC/ μ L	[29]
μ	Background RBC production and RBC/iRBC mortality rate	0.025	day ⁻¹	[57]
p	Infection probability of RBC by merozoite invasion	8×10^{-6}	day ⁻¹	[91]
α	Asexual iRBC development period	1	day	[92]
α_g	Sexual iRBC development period	2	day	[93]
β	Burst size; number of merozoites produced per iRBC	5.721	merozoites	[94]
μ_m	Background merozoite mortality rate	48	day ⁻¹	[91]
μ_g	Background gametocyte mortality rate	4	day ⁻¹	[93]
I_0	Initial iRBC inoculum size	43.860	iRBC/ μ L	[23]
sp	Shape parameter dictating iRBC synchrony	1	NA	[89]
ψ_n	Activation strength for indiscriminate immunity	16.692	NA	[41]
ψ_w	Activation strength for targeted immunity	0.843	NA	[41]
ϕ_n	Half-life of indiscriminate immunity	0.035	day	[41]
ϕ_w	Half-life of targeted immunity	550.842	day	[41]
ι	Minimum iRBC concentration to induce maximum immunity	2.18×10^6	RBC/ μ L	[29]
ρ	Maximum proportion of RBC recovered per day	0.263	NA	[41]

transmission investment is transformed via a linear combination of Heaviside step functions (H) from the R package *CRONE* [95] that constrains the cue value to be between zero and the maximum value (max) defined in Table 3. In summary, the transmission investment function is defined as

$$c[f_t(t - \alpha)] = \exp(-\exp(B_3[f_t(t - \alpha)])), \quad (14)$$

where the Heaviside-transformed environmental cue $f_t(t - \alpha)$ is given by

$$f_t(t - \alpha) = H[f(t - \alpha)]f(t - \alpha) + H[f(t - \alpha) - \max][\max - f(t - \alpha)]. \quad (15)$$

For the best time-varying reaction norm, we defined the perceived cue as the time since infection, which has a range of 0-20 days and a step size of 0.001 (same as the simulation time step). In this case,

$$c(t) = \exp(-\exp(B_3(t))). \quad (16)$$

To incorporate dual cue perception, we used the *mgcv* package [96] to model transmission investment as the response variable of a generalized additive model. This model incorporates a full tensor product smooth of two environmental cues with nine degrees of freedom, represented as follows:

$$c[f_{t1}(t - \alpha), f_{t2}(t - \alpha)] = \exp(-\exp(\sum_{i=1}^3 \sum_{j=1}^3 \zeta_{ij} B_{3,1}[f_{t1}(t - \alpha)] B_{3,2}[f_{t2}(t - \alpha)])), \quad (17)$$

where transmission investment is governed by two Heaviside-transformed cues, f_{t1} and f_{t2} , perceived α days ago. The transmission investment surface (i.e., 2-dimensional reaction norm) is generated by taking the tensor product of the basis-splines of two cues ($B_{3,1}$ and $B_{3,2}$). The shape of this spline surface is defined by the model coefficients ζ_{ij} .

Quantifying parasite fitness

To quantify parasite fitness, we calculated the cumulative transmission potential of different transmission investment strategies, following the method described by Greischar *et al.* [23]. For a given time step, transmission potential, $\tau(t)$, is defined as the probability that a mosquito vector becomes infected after biting an infected host and has been shown empirically to be a sigmoid function of gametocyte density in this parasite species [10],

$$\tau(t) = \frac{3.08 \times 10^{-6} G(t)^{1.56}}{1 + 3.08 \times 10^{-6} G(t)^{1.56}}. \quad (18)$$

The cumulative transmission potential, or overall fitness of the parasite, is calculated by integrating transmission potential over the entire infection duration of δ days:

$$\text{fitness} = \int_0^\delta \tau(\omega) d\omega. \quad (19)$$

This fitness metric assumes that the mosquito population is at a constant abundance and continuously biting, for the duration of the infection. Although this is a simplification of the evolutionary pressures faced by malaria parasites [35], it enables us to focus our investigation at the within-host level, where transmission investment “decisions” are being made and environmental cues are being perceived by parasites. In our numerical simulations, we estimated fitness with a step size of 0.001 days unless stated otherwise.

Cue choice and perception

Transmission investment has been shown to vary in response to within-host competition [11, 12] and RBC age structure and availability [13, 17–19, 24], lysophosphatidylcholine level (a key component of the RBC plasma membrane) and exosome-like particles [97, 98]. Whether parasites are responding to these cues directly, or if they are instead a proxy for changes in some other within-host environmental factor is unclear. To capture some of the diversity of putative cues, we defined transmission investment as a function of one of ten different host and parasite-centric measures (Table 3). Additionally, we implemented dual cue models, allowing parasites to simultaneously perceive two different cues. As transmission investment in *P. falciparum* varies non-linearly with cue strength (e.g., *P. falciparum* in response to lysophosphatidylcholine concentration [97]), and within-host dynamics are known to vary only when infectious dose changes by orders of magnitude [29], we introduced non-linear cue perception, enabling parasites to perceive cues either unaltered or $\log_{10}$ -transformed. We excluded merozoite density as a cue due to its short half-life and the rapid pace of RBC invasion [99]. We also did not include measurements of immunity as cues, given that our model does not track individual components of immunity (rather, it tracks immune killing, phenomenologically), nor do we have sufficient empirical evidence detailing the relationship between different immunity components and transmission investment.

For all cues, we adopt a standardized range that spans the values typically observed when infecting immunologically naive mice with *P. chabaudi* [11], given by the minimum and maximum values in Table 3. All values that exceed the maximum bounds are “perceived” by the parasite as the maximum bound via the modified Heaviside transformation function.

Table 3. Within-host environmental cues tested and their range.

Cue (density per μL)	Abbreviation	Min. value	Max. value
Asexual iRBC	I	0	6×10^6
Asexual iRBC $\log_{10}$	I log	0	6.78
Sexual iRBC	Ig	0	6×10^6
Sexual iRBC $\log_{10}$	Ig log	0	6.78
Asexual iRBC + sexual iRBC	Total I	0	6×10^6
Asexual iRBC $\log_{10}$ + sexual iRBC $\log_{10}$	Total I log	0	6.78
Gametocyte	G	0	6×10^4
Gametocyte $\log_{10}$	G log	0	4.78
RBC	R	10^6	10^7
RBC $\log_{10}$	R log	6	7

Fitness optimization

Acute *P. chabaudi* infection tends to resolve within 20 days [100]. For all models, we therefore confined our simulations and optimizations to a 20-day infection period using the parameters detailed in Table 2 and using the *deSolve* package in R [101]. To ensure the convergence and continuity of output dynamics, we simulated the models with a small time step size of 0.001 days using the vode solver [102]. For the dual cue model, we used a larger time step of 0.01 days (which does not greatly alter parasite dynamics but improves computational speed immensely).

To investigate how transmission investment strategies affect parasite fitness, we performed numerical optimization of the basis-spline (single cue) or generalized additive model coefficients (dual cue) to maximize the cumulative transmission potential. We optimized the models using two methods to increase the chances of obtaining the global fitness optima. First, we performed local optimization following the methodology outlined by Greischar *et al.* [23] with the same arbitrary initial values of 0.5 for all spline coefficients, but using the more efficient parallelized L-BFGS-B algorithm (Limited-Memory Broyden–Fletcher–Goldfarb–Shanno algorithm-B) implemented in the *OptimParallel* package [103]. Second, we optimized each model using a hybrid approach to avoid convergence to a local fitness maximum. This technique involved initial global optimization using the Differential Evolution (DE) algorithm from the *DEoptim* package [104] (maximum iterations = 500, step tolerance = 50, differential weighting factor = 0.8, cross-over constant = 0.5 (single cue) or 0.9 (dual cue), population size = 40 (single cue) or 90 (dual cue), time step = 0.01 (single cue) or 0.05 (dual cue)). DE is an efficient and robust algorithm with comparable/better performance relative to other stochastic optimization algorithms [105]. Following optimization by the Differential Evolution algorithm, we further optimized the objective function using the L-BFGS-B algorithm to pinpoint a probable fitness maximum. For the differential Evolution algorithm, we used a parameter bound of $(-5, -500, -500, -500)$ and $(5, 500, 5000, 500)$ for the single cue model and $(-10, -500, -1000, -1000, -250, -500, -1000, -500, -250)$ and $(10, 500, 1000, 1000, 250, 500, 1000, 500, 250)$ for the dual cue model. These bounds effectively encompass the majority of basis-spline flexibility. From the two optimization results, we selected the parameter sets that resulted in the highest fitness.

For dual cue models, we implemented an additional optimization method that relies on the single cue models to inform the initial starting point. For dual cue models, we reasoned that the best single cue strategy out of the two provides a theoretical baseline fitness. In other words, the worst a dual-cue strategy should do is as good as the best

single cue (out of the two cues), while being unresponsive to the second cue. As such, we applied the L-BFGS-B algorithm to derive a dual cue strategy that “resembles” the best single cue strategy, specifically by minimizing an objective function with two terms: (i) the summed absolute difference between the dual and single cue reaction norms, and (ii) the total standard deviation in transmission investment across the worse cue. Inclusion of the second term ensures that transmission investment is not affected by variations in the worse cue. Together, the objective function is represented by:

$$\sum_{i=1}^{500} |\bar{c}_{i \text{ dual}} - c_{i \text{ single}}| + \sum_{i=1}^{500} \sum_{j=1}^{500} \sigma_{c_{ij} \text{ dual}}. \quad (20)$$

We first discretized the cue ranges into 500 evenly spaced values ($i = 1 \dots 500$, better performing cue; $j = 1 \dots 500$, worse performing cue). For each cue value with the index i , we calculate the deviation of the dual cue model using the absolute difference between $\bar{c}_{i \text{ dual}}$, the dual cue transmission investment averaged over the worse cue, and $c_{i \text{ single}}$, the corresponding transmission investment in the single cue model. The objective function also quantified the extent to which the worse cue affects transmission investment, specifically by summing the standard deviation across the worse cue value $\sum_{j=1}^{500} \sigma_{c_{ij} \text{ dual}}$ for each cue value indexed i .

Following this initial optimization, the resultant parameter is set as the starting point for an additional round of local optimization using the L-BFGS-B algorithm to maximize parasite fitness (same parameters as mentioned above).

As a further validation step, we compared the fitness of each putative single cue optimal strategy to the fitness obtained by 1000 randomly generated strategies. These random strategies were generated within the predefined bounds of $(-5, -500, -500, -500)$ and $(5, 500, 5000, 500)$, established previously to encompass much of the spline space. In no case was the putative best strategy outperformed by a random one (S1 Fig).

Semi-stochastic infection model

In order to explore the impact of individual infection-level variation on putative optimal transmission investment strategies, we incorporated the following random variables in the within-host model: maximum proportion of RBC deficit recovered per day (ρ), parasite burst size (β), immunity activation strengths (ψ_n and ψ_w), and immunity half-lives (ϕ_n and ϕ_w). For each iteration, random variables were drawn from 167 Markov chains (selected at regular intervals from a pool of 16,000 chains) used to estimate the posterior distribution in [41]. For the maximum proportion of RBC recovered per day and parasite burst size, the posterior distributions (originally centered at zero) are re-centered at the deterministic model values of 0.263 and 5.721, respectively. Since the paper from which we draw parameters explored the impact of dose [41], we only considered estimates pertaining to hosts infected with an initial dose of 10^4 parasites (constituting around 1000 parameter values), which corresponds to our choice for this model parameter (Table 2).

For each optimal transmission investment strategy (i.e., the best strategy for each cue or dual cue combination), we performed 20-day simulations with (i) all parameters allowed to vary from their deterministic default and (ii) only a single parameter varying. Here, we used a time step of 0.01 days, which does not change fitness substantially relative to performing the simulation with a time step of 0.001 days. [snip]

Disease mapping of laboratory mice infections

To explore the dynamics of host and parasite-centric cues in an *in vivo* setting, we curated data from publicly available sources of *P. chabaudi* infection dynamics in mice [11,42–45,47]. The curated dataset includes experiments that met the following criteria: a) single parasite strain infection, b) no administration of anti-malarial drugs, and c) availability of data on iRBC, RBC, and gametocyte densities. We then visualized two-dimensional disease maps (e.g., iRBC vs. RBC path plots) using the R package *ggplot2* [106].

Quantifying parasite virulence

We used the area encompassed within a disease map (where the x-axis displays total iRBC density and the y-axis displays uninfected RBC density, as in [46]) as a proxy for parasite virulence. To calculate this area, we first used the *chull* function [107] to define the convex hull, the smallest set of coordinates that encloses the points within the infection trajectory. We then used the *areapl* function from the *splancs* package [108] to quantify the area of the convex hull.

Supporting information

S1 Fig. Validation of transmission investment optimization. We conducted 1000 infection simulations for parasites sensing different cues, with each simulation using randomly generated transmission investment strategies. The fitness values of these simulations were compared to the fitness of parasites adopting the optimal transmission investment strategy. Each histogram represents the difference between the optimized fitness and the fitness obtained from the randomly generated strategy. The fitness of the random strategies is always lower.

S2 Fig. Transmission investment dynamics produced by parasites sensing different dual cue combinations. Transmission investment is plotted for all possible dual cue combinations, with heatmaps ranked in descending order (from top to bottom) based on their associated cumulative transmission potential.

S3 Fig. Terminal investment increases the fitness of dual cue-perceiving parasites. The cumulative transmission potential of parasites sensing time, the best performing dual cue combination (R log & I log), or the individual single cues (I log, R log) are indicated with a black cross, blue square, orange circle, and yellow triangle, respectively. The cumulative transmission potential is normalized with respect to the fitness of the best time-varying strategy. Note that time, the dual cue combination, and I log all permit an initial delay in transmission investment, while R log does not; the fitness advantage of this delayed strategy can be clearly viewed on these plots as fitness accrues faster. Only time and the dual cue combination lead to terminal investment, leading to additional fitness gains from day 18 onward.

S4 Fig. Logged RBC and logged iRBC form a looping trajectory *in silico*. Graphs display the two-dimensional reaction norm and infection trajectories for parasite sensing time with nine degrees of freedom. The trajectories depict the dynamics of the cues (indicated on the x- and y-axes) along with the corresponding transmission

investment (represented by the color of the line) **Note that the dynamics of gametocytes and sexual iRBC are hard to display completely given that these densities start off at zero.**[snip]. The distance between each hollow circle represents one day.[snip]

S5 Fig. RBC/logged RBC and logged iRBC form a looping trajectory *in vivo*. To identify combinations of within-host variables that exhibit a looping relationship, we plotted all possible permutations of experimentally derived *P. chabaudi* variables. Each line represents the dynamics of a single strain of infecting parasite. The line colour indicates the progression of time.

S6 Fig. Posterior distribution of parameter values used for the semi-stochastic model. The histograms display the distribution of 1000 parameter values drawn from 167 Markov chains derived by Kamiya *et al* [41]. The parameter values used in the deterministic model are indicated by the dotted line.

S7 Fig. Perceiving the optimal dual cue combinations increases parasite susceptibility to environmental and developmental stochasticity. We conducted 1000 simulations, each lasting 20 days, in which parasites adopted the optimal transmission investment strategy (from the deterministic model) using a semi-stochastic model where only one parameter was randomly drawn from the posterior distributions (see S6 Fig). For each parameter that is varied (indicated in the facet panels), we plot the geometric mean of the fitness on the y-axis and the fitness from the deterministic model on the x-axis. All fitness values are normalized with respect to the best time-varying strategy. The dashed line represents a 1:1 relationship between the x-axis and the y-axis; the line is not visible on the plot for burst size, since variation in this parameter has the greatest negative influence on fitness. [snip]

Acknowledgments

We thank members of the Mideo Lab for useful feedback on the project.

References

1. Jones OR, Scheuerlein A, Salguero-Gómez R, Camarda CG, Schaible R, Casper BB, et al. Diversity of ageing across the tree of life. *Nature*. 2014;505(7482):169–173. doi:10.1038/nature12789.
2. Gadgil M, Bossert WH. Life Historical Consequences of Natural Selection. *The American Naturalist*. 1970;104(935):1–24. doi:10.1086/282637.
3. Clutton-Brock TH, Albon SD, Guinness FE. Maternal dominance, breeding success and birth sex ratios in red deer. 1984;308(5957):358–360. doi:https://doi.org/10.1038/308358a0.
4. McNamara JM, Houston AI. State-dependent life histories. *Nature*. 1996;380(6571):215–221. doi:10.1038/380215a0.
5. Cotter SC, Ward RJS, Kilner RM. Age-specific reproductive investment in female burying beetles: independent effects of state and risk of death. *Functional Ecology*. 2011;25(3):652–660. doi:10.1111/j.1365-2435.2010.01819.x.

6. Velando A, Drummond H, Torres R. Senescent birds redouble reproductive effort when ill: confirmation of the terminal investment hypothesis. *Proceedings of the Royal Society B: Biological Sciences*. 2006;273(1593):1443–1448. doi:10.1098/rspb.2006.3480.
7. Carter LM, Kafsack BFC, Llinás M, Mideo N, Pollitt LC, Reece SE. Stress and sex in malaria parasites: Why does commitment vary? *Evolution, Medicine, and Public Health*. 2013;2013(1):135–147. doi:10.1093/EMPH/EOT011.
8. Churcher TS, Bousema T, Walker M, Drakeley C, Schneider P, Ouédraogo AL, et al. Predicting mosquito infection from *Plasmodium falciparum* gametocyte density and estimating the reservoir of infection. *eLife*. 2013;2:e00626. doi:10.7554/eLife.00626.
9. Bell AS, Huijben S, Paaijmans KP, Sim DG, Chan BHK, Nelson WA, et al. Enhanced Transmission of Drug-Resistant Parasites to Mosquitoes following Drug Treatment in Rodent Malaria. *PLOS ONE*. 2012;7(6):e37172. doi:10.1371/journal.pone.0037172.
10. Huijben S, Nelson WA, Wargo AR, Sim DG, Drew DR, Read AF. Chemotherapy, within-host ecology and the fitness of drug-resistant malaria parasites. *Evolution*. 2010;64(10):2952–2968. doi:10.1111/J.1558-5646.2010.01068.X.
11. Pollitt LC, Mideo N, Drew DR, Schneider P, Colegrave N, Reece SE. Competition and the Evolution of Reproductive Restraint in Malaria Parasites. *The American Naturalist*. 2011;177(3):358–367. doi:10.1086/658175.
12. Cameron A, Reece SE, Drew DR, Haydon DT, Yates AJ. Plasticity in transmission strategies of the malaria parasite, *Plasmodium chabaudi*: environmental and genetic effects. *Evolutionary Applications*. 2013;6(2):365. doi:10.1111/EVA.12005.
13. Schneider P, Greischar MA, Birget PLG, Repton C, Mideo N, Reece SE. Adaptive plasticity in the gametocyte conversion rate of malaria parasites. *PLOS Pathogens*. 2018;14(11):e1007371. doi:10.1371/JOURNAL.PPAT.1007371.
14. Williams JL. Stimulation of *Plasmodium falciparum* gametocytogenesis by conditioned medium from parasite cultures. *The American Journal of Tropical Medicine and Hygiene*. 1999;60(1):7–13. doi:10.4269/ajtmh.1999.60.7.
15. Dyer M, Day Karen P . Regulation of the rate of asexual growth and commitment to sexual development by diffusible factors from in vitro cultures of *Plasmodium falciparum*. *The American Journal of Tropical Medicine and Hygiene*. 2003;68(4):403–409. doi:10.4269/AJTMH.2003.68.403.
16. Mantel PY, Hoang A, Goldowitz I, Potashnikova D, Hamza B, Vorobjev I, et al. Malaria-Infected Erythrocyte-Derived Microvesicles Mediate Cellular Communication within the Parasite Population and with the Host Immune System. *Cell Host & Microbe*. 2013;13(5):521–534. doi:10.1016/J.CHOM.2013.04.009.
17. Trager W, Gill GS. Enhanced Gametocyte Formation in Young Erythrocytes by *Plasmodium falciparum* In Vitro. *The Journal of Protozoology*. 1992;39(3):429–432. doi:10.1111/j.1550-7408.1992.tb01476.x.

18. Trager W, Gill GS, Lawrence C, Nagel RL. *Plasmodium falciparum*: Enhanced Gametocyte Formation *in Vitro* in Reticulocyte-Rich Blood. *Experimental Parasitology*. 1999;91(2):115–118. doi:10.1006/expr.1998.4347.
19. Reece SE, Duncan AB, West SA, Read AF. Host cell preference and variable transmission strategies in malaria parasites. *Proceedings of the Royal Society B: Biological Sciences*. 2005;272(1562):511–517. doi:10.1098/rspb.2004.2972.
20. Buckling A, Crooks L, Read A. *Plasmodium chabaudi*: Effect of Antimalarial Drugs on Gametocytogenesis. *Experimental Parasitology*. 1999;93(1):45–54. doi:10.1006/expr.1999.4429.
21. Koella JC, Antia R. Optimal Pattern of Replication and Transmission for Parasites with Two Stages in Their Life Cycle. *Theoretical Population Biology*. 1995;47(3):277–291. doi:10.1006/tpbi.1995.1012.
22. Reece SE, Ramiro RS, Nussey DH. Synthesis: Plastic parasites: sophisticated strategies for survival and reproduction? *Evolutionary Applications*. 2009;2(1):11–23. doi:10.1111/j.1752-4571.2008.00060.x.
23. Greischar MA, Mideo N, Read AF, Bjørnstad ON. Predicting optimal transmission investment in malaria parasites. *Evolution*. 2016;70(7):1542–1558. doi:10.1111/EVO.12969.
24. Birget PLG, Repton C, O'Donnell AJ, Schneider P, Reece SE. Phenotypic plasticity in reproductive effort: malaria parasites respond to resource availability. *Proceedings of the Royal Society B: Biological Sciences*. 2017;284(1860). doi:10.1098/RSPB.2017.1229.
25. Clutton-Brock TH. Reproductive Effort and Terminal Investment in Iteroparous Animals. <https://doi.org/10.1086/284198>. 2015;123(2):212–229. doi:10.1086/284198.
26. Eichner M, Diebner HH, Molineaux L, Collins WE, Jeffery GM, Dietz K. Genesis, sequestration and survival of *Plasmodium falciparum* gametocytes: parameter estimates from fitting a model to malariatherapy data. *Transactions of The Royal Society of Tropical Medicine and Hygiene*. 2001;95(5):497–501. doi:10.1016/S0035-9203(01)90016-1.
27. Greischar MA, Mideo N, Read AF, Bjørnstad ON. Quantifying Transmission Investment in Malaria Parasites. *PLOS Computational Biology*. 2016;12(2):e1004718. doi:10.1371/JOURNAL.PCBI.1004718.
28. Pianka ER, Parker WS. Age-Specific Reproductive Tactics. *The American Naturalist*. 1975;109(968):453–464. doi:10.1086/283013.
29. Timms R, Colegrave N, Chan BHK, Read AF. The effect of parasite dose on disease severity in the rodent malaria *Plasmodium chabaudi*. *Parasitology*. 2001;123(Pt 1):1–11. doi:10.1017/S0031182001008083.
30. Metcalf CJE, Graham AL, Huijben S, Barclay VC, Long GH, Grenfell BT, et al. Partitioning regulatory mechanisms of within-host malaria dynamics using the effective propagation number. *Science*. 2011;333(6045):984–988. doi:10.1126/SCIENCE.1204588.

31. Barclay VC, Råberg L, Chan BHK, Brown S, Gray D, Read AF. CD4+T cells do not mediate within-host competition between genetically diverse malaria parasites. *Proceedings of the Royal Society B: Biological Sciences*. 2008;275(1639):1171–1179. doi:10.1098/rspb.2007.1713.
32. McKenzie FE, Bossert WH. An integrated model of *Plasmodium falciparum* dynamics. *Journal of Theoretical Biology*. 2005;232(3):411–426. doi:10.1016/j.jtbi.2004.08.021.
33. Bousema T, Drakeley C. Epidemiology and infectivity of *Plasmodium falciparum* and *Plasmodium vivax* gametocytes in relation to malaria control and elimination. *Clinical Microbiology Reviews*. 2011;24(2):377–410. doi:10.1128/CMR.00051-10.
34. Cao P, Collins KA, Zaloumis S, Wattanakul T, Tarning J, Simpson JA, et al. Modeling the dynamics of *Plasmodium falciparum* gametocytes in humans during malaria infection. *eLife*. 2019;8:e49058. doi:10.7554/eLife.49058.
35. Greischar MA, Beck-Johnson LM, Mideo N. Partitioning the influence of ecology across scales on parasite evolution. *Evolution*. 2019;73(11):2175–2188. doi:10.1111/EVO.13840.
36. Early AM, Camponovo F, Pelleau S, Cerqueira GC, Lazrek Y, Volney B, et al. Declines in prevalence alter the optimal level of sexual investment for the malaria parasite *Plasmodium falciparum*. *Proceedings of the National Academy of Sciences*. 2022;119(30):e2122165119. doi:10.1073/pnas.2122165119.
37. Lensen A, Bril A, van de Vegte M, van Gemert GJ, Eling W, Sauerwein R. *Plasmodium falciparum*: infectivity of cultured, synchronized gametocytes to mosquitoes. *Experimental Parasitology*. 1999;91(1):101–103. doi:10.1006/expr.1998.4354.
38. de Boor C. A Practical Guide to Splines. 1st ed. No. 27 in Applied Mathematical Sciences. Springer New York; 2001.
39. Dixon MWA, Thompson J, Gardiner DL, Trenholme KR. Sex in *Plasmodium*: a sign of commitment. *Trends in Parasitology*. 2008;24(4):168–175. doi:10.1016/J.PT.2008.01.004.
40. Birget PLG, Greischar MA, Reece SE, Mideo N. Altered life history strategies protect malaria parasites against drugs. *Evolutionary Applications*. 2018;11(4). doi:10.1111/eva.12516.
41. Kamiya T, Greischar MA, Schneider DS, Mideo N. Uncovering drivers of dose-dependence and individual variation in malaria infection outcomes. *PLOS Computational Biology*. 2020;16(10):e1008211. doi:10.1371/JOURNAL.PCBI.1008211.
42. Huijben S, Sim DG, Nelson WA, Read AF. Data from: The fitness of drug-resistant malaria parasites in a rodent model: multiplicity of infection. Dryad. 2011;.
43. Huijben S, Nelson WA, Wargo AR, Sim DG, Drew DR, Read AF. Data from: Chemotherapy, within-host ecology and the fitness of drug-resistant malaria parasites. Dryad. 2013;.

44. Huijben S, Bell AS, Sim DG, Tomasello D, Mideo N, Day T, et al. Data from: Aggressive chemotherapy and the selection of drug resistant pathogens. Dryad. 2014;.
45. Huijben S, Chan BHK, Nelson WA, Read AF. Data from: The impact of within-host ecology on the fitness of a drug-resistant parasite. Dryad. 2018;.
46. Torres BY, Oliveira JHM, Thomas Tate A, Rath P, Cumnock K, Schneider DS. Tracking Resilience to Infections by Mapping Disease Space. PLOS Biology. 2016;14(4):e1002436. doi:10.1371/JOURNAL.PBIO.1002436.
47. Reece SE, Drew DR, Gardner A. Sex ratio adjustment and kin discrimination in malaria parasites. Nature. 2008;453(7195). doi:10.1038/nature06954.
48. Moran NA. The Evolutionary Maintenance of Alternative Phenotypes. The American Naturalist. 1992;139(5):971–989. doi:10.1086/285369.
49. Scheiner SM. The genetics of phenotypic plasticity. VII. Evolution in a spatially-structured environment. Journal of Evolutionary Biology. 1998;11(3):303–320. doi:10.1046/j.1420-9101.1998.11030303.x.
50. De Jong G. Unpredictable selection in a structured population leads to local genetic differentiation in evolved reaction norms. Journal of Evolutionary Biology. 1999;12(5). doi:10.1046/j.1420-9101.1999.00118.x.
51. Schlaepfer MA, Runge MC, Sherman PW. Ecological and evolutionary traps. Trends in Ecology & Evolution. 2002;17(10):474–480. doi:10.1016/S0169-5347(02)02580-6.
52. Reed TE, Robin SW, Schindler DE, Hard JJ, Kinnison MT. Phenotypic plasticity and population viability: The importance of environmental predictability. Proceedings of the Royal Society B: Biological Sciences. 2010;277(1699). doi:10.1098/rspb.2010.0771.
53. Tufto J. The evolution of plasticity and nonplastic spatial and temporal adaptations in the presence of imperfect environmental cues. American Naturalist. 2000;156(2). doi:10.1086/303381.
54. Antia R, Yates A, de Roode JC. The dynamics of acute malaria infections. I. Effect of the parasite's red blood cell preference. Proceedings of the Royal Society B: Biological Sciences. 2008;275(1641):1449–1458. doi:10.1098/rspb.2008.0198.
55. Birget PLG, Kimberley F Prior, Savill NJ, Steer L, Reece SE. Plasticity and genetic variation in traits underpinning asexual replication of the rodent malaria parasite, *Plasmodium chabaudi*. Malaria Journal. 2019;18(1):222. doi:10.1186/s12936-019-2857-0.
56. Haydon DT, Matthews L, Timms R, Colegrave N. Top-down or bottom-up regulation of intra-host blood-stage malaria: do malaria parasites most resemble the dynamics of prey or predator? Proceedings of the Royal Society B: Biological Sciences. 2003;270(1512):289–298. doi:10.1098/rspb.2002.2203.
57. Miller MR, Råberg L, Read AF, Savill NJ. Quantitative Analysis of Immune Response and Erythropoiesis during Rodent Malarial Infection. PLOS Computational Biology. 2010;6(9):e1000946. doi:10.1371/JOURNAL.PCBI.1000946.

58. Kamiya T, Davis NM, Greischar MA, Schneider D, Mideo N. Linking functional and molecular mechanisms of host resilience to malaria infection. *eLife*. 2021;10. doi:10.7554/ELIFE.65846.
59. Lewontin RC, Cohen D. On population growth in a randomly varying environment. *Proceedings of the National Academy of Sciences of the United States of America*. 1969;62(4):1056–1060. doi:10.1073/pnas.62.4.1056.
60. Gillespie J. Polymorphism in random environments. *Theoretical Population Biology*. 1973;4(2):193–195. doi:10.1016/0040-5809(73)90028-2.
61. Pak D, Kamiya T, Greischar MA. Proliferation in malaria parasites: How resource limitation can prevent evolution of greater virulence. *Evolution*. 2024;78(7):1287–1301. doi:10.1093/evolut/qpae057.
62. Schneider P, Reece SE. The private life of malaria parasites: Strategies for sexual reproduction. *Molecular and Biochemical Parasitology*. 2021;244:111375. doi:10.1016/J.MOLBIOPARA.2021.111375.
63. Pigliucci M. Evolution of phenotypic plasticity: where are we going now? *Trends in Ecology & Evolution*. 2005;20(9):481–486. doi:10.1016/j.tree.2005.06.001.
64. Kochin BF, Bull JJ, Antia R. Parasite Evolution and Life History Theory. *PLOS Biology*. 2010;8(10):e1000524. doi:10.1371/journal.pbio.1000524.
65. McKenzie FE, Bossert WH. The Optimal Production of Gametocytes by *Plasmodium falciparum*. *Journal of Theoretical Biology*. 1998;193(3):419–428. doi:10.1006/jtbi.1998.0710.
66. Mideo N, Day T. On the evolution of reproductive restraint in malaria. *Proceedings of the Royal Society B: Biological Sciences*. 2008;275(1639):1217–1224. doi:10.1098/RSPB.2007.1545.
67. Charvin G, Oikonomou C, Siggia ED, Cross FR. Origin of Irreversibility of Cell Cycle Start in Budding Yeast. *PLOS Biology*. 2010;8(1):e1000284. doi:10.1371/journal.pbio.1000284.
68. Melen GJ, Levy S, Barkai N, Shilo B. Threshold responses to morphogen gradients by zero-order ultrasensitivity. *Molecular Systems Biology*. 2005;1(1):2005.0028. doi:10.1038/msb4100036.
69. Schneider DS. Tracing Personalized Health Curves during Infections. *PLOS Biology*. 2011;9(9):e1001158. doi:10.1371/journal.pbio.1001158.
70. Kasumovic MM. The multidimensional consequences of the juvenile environment: towards an integrative view of the adult phenotype. *Animal Behaviour*. 2013;85(5):1049–1059. doi:10.1016/j.anbehav.2013.02.009.
71. Dore AA, McDowall L, Rouse J, Bretman A, Gage MJG, Chapman T. The role of complex cues in social and reproductive plasticity. *Behavioral Ecology and Sociobiology*. 2018;72(8):124. doi:10.1007/s00265-018-2539-x.
72. Westneat DF, Potts LJ, Sasser KL, Shaffer JD. Causes and Consequences of Phenotypic Plasticity in Complex Environments. *Trends in Ecology & Evolution*. 2019;34(6):555–568. doi:10.1016/j.tree.2019.02.010.
73. Moller AP, Pomiankowski A. Why have birds got multiple sexual ornaments? *Behavioral Ecology and Sociobiology*. 1993;32(3):167–176. doi:10.1007/BF00173774.

74. Getty T. The maintenance of phenotypic plasticity as a signal detection problem. *The American Naturalist*. 1996;148(2):378–385.
75. DeWitt TJ. Costs and limits of phenotypic plasticity: Tests with predator-induced morphology and life history in a freshwater snail. *Journal of Evolutionary Biology*. 1998;11(4). doi:10.1007/s000360050100.
76. Anderson RM, May RM. Coevolution of hosts and parasites. *Parasitology*. 1982;85(2):411–426. doi:10.1017/S0031182000055360.
77. Alizon S, Hurford A, Mideo N, Van Baalen M. Virulence evolution and the trade-off hypothesis: history, current state of affairs and the future. *Journal of Evolutionary Biology*. 2009;22(2):245–259. doi:10.1111/j.1420-9101.2008.01658.x.
78. Mackinnon MJ, Read AF. Genetic Relationships between Parasite Virulence and Transmission in the Rodent Malaria *Plasmodium chabaudi*. *Evolution*. 1999;53(3):689–703. doi:10.1111/j.1558-5646.1999.tb05364.x.
79. Mackinnon MJ, Gaffney DJ, Read AF. Virulence in rodent malaria: host genotype by parasite genotype interactions. *Infection, Genetics and Evolution*. 2002;1(4):287–296. doi:10.1016/S1567-1348(02)00039-4.
80. Mackinnon MJ, Read AF. The effects of host immunity on virulence–transmissibility relationships in the rodent malaria parasite *Plasmodium chabaudi*. *Parasitology*. 2003;126(2):103–112. doi:10.1017/S003118200200272X.
81. Paul R, Lafond T, Müller-Graf C, Nithiuthai S, Brey P, Koella J. Experimental evaluation of the relationship between lethal or non-lethal virulence and transmission success in malaria parasite infections. *BMC Evolutionary Biology*. 2004;4(1):30. doi:10.1186/1471-2148-4-30.
82. Regoes RR, Nowak MA, Bonhoeffer S. Evolution of Virulence in a Heterogeneous Host Population. *Evolution*. 2000;54(1):64–71. doi:10.1111/j.0014-3820.2000.tb00008.x.
83. Williams PD. New Insights into Virulence Evolution in Multigroup Hosts. *The American Naturalist*. 2012;179(2):228–239. doi:10.1086/663690.
84. White PS, Choi A, Pandey R, Menezes A, Penley M, Gibson AK, et al. Host heterogeneity mitigates virulence evolution. *Biology Letters*. 2020;16(1):20190744. doi:10.1098/rsbl.2019.0744.
85. Hawking F, Wilson ME, Gammage K. Evidence for cyclic development and short-lived maturity in the gametocytes of *Plasmodium falciparum*. *Transactions of The Royal Society of Tropical Medicine and Hygiene*. 1971;65(5):549–559. doi:10.1016/0035-9203(71)90036-8.
86. Padilla DK, Adolph SC. Plastic inducible morphologies are not always adaptive: The importance of time delays in a stochastic environment. *Evolutionary Ecology*. 1996;10(1):105–117. doi:10.1007/BF01239351.
87. Hoi AG, Greischar MA, Mideo N. Limited impact of within-vector ecology on the evolution of malaria parasite transmission investment. *Frontiers in Malaria*. 2024;2. doi:10.3389/fmala.2024.1392060.

88. Mukherjee S, Bassler BL. Bacterial quorum sensing in complex and dynamically changing environments. *Nature reviews Microbiology*. 2019;17(6):371–382. doi:10.1038/s41579-019-0186-5.
89. Greischar MA, Read AF, Bjørnstad ON. Synchrony in malaria infections: How intensifying within-host competition can be adaptive. *American Naturalist*. 2014;183(2). doi:10.1086/674357.
90. Chang KH, Tam M, Stevenson MM. Modulation of the Course and Outcome of Blood-Stage Malaria by Erythropoietin-Induced Reticulocytosis. *The Journal of Infectious Diseases*. 2004;189(4):735–743. doi:10.1086/381458.
91. Mideo N, Barclay VC, Chan BHK, Savill NJ, Read AF, Day T. Understanding and predicting strain-specific patterns of pathogenesis in the rodent malaria *Plasmodium chabaudi*. *American Naturalist*. 2008;172(5):214–238. doi:doi/10.1086/591684.
92. Landau I, Boulard Y. Life Cycles and Morphology. In: Killick-Kendrick R, Peters W, editors. *Rodent Malaria*. Academic Press; 1978. p. 53–84.
93. Gautret P, Miltgen F, Gantier JC, Chabaud AG, Landau I. Enhanced gametocyte formation by *Plasmodium chabaudi* in immature erythrocytes: Pattern of production, sequestration, and infectivity to mosquitoes. *Journal of Parasitology*. 1996;82(6). doi:10.2307/3284196.
94. Mideo N, Savill NJ, Chadwick W, Schneider P, Read AF, Day T, et al. Causes of variation in malaria infection dynamics: Insights from theory and data. *American Naturalist*. 2011;178(6). doi:10.1086/662670.
95. Smith E, Evans G, Foadi J. An effective introduction to structural crystallography using 1D Gaussian atoms. *European Journal of Physics*. 2017;38(6):065501. doi:10.1088/1361-6404/AA8188.
96. Wood SN. mgcv: GAMs and generalized ridge regression for R. *R News*. 2001;1.
97. Brancucci NMB, Gerdt JP, Wang C, Niz MD, Philip N, Adapa SR, et al. Lysophosphatidylcholine Regulates Sexual Stage Differentiation in the Human Malaria Parasite *Plasmodium falciparum*. *Cell*. 2017;171(7):1532–1544.e15. doi:10.1016/J.CELL.2017.10.020.
98. Regev-Rudzki N, Wilson DW, Carvalho TG, Sisqueira X, Coleman BM, Rug M, et al. Cell-Cell Communication between Malaria-Infected Red Blood Cells via Exosome-like Vesicles. *Cell*. 2013;153(5):1120–1133. doi:10.1016/J.CELL.2013.04.029.
99. Boyle MJ, Wilson DW, Richards JS, Riglar DT, Tetteh KKA, Conway DJ, et al. Isolation of viable *Plasmodium falciparum* merozoites to define erythrocyte invasion events and advance vaccine and drug development. *Proceedings of the National Academy of Sciences of the United States of America*. 2010;107(32). doi:10.1073/pnas.1009198107.
100. Stephens R, Culleton RL, Lamb TJ. The contribution of *Plasmodium chabaudi* to our understanding of malaria. *Trends in Parasitology*. 2012;28(2). doi:10.1016/j.pt.2011.10.006.
101. Soetaert K, Petzoldt T, Setzer RW. Solving Differential Equations in R: Package deSolve. *Journal of Statistical Software*. 2020;33(9):1–25. doi:10.18637/jss.v033.i09.

102. Brown PN, Byrne GD, Hindmarsh AC. VODE: A Variable-Coefficient ODE Solver. <http://dxdoiorg/101137/0910062>. 2006;10(5):1038–1051. doi:10.1137/0910062.
103. Gerber F, Furrer R. optimParallel: An R Package Providing a Parallel Version of the L-BFGS-B Optimization Method. *The R Journal*. 2019;11(1):352–358. doi:10.32614/RJ-2019-030.
104. Mullen KM, Ardia D, Gil DL, Windover D, Cline J. DEoptim: An R package for global optimization by differential evolution. *Journal of Statistical Software*. 2011;40(6). doi:10.18637/jss.v040.i06.
105. Lampinen J, Storn R. Differential Evolution. In: Onwubolu GC, Babu BV, editors. *New Optimization Techniques in Engineering*. Studies in Fuzziness and Soft Computing. Berlin, Heidelberg: Springer; 2004. p. 123–166. Available from: https://doi.org/10.1007/978-3-540-39930-8_6.
106. Wickham H. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York; 2016. Available from: <https://ggplot2.tidyverse.org>.
107. R Core Team. *R: A language and environment for statistical computing*; 2021. Available from: <https://www.r-project.org/>.
108. Bivand R, Rowlingson B, Diggle P, Petris G, Eglen S. *splancs: Spatial and Space-Time Point Pattern Analysis*; 2024. Available from: <https://CRAN.R-project.org/package=splancs>.