

THE STABLE ADJUNCTION IN $\mathbb{A}^1$ -HOMOTOPY THEORY

AJAY SRINIVASAN

ABSTRACT. This paper demonstrates a monadicity result for the adjunction between the suspension spectrum and zeroth space functors in motivic stable homotopy theory. By collecting enough preliminary results, we are able to plug motivic stable homotopy theory into the general monadicity theorem of [KMZ26] to deduce the main results. This pursuit requires the verification of six properties of simplicial motivic homotopy theory, the hardest of which is a weak commutativity result between the zeroth space functor and realization of simplicial motivic spectra. Also essential to the main results is the elaboration presented here of how monadic algebras can be linked under an op-lax map of monads. We believe that an understanding of these simplicial properties, and the monadic framework explored here, may be helpful towards proving an operadic recognition principle for motivic infinite loop spaces.

CONTENTS

Introduction	1
Acknowledgements	4
1. Preliminaries	4
1.1. The Morel-Voevodsky framework	4
1.2. Po Hu's category of coordinate-free motivic spectra	6
1.3. The $\mathbb{A}^1$ -local stable model structure	9
1.4. The $(\Sigma^\infty, \Omega^\infty)$ adjunction	10
1.5. Simplicial axioms	13
2. The proofs of Theorems 0.1 and 0.2	13
2.1. Proof of the crude monadicity theorem	13
2.2. Op-lax maps of monads and abstract linking theory	15
2.3. Simplicial Γ -algebras and sharpened homotopical monadicity	20
3. Verification of the simplicial axioms	22
3.1. The easy axioms	22
3.2. The hard axiom	25
3.3. Realization-fibrations and the proof of Theorem 3.10	28
References	32

INTRODUCTION

Check. At the crux of stable homotopy theory is the adjunction between the suspension spectrum and zeroth space functors. The same is true in motivic stable

homotopy theory. This article analyzes the monadic structure of that stable adjunction in motivic homotopy theory. Consider a left adjoint functor $F: \mathcal{C} \rightarrow \mathcal{D}$ with right adjoint $G: \mathcal{D} \rightarrow \mathcal{C}$. The functor $\Gamma := GF$ on $\mathcal{C}$ has a canonical monad structure, where the product is induced by the counit ε of the adjunction. Γ is called the *adjunction monad*. Let $\Gamma[\mathcal{C}]$ denote the category of Γ -algebras in $\mathcal{C}$. For every $Y \in \mathcal{D}$, GY has a canonical Γ -action given by $G\varepsilon$, and as such G induces a functor $G_\Gamma: \mathcal{D} \rightarrow \Gamma[\mathcal{C}]$. G_Γ is also a right adjoint, with a left adjoint F_Γ . This left adjoint is quite distinct from F —it is a coequalized version of F with respect to Γ -actions. The *monadicity* of the adjunction (F, G) , in the general sense that we use the word here, will refer to the extent to which the right adjoint $G_\Gamma: \mathcal{D} \rightarrow \Gamma[\mathcal{C}]$ is an equivalence of categories. The result in [Bec25] provides conditions for (F_Γ, G_Γ) to be an adjoint equivalence. The resulting property is often called Beck monadicity or *classical* monadicity. However in many settings, including ours, the conditions for classical monadicity fail and a specificied homotopical approach is more enlightening. The recent framework of [KMZ26] provides one roadmap to such a form of *homotopical* monadicity. Roughly speaking, this paper here establishes that the stable adjunction in motivic homotopy theory is monadic *up to linked $\mathbb{A}^1$ -homotopy*. The analogous result in topology has roots dating back to [May72; May09], but is dealt more carefully in the general framework in [KMZ26]. The approach here is to collect enough results to plug motivic stable homotopy theory into the machine of [KMZ26]. Through our development, we resolve what may be an important step towards a conjectural operadic recognition principle for motivic infinite loop spaces [Voe00].

Let $\mathcal{T}$ denote the Morel-Voevodsky model category of algebraic spaces, this will be our category of *spaces*. In analogy with sequential Ω -spectra in topology, by *spectra* we could pick Voevodsky’s coordinatized $\mathbb{T}$ -spectra. However, for convenience with respect to the loop and suspension adjunction, we instead must work in a coordinate-free setting developed by [Hu03]. We will call Hu’s model category of coordinate-free spectra $\mathcal{S}$, and refer to its objects as *spectra*. The functor $\Omega^\infty: \mathcal{S} \rightarrow \mathcal{T}$, obtained by taking the “zeroth” space of a spectrum, is right adjoint. The *infinite loop spaces* will be those spaces that are in the image of Ω^∞ . The left adjoint of Ω^∞ , denoted $\Sigma^\infty: \mathcal{T} \rightarrow \mathcal{S}$, is the motivic suspension spectrum in our setting. The adjunction $(\Sigma^\infty, \Omega^\infty)$ is what we refer to as the *stable adjunction* in motivic homotopy theory, and it is the central object of study in this article.

Let $\Gamma = \Omega^\infty \Sigma^\infty$ denote the monad associated to the adjunction $(\Sigma^\infty, \Omega^\infty)$. Our first theorem utilizes the framework of [KMZ26] to produce a first step towards homotopical monadicity for the stable motivic adjunction. Analogous to the topological story, the main results here pertain to the subcategory $\mathcal{S}_c$ of *connective* spectra (also called *very effective* spectra elsewhere), which forms the homologically positive part of a t -structure on $\mathcal{S}$.

Theorem 0.1. *There is a functor $\mathbb{E}: \Gamma[\mathcal{T}] \rightarrow \mathcal{S}_c$ and a functor $\text{Bar}: \Gamma[\mathcal{T}] \rightarrow \mathcal{T}$, denoted $Y \mapsto \bar{Y}$, with the following properties.*

- (1) $\mathbb{E}: \Gamma[\mathcal{T}] \rightarrow \mathcal{S}_c$ and $\Omega_\Gamma^\infty: \mathcal{S}_c \rightarrow \Gamma[\mathcal{T}]$ preserve weak equivalences.
- (2) For $E \in \mathcal{S}_c$ and $Y \in \Gamma[\mathcal{T}]$, we have a natural homotopy equivalence ζ and weak equivalences γ and ξ in the following directions:

$$Y \xleftarrow{\zeta} \bar{Y} \xrightarrow{\gamma} \Omega_\Gamma^\infty \mathbb{E} Y$$

$$\mathbb{E}\Omega_{\Gamma}^{\infty}E \xrightarrow{\xi} E.$$

Theorem 0.1 however, does not establish an equivalence between the homotopy categories of Γ -algebras and connective spectra, and as such cannot be branded as homotopical monadicity. The functor Bar in the theorem alludes to the two-sided monadic bar resolution of Γ -algebras, built as the realization of a simplicial Γ -algebra—and as such we would have liked a functor $\text{Bar}: \Gamma[\mathcal{T}] \rightarrow \Gamma[\mathcal{T}]$. Similarly, we may have also liked for the weak equivalences in the theorem to be maps of Γ -algebras. In such a setting, we could immediately descend to homotopy categories to conclude that $(\mathbb{E}, \Omega_{\Gamma}^{\infty})$ induces an equivalence between homotopy categories. Unfortunately, due to a profound noncommutativity between Ω^{∞} and the realization of simplicial objects, there is no canonical Γ -action on $\bar{Y}$ for a Γ -algebra Y . So we do not quite have monadicity up to homotopy.

We resolve this gap by developing an abstract linking theory for monads connected by op-lax maps. Instantiating this abstract theory to our setting, we are able to concretely compare Γ -algebras and simplicial Γ -algebras. As a consequence, we show that although $\bar{Y}$ is not a Γ -algebra, it is the realization of a simplicial Γ -algebra whose algebra structure is *linked* via realization to that of the Γ -algebras $\Omega_{\Gamma}^{\infty}\mathbb{E}Y$ and Y . We also introduce the idea of *linked* maps between algebras. These are weakenings of algebra maps that still retain some monadic structure, by virtue of being maps *over* a linked algebra. Our second theorem is a strengthening of Theorem 0.1 with this linked language.

Theorem 0.2. *Let (Y, θ) be a Γ -algebra. There are maps*

$$Y \xleftarrow{\zeta} \bar{Y} \xrightarrow{\gamma} \Omega_{\Gamma}^{\infty}\mathbb{E}Y$$

where the following hold.

- (1) *Forgetting monadic structures, ζ is a homotopy equivalence and γ is a weak equivalence.*
- (2) *ζ and γ are maps that link $\bar{Y}$ to the Γ -algebras Y and $\Omega_{\Gamma}^{\infty}\mathbb{E}Y$ respectively.*
- (3) *For ν the homotopy inverse of ζ , the composite $\gamma \circ \nu$ is a linked weak equivalence between Y and $\Omega_{\Gamma}^{\infty}\mathbb{E}Y$.*

Theorem 0.2 allows us to make the following concrete statement about homotopy categories, a precisification of being “monadic up to linked $\mathbb{A}^1$ -homotopy.”

Corollary 0.3. *The full subcategory of Γ -algebras in $\mathcal{T}$ localized at the linked weak equivalences is equivalent to the homotopy category of connective spectra.*

The paper is structured as follows. In Section 1 we discuss preliminaries. Sections 1.1, 1.2, and 1.3 offer relevant background on Morel-Voevodsky algebraic spaces and on Po Hu’s coordinate-free spectra. Section 1.4 defines the stable adjunction and elaborates on some of its formal and homotopical properties, particularly with respect to the category $\mathcal{S}_c$ of connective spectra. The proofs of the main theorems, owing to [KMZ26], rely on methods from simplicial homotopy theory. In Section 1.5, we outline five properties of simplicial objects (**SA1-SA5**) that are essential to the proofs. The verification of these properties constitute all of Section 3. Properties **SA1-SA5** are easier to prove of the lot, and are verified in Section 3.1. The hardest and perhaps conceptually most insightful property in the list is **SA6**, which describes the weak commutativity of realization and Ω^{∞} . This is precisely the

source for the weak commutativity of Γ and $|-|$. The analogous result in topology plays a crucial role in the operadic recognition principle for infinite loop spaces [May72], and as such we believe resolves an important step towards an operadic recognition principle for motivic infinite loop spaces. Pursuing this line of inquiry towards operadic monads is the subject of a sequel paper. **SA6** is proved in Section 3.2 and Section 3.3, where Rezk’s reformulation of the compatibility of realizations with homotopy pullbacks [Rez14] plays a central part. With the preliminaries in place, Section 2 develops the required bar resolutions of Γ -algebras, and an abstract linking theory to complete the proofs of Theorems 0.1 and 0.2.

ACKNOWLEDGEMENTS

Much of this article was developed while I was a visitor at the University of Chicago in the summers of 2024 and 2025. I am indebted to Peter May for his support through the summer program at Chicago, and for his efforts in this project, which was very much joint work with him. My heartfelt gratitude also goes out to Alicia Lima, who co-advised me through the summer of 2024. I thank Paul Goerss for clarifications on simplicial path-space fibrations and Aravind Asok for pointing me to references on the commutativity of realization and homotopy pullback, and for continued discussion over the years on the social history of mathematics. Any errors in this article are mine alone.

1. PRELIMINARIES

1.1. The Morel-Voevodsky framework. This section recalls relevant details of the model category of algebraic spaces constructed by Morel and Voevodsky in [Voe96; MV99; Mor12]. Let k be a perfect field and S a smooth affine integral scheme of finite type over k (in particular, we ask that $S = \mathrm{Spec}(R)$ for a Noetherian integral domain R). We define Sm_S to be the category of smooth schemes of finite type over S . Whenever we refer to a scheme, we mean an object of Sm_S . The category Sm_S is infeasible for homotopy theory since it is not cocomplete, so one cannot use the objects of Sm_S to mean “spaces.” The fix for this is to consider simplicial sheaves over Sm_S as spaces. To speak of sheaves, one endows Sm_S with an appropriate topology. [MV99] identified the Nisnevich topology, finer than the Zariski but coarser than the Étale topology on Sm_S , to be a convenient choice. It is defined so that the covering sieves are those families of étale morphisms $\varphi_i: \{U_i\} \rightarrow X$ such that for any $x \in X$, there exists i and $u \in U_i$ with the corresponding map on residue fields being an isomorphism (that maps to x with the same residue field). We write $(\mathrm{Sm}_S)_{\mathrm{Nis}}$ to denote the site of Sm_S equipped with the Nisnevich topology. The definition of the Morel-Voevodsky category of spaces over S is then¹:

$$\mathrm{Spc}(S) := \mathrm{sSh}((\mathrm{Sm}_S)_{\mathrm{Nis}}).$$

We can also define the category of based spaces over S to be $\mathrm{Spc}(S)_\bullet = S \downarrow \mathrm{Spc}(S)$. We write $\mathcal{T}$ to mean this category and we call its objects spaces. The presheaf represented by an object $X \in \mathrm{Sm}_S$ is in fact a sheaf in the Nisnevich topology (this is due to [SGA4V2, VII.2a]). We also use $\mathcal{T}_k$ to mean the category

¹Notation: for a Grothendieck site $(\mathcal{C}, \tau)$ we write $\mathrm{sSh}(\mathcal{C}_\tau)$ to mean the category of simplicial **Set**-valued sheaves on $\mathcal{C}$. When there is no confusion, we will call the objects of $\mathrm{sSh}(\mathcal{C}_\tau)$ simplicial sheaves on $\mathcal{C}$.

$\mathrm{Spc}(k)_\bullet$ of spaces over k .

Now we look at some categorical properties of $\mathcal{T}$ before moving on to its homotopical properties and its model structure. First, $\mathcal{T}$ is complete and cocomplete. Second, $\mathcal{T}$ is tensored and cotensored over itself. That is to say, there is a smash product $\wedge: \mathcal{T} \times \mathcal{T} \rightarrow \mathcal{T}$, and there are internal Hom-space functors $\underline{\mathrm{Hom}}_{\mathcal{T}}(X, -)$ (right adjoint to $- \wedge X: \mathcal{T} \rightarrow \mathcal{T}$) defined in the usual way. Finally, by the co-Yoneda lemma we know that every presheaf over Sm_S is a small colimit of schemes. Passing to sheafification, which is left adjoint, we can conclude that every Nisnevich sheaf over Sm_S is a small colimit of schemes. Up to homotopy, we can make a similar statement about how to build simplicial sheaves. The following compactness property [Hu03, Lemma 17.2] is often used in conjunction with co-Yoneda type formulae to reduce some Hom-computations down to the scheme level (for example, with the case of spectrification in Section 1.2).

Proposition 1.1. *Every object of Sm_S is compact in $\mathcal{T}$*

We now recall aspects of the model structure on $\mathcal{T}$ and the notion of an $\mathbb{A}^1$ -homotopy defined in [MV99]. We write $\mathbb{A}^1$ to mean $\mathbb{A}_S^1$, the affine line over S . Two maps $f, g: X \rightarrow Y$ in $\mathcal{T}$ are said to be $\mathbb{A}^1$ -homotopic if there is a map $h: \mathbb{A}^1 \times X \rightarrow Y$ in $\mathrm{Spc}(k)$, such that the composite $X \xrightarrow{\iota_0} \mathbb{A}^1 \times X \xrightarrow{h} Y$ is f , and $X \xrightarrow{\iota_1} \mathbb{A}^1 \times X \rightarrow Y$ is g (where ι_j is induced by the inclusion of S into $\mathbb{A}_S^1$ as j).

Definition 1.2. A map $f: X \rightarrow Y$ in $\mathcal{T}$ is said to be a:

- (1) simplicial cofibration, if it is a monomorphism.
- (2) simplicial weak equivalence if for every $x^* \in \mathrm{Sm}_S$, the induced map $f^*: x^*(X) \rightarrow x^*(Y)$ is a weak equivalence of simplicial sets.²
- (3) simplicial fibration if it has the right lifting property (RLP) with respect to all trivial simplicial cofibrations.

The above definitions provide a model structure on $\mathcal{T}$ known as the simplicial model structure [Jar87, Corollary 2.7]. Denote by $\mathcal{H}_s(\mathcal{T})$ the homotopy category of $\mathcal{T}$ with respect to the simplicial model structure. The appropriate model structure that $\mathbb{A}^1$ -homotopy theory is set in comes from localizing the simplicial model structure with respect to $\mathbb{A}^1$ -homotopy. The first step towards such a localization is the notion of $\mathbb{A}^1$ -local spaces.

Definition 1.3. We say $X \in \mathcal{T}$ is $\mathbb{A}^1$ -local if for every $Y \in \mathcal{T}$, the map on hom-classes in the simplicial homotopy category $[Y, X]_{\mathcal{H}_s(\mathcal{T})} \rightarrow [Y \times \mathbb{A}^1, X]_{\mathcal{H}_s(\mathcal{T})}$ induced by the projection $Y \times \mathbb{A}^1 \rightarrow Y$ is a bijection.

Definition 1.4. A map $f: X \rightarrow Y$ in $\mathcal{T}$ is said to be an

- (1) $\mathbb{A}^1$ -cofibration if it is a monomorphism.
- (2) $\mathbb{A}^1$ -weak equivalence if for any $\mathbb{A}^1$ -local $Z \in \mathcal{T}$, the map on homotopy classes $f^*: [Y, Z]_{\mathcal{H}_s(\mathcal{T})} \rightarrow [X, Z]_{\mathcal{H}_s(\mathcal{T})}$ is a bijection.
- (3) $\mathbb{A}^1$ -fibration if it has the right lifting property with respect to all trivial cofibrations.

The $\mathbb{A}^1$ -local definitions above prescribe a model structure on $\mathcal{T}$ that admits functorial factorization [MV99, Theorem 2.21; Jar87, Section 2]. We denote by

²Here we think of an object x of a site S as a functor $x^*: \mathrm{Sh}(S) \rightarrow \mathbf{Set}$.

$\mathcal{H}(\mathcal{T})$ the homotopy category of $\mathcal{T}$ associated to this model structure. As promised earlier, we have the following characterization of a space up to simplicial weak equivalence as a colimit [Hu03, Lemma 17.3].

Proposition 1.5. *Every object in $\mathcal{T}$ is simplicially weak equivalent to a homotopy colimit of small colimits of smooth schemes.*

We close by discussing the $\mathbb{A}^1$ -homotopy sheaves, $\pi_n^{\mathbb{A}^1}(-, *)$, of a space. Let $X \in \mathcal{T}$. For $n = 0$, define $\pi_0^{\mathbb{A}^1}(X)$ to be the sheaf associated to the presheaf $U \mapsto [U, X]_{\mathcal{H}(\mathcal{T})}$ (for $U \in \text{Sm}_S$). For $n \geq 1$, and the given basepoint $*$: $S \hookrightarrow X$, define $\pi_n^{\mathbb{A}^1}(X, *)$ to be the sheaf associated to the presheaf $U \mapsto [\Sigma^n U_+, X]_{\mathcal{H}(\mathcal{T})}$. Here Σ^n denotes the simplicial suspension. We have the following motivic analog of Whitehead's theorem [MV99, Proposition 2.14].

Theorem 1.6. *A map of spaces $f: X \rightarrow Y$ in $\mathcal{T}$ is an $\mathbb{A}^1$ -weak equivalence iff it induces isomorphisms of sheaves $\pi_0^{\mathbb{A}^1}(X) \xrightarrow{\sim} \pi_0^{\mathbb{A}^1}(Y)$, and $\pi_n^{\mathbb{A}^1}(X, *) \xrightarrow{\sim} \pi_n^{\mathbb{A}^1}(Y, *)$ for every choice of basepoints $*$ for X and $*$ for Y (as objects in $\text{Spc}(S)$).*

A most helpful computational property of the $\mathbb{A}^1$ -motivic homotopy sheaves is that of the long exact sequence associated to a homotopy fiber sequence analogous to the one in topology.

Theorem 1.7. *If $F \rightarrow E \rightarrow B$ is a homotopy fiber sequence of spaces, then there is a natural long exact sequence of homotopy sheaves:*

$$\cdots \rightarrow \pi_{n+1}^{\mathbb{A}^1}(B) \rightarrow \pi_n^{\mathbb{A}^1}(F) \rightarrow \pi_n^{\mathbb{A}^1}(E) \rightarrow \pi_n^{\mathbb{A}^1}(B) \rightarrow \cdots$$

This is a natural long exact sequence first obtained at the presheaf level, passed through Nisnevich sheafification (which is exact). In the particular case of an inclusion of a subspace $\iota: A \hookrightarrow Y$, and the homotopy fiber of this inclusion, we get an analog of the long exact sequence of relative homotopy groups.

Fixing terminology for the rest of the paper, we say that $X \in \mathcal{T}$ is *connected* if $\pi_0^{\mathbb{A}^1}(X)$ is terminal in the category of sheaves of sets (i.e. a constant sheaf at a singleton set). Similarly we say $X \in \mathcal{T}$ is *n -connected* if $\pi_q^{\mathbb{A}^1}(X)$ is terminal for $q \leq n$ (in the category of sheaves of sets for $q = 0$, of groups for $q = 1$, and of abelian groups for $q \geq 2$). We say that a map $f: X \rightarrow Y$ in $\mathcal{T}$ is *n -connected* if its homotopy fiber Ff is $(n-1)$ -connected. By Theorem 1.7 this means that f induces isomorphisms $\pi_q^{\mathbb{A}^1}(X) \xrightarrow{\sim} \pi_q^{\mathbb{A}^1}(Y)$ for $q < n$ and a surjection $\pi_n^{\mathbb{A}^1}(X) \twoheadrightarrow \pi_n^{\mathbb{A}^1}(Y)$. We say that a pair (X, A) (i.e. an inclusion $\iota: A \rightarrow X$ in $\mathcal{T}$) is *n -connected* if ι is n -connected.

1.2. Po Hu's category of coordinate-free motivic spectra. In this section, we recall the construction of the coordinate-free spectra of [Hu03]. Recall that our base scheme was $S = \text{Spec}(R)$ for a Noetherian integral domain R . A natural choice for our coordinate-free indexing universe would be an infinite-dimensional affine space over S . We define our universe to be the countably infinite-dimensional R -module $\mathcal{U} = \bigoplus_{\infty} R$.³ We say Z is a *finite dimensional subspace* of $\mathcal{U}$ if it is a finitely generated projective submodule of $\mathcal{U}$ such that the inclusion $Z \hookrightarrow \mathcal{U}$ splits.

³For our purposes, it is okay to fix a universe here. [Hu03, Chapter 8] presents the theory for changes of universe.

Fix a basis $\{e_1, e_2, \dots\}$ for $\mathcal{U}$ and define T_n to be the free submodule generated by $\{e_1, \dots, e_n\}$ so that $\mathcal{U} \cong \operatorname{colim}_n T_n$ as an R -module. Now to think of $\mathcal{U}$ as a space over S , it suffices to note that every finitely generated projective R -module can be thought of as a vector bundle over S . Each T_n then corresponds to the trivial bundle $\mathbb{A}_S^n$, the n -dimensional affine space over S . To define $\mathcal{U}$ as a space (over S), we only need to write $\mathcal{U} \cong \operatorname{colim}_n T_n \cong \operatorname{colim}_n \mathbb{A}_S^n \cong \mathbb{A}_S^\infty$. Note that $\mathcal{U}$ is not really an object of Sm_S , but it is an ind-scheme over S . Similarly, a finitely generated projective submodule Z of $\mathcal{U}$ can be thought of as a finite-dimensional subbundle of $\mathcal{U}$ over S .

Our spectra will be indexed over the cofinite subspaces of $\mathcal{U}$. These are the subspaces of $\mathcal{U}$ of finite codimension, i.e. the split projective submodules of finite codimensions thought of as vector bundles over S . To put this definition on rigorous footing, P. Hu first defines the Grassmannian $Gr_S^{cof}(\mathcal{U})$ of cofinite subspaces of $\mathcal{U}$. We begin with the Grassmannian $Gr_k^{cof}(\mathcal{U}_k)$ over k of cofinite subspaces of $\mathbb{A}_k^\infty$, and write $Gr_S^{cof}(\mathcal{U}) := Gr_k^{cof}(\mathcal{U}_k) \times_{\operatorname{Spec}(k)} S$. For $N \in \mathbb{N}$ and $m \leq N$, define the Grassmannian $Gr_k^m(\mathbb{A}_k^N)$ to be the space (over k) of subspaces of $\mathbb{A}_k^N$ whose direct sum with $\mathbb{A}_k^m$ is $\mathbb{A}_k^N$ (here $\mathbb{A}_k^m \hookrightarrow \mathbb{A}_k^N$ by the inclusion into the first m coordinates of $\mathbb{A}_k^N$). Identifying $\mathcal{U}_k$ with $\operatorname{colim}_N \mathbb{A}_k^N$, we then define $Gr_k^{cof}(\mathcal{U}_k)$ to be the colimit $\operatorname{colim}_m \lim_{N \geq m} Gr_k^m(\mathbb{A}_k^N)$. The limit is taken over the restriction maps $Gr_k^m(\mathbb{A}_k^{N+1}) \rightarrow Gr_k^m(\mathbb{A}_k^N)$ taking V to $V \cap \mathbb{A}_k^N$ (here we think of $\mathbb{A}_k^N \subset \mathbb{A}_k^{N+1}$ as inclusion into the first N coordinates). We say that a map $V: S \rightarrow Gr_k^{cof}(\mathcal{U}_k)$ over k is a cofinite subspace of $\mathcal{U}$, in other words the cofinite subspaces of $\mathcal{U}$ precisely correspond to the split projective submodules of $\mathcal{U}$ of finite codimension. We similarly define an n -dimensional subspace of $\mathbb{A}_S^N$ to be a map $Z: S \rightarrow Gr_k(n, \mathbb{A}_k^N)$ (here $Gr_k(n, \mathbb{A}_k^N)$ is the Grassmannian of n -dimensional subspaces over k of $\mathbb{A}_k^N$). Again, this is just to say that the finite-dimensional subspaces are the finite-dimensional split projective submodules. For U an affine space over S , and subspaces V, W of U with trivial intersection, we denote by $V \oplus W$ the internal direct sum of V and W .

We are now ready to define our indexing category. Define the category of cofinite subspaces of $\mathcal{U}$, written $\mathcal{C}(\mathcal{U})$, to be the category whose objects are cofinite subspaces of $\mathcal{U}$ and morphisms $Z: U \rightarrow V$ are given by finite-dimensional subspaces $Z \subset U$ such that $V \oplus Z = U$. The composite of morphisms $Z: U \rightarrow V$ and $T: V \rightarrow W$ is the finite-dimensional subspace $Z \oplus T$ of U (note that $W \oplus (Z \oplus T) = U$). [Hu03, Lemma 2.3] shows that $\mathcal{C}(\mathcal{U})$ is a small directed category.

Let Z be a finite-dimensional space (alias finite-dimensional split projective module) over S . Thought of as a space, Z looks like a vector bundle over S equipped with a rational point $0: S \rightarrow Z$ (alias the zero-section of Z as a vector bundle). We define the space S^Z as the pushout

$$\begin{array}{ccc} Z \setminus \{0\} & \longrightarrow & Z \\ \downarrow & & \downarrow \\ S & \longrightarrow & S^Z. \end{array}$$

We think of S^Z as the one-point compactification of Z over S . For $X \in \mathcal{T}$, we write $\Omega^Z X$ to mean $\underline{\text{Hom}}_{\mathcal{T}}(S^Z, X)$. We will also denote the adjoint construction $S^Z \wedge X$ by $\Sigma^Z X$.

Definition 1.8. A *prespectrum* D is a family of spaces $\{D_U\}_{U \in \mathcal{C}(\mathcal{U})}$, together with morphisms $\rho_{V,Z}^U: D_U \rightarrow \Omega^Z D_V$ (for every morphism $Z: U \rightarrow V$ in $\mathcal{C}(\mathcal{U})$) satisfying the following conditions:

- a) $\rho_{U,0}^U = \text{id}_{D_U}$
- b) For every $Z: U \rightarrow V$, and $T: V \rightarrow W$ in $\mathcal{C}(\mathcal{U})$,

$$\rho_{W,Z \oplus T}^U = (\Omega^Z \rho_{W,T}^V) \circ \rho_{V,Z}^U.$$

The morphisms $\rho_{V,Z}^U$ are called the *structure maps* of D . Given prespectra D and D' , we call a collection of maps $\{f_U: D_U \rightarrow D'_U\}_{U \in \mathcal{C}(\mathcal{U})}$ whose elements are compatible with the structure maps, a map of prespectra from D to D' , written $f: D \rightarrow D'$. We will denote by $p\mathcal{S}$ the category of prespectra.

Definition 1.9. We say that a prespectrum E is a *spectrum* if its structure maps are all isomorphisms. A map of prespectra between two spectra is called a map of spectra. We denote by $\mathcal{S}$ the category of spectra.

All limits in $\mathcal{S}$ (and all limits and colimits in $p\mathcal{S}$) can be computed spacewise, just as in the topological case. The question of colimits, can be answered as follows. There is a forgetful functor $R: \mathcal{S} \rightarrow p\mathcal{S}$, given by forgetting that the structure maps are isomorphisms. R admits a left adjoint L that we call *spectrification*. Thanks to the Nisnevich topology (namely the compactness of S^Z as an object in $\mathcal{T}$ 1.1), the story here is much simpler than in topology where one must first pass through inclusion prespectra and Freyd's adjoint functor theorem.

Proposition 1.10. *There exists a functor $L: p\mathcal{S} \rightarrow \mathcal{S}$ called spectrification, defined by $D \mapsto LD$ where $(LD)_U = \text{colim}_{(V,Z) \in U \downarrow \mathcal{C}(\mathcal{U})} \Omega^Z D_V$ with prescribed structure maps $\rho_{W,T}^U: (LD)_U \xrightarrow{\sim} \Omega^T (LD)_W$. Here the colimit is taken over the maps $\Omega^Z D_V \rightarrow \Omega^{Z'} D_{V'}$ given by $\Omega^Z \rho_{V',T}^{V'}$ for maps $T: (V, Z) \rightarrow (V', Z')$.*

Proof. We only demonstrate why $\{(LD)_U\}_{U \in \mathcal{C}(\mathcal{U})}$ is actually a spectrum. It suffices to prescribe structure maps $\rho_{W,T}^U: (LD)_U \xrightarrow{\sim} \Omega^T (LD)_W$ for $T: U \rightarrow W$ in $\mathcal{C}(\mathcal{U})$. We begin with $(LD)_U$.

$$\begin{aligned} (LD)_U &\cong \text{colim}_{(V,Z) \in U \downarrow \mathcal{C}(\mathcal{U})} \Omega^{T \oplus Z} D_V \\ &\cong \Omega^T \text{colim}_{(V,Z) \in U \downarrow \mathcal{C}(\mathcal{U})} \Omega^Z D_V \\ &= \Omega^T (LD)_W \end{aligned}$$

Here we have used Proposition 1.1 to see that S^T is compact in $\mathcal{T}$, allowing us to commute Ω^T and the colimit. ■

As a left adjoint, spectrification commutes with colimits. Therefore we may compute colimits in $\mathcal{S}$ by first computing them at the prespectrum level and then spectrifying. The smash product $X \wedge E$ of a space X with a spectrum E is given by taking smash products spacewise and then spectrifying. We also have the function spectrum $F(X, E)$ defined by $F(X, E)_U = \underline{\text{Hom}}_{\mathcal{T}}(X, E_U)$ for each $U \in \mathcal{C}(\mathcal{U})$. These constructions show that $\mathcal{S}$ is tensored and cotensored over $\mathcal{T}$. One might wonder how the category $\mathcal{S}$ compares with Voevodsky's original construction of

the category of sequential $\mathbb{T}$ -spectra in [Voe96]. These concerns are addressed in [Hu03, Proposition 3.4 and Theorem 4.5]. These categories are in fact on the nose equivalent.

1.3. The $\mathbb{A}^1$ -local stable model structure. In this section, we recall P. Hu's definition of the stable $\mathbb{A}^1$ -local model structure on $\mathcal{S}$. There is first, a stable *simplicial* model structure on spectra, which itself comes from levelwise and stable simplicial model structures on the *prespectrum* level.

Definition 1.11. We say a map $f: D \rightarrow E$ in $p\mathcal{S}$ is a levelwise simplicial weak equivalence or fibration if for every $V \in \mathcal{C}(\mathcal{U})$, the V -th space map $f_V: D_V \rightarrow E_V$ is a simplicial weak equivalence or simplicial fibration of spaces (over k) respectively. We say f is a levelwise simplicial cofibration if it satisfies the left lifting property (LLP) with respect to every levelwise simplicial acyclic fibrations.

[Hu03, Proposition 5.10 and Chapter 15] show using the small objects argument that the distinguished classes of maps in the above definitions form a model structure on $p\mathcal{S}$. We call this model structure the *levelwise simplicial model structure* on $p\mathcal{S}$.

Definition 1.12. Let $f: D \rightarrow E$ be a map in $p\mathcal{S}$. We say f is a:

- (1) stable simplicial cofibration if it is a levelwise simplicial cofibration.
- (2) stable simplicial weak equivalence if it induces a bijection of colimits:

$$\operatorname{colim}_{(W,Z) \in V \downarrow \mathcal{C}(\mathcal{U})} [\Sigma^Z X_+, D_W]_{\mathcal{H}_s(\mathcal{T}_k)} \cong \operatorname{colim}_{(W,Z) \in V \downarrow \mathcal{C}(\mathcal{U})} [\Sigma^Z X_+, E_W]_{\mathcal{H}_s(\mathcal{T}_k)}$$

for every $V \in \mathcal{C}(\mathcal{U})$ and every $X \in \operatorname{Sms}$.

- (3) stable simplicial fibration if it satisfies the RLP with respect to all stable simplicial acyclic cofibrations.

The Bousfield-Friedlander theorem of [BF78, Theorem A.7] shows that the above definitions form a model structure on $p\mathcal{S}$. We call this model structure the *stable simplicial model structure* on $p\mathcal{S}$. We can now define the stable simplicial model structure on $\mathcal{S}$ by the following definitions.

Definition 1.13. Let $f: D \rightarrow E$ be a map in $\mathcal{S}$. We say f is a stable simplicial cofibration, weak equivalence, or fibration if it is one in the stable simplicial model structure on $p\mathcal{S}$.

The small objects argument from [Hu03, Chapter 15] applied to the above definitions show that they prescribe a model structure on $\mathcal{S}$. We call this model structure the *stable simplicial model structure* on $\mathcal{S}$, and we denote the associated homotopy category of $\mathcal{S}$ by $\mathcal{H}_s(\mathcal{S})$.

To $\mathbb{A}^1$ -localize the stable simplicial model structure on $\mathcal{S}$, we first recall P. Hu's notion of an $\mathbb{A}^1$ -homotopy in $\mathcal{S}$. Consider two maps $f, g: E \rightarrow G$ in $\mathcal{S}$. We say f and g are $\mathbb{A}^1$ -homotopic if there is a map $h: E \wedge \mathbb{A}_+^1 \rightarrow G$ whose composition with the maps $\iota_0, \iota_1: E = E \wedge S^0 \rightarrow E \wedge \mathbb{A}_+^1$ (one sending the non-base-point of S^0 to 0, and the other sending that point to 1 in $\mathbb{A}^1$) can be identified with f and g respectively. We define $\mathbb{A}^1$ -local spectra analogous to $\mathbb{A}^1$ -local spaces.

Definition 1.14. A spectrum $G \in \mathcal{S}$ is termed $\mathbb{A}^1$ -local if for every $E \in \mathcal{S}$, the map $E \wedge \mathbb{A}_+^1 \rightarrow E$ induced by the projection map $\mathbb{A}_+^1 \rightarrow S^0$ induces a bijection

of homotopy classes:

$$[E, G]_{\mathcal{H}_s(\mathcal{S})} \longrightarrow [E \wedge \mathbb{A}_+^1, G]_{\mathcal{H}_s(\mathcal{S})}$$

Definition 1.15. Let $f: E \longrightarrow E'$ be a map in $\mathcal{S}$. We say that f is an:

- (1) $\mathbb{A}^1$ -cofibration if it is a stable simplicial cofibration.
- (2) $\mathbb{A}^1$ -weak equivalence if for every $\mathbb{A}^1$ -local G , the map of homotopy classes $[E', G]_{\mathcal{H}_s(\mathcal{S})} \longrightarrow [E, G]_{\mathcal{H}_s(\mathcal{S})}$ is a bijection.
- (3) $\mathbb{A}^1$ -fibration if it satisfies the RLP with respect to every acyclic $\mathbb{A}^1$ -cofibration.

The above definitions form a model structure on $\mathcal{S}$ that we call the *stable $\mathbb{A}^1$ -local model structure* on $\mathcal{S}$ [Hu03, Theorem 6.4]. The proof is mostly analogous to that of [MV99, Theorem 2.2.21]. We denote by $\mathcal{H}(\mathcal{S})$ the homotopy category of $\mathcal{S}$ associated to the stable $\mathbb{A}^1$ -local model structure.

1.4. The $(\Sigma^\infty, \Omega^\infty)$ adjunction. In this section, we define the adjunction $(\Sigma^\infty, \Omega^\infty)$ and study some of its properties. For any $V \in \mathcal{C}(\mathcal{U})$, there is the V -th space functor $\Omega_V^\mathcal{U}: \mathcal{S} \longrightarrow \mathcal{T}$ given by $\Omega_V^\mathcal{U} E = E_V$. This functor has a left adjoint $\Sigma_V^\mathcal{U}: \mathcal{T} \longrightarrow \mathcal{S}$, called the V -th shift desuspension of the suspension spectrum, whose construction we now recall. Let $X \in \mathcal{T}$. For each $U \in \mathcal{C}(\mathcal{U})$, define the space $D_V(X)_U = \bigvee_{Z \oplus U = V} \Sigma^Z X$, and $D_V(X)_U = S$ (a point) otherwise. If $W \subset U \subset V$ (so that $V \longrightarrow U \longrightarrow W$ in $\mathcal{C}(\mathcal{U})$) such that $W \oplus T = U$, then for any Z with $Z \oplus U = V$, we have $(T \oplus Z) \oplus W = V$. This furnishes compatible maps:

$$\Sigma^T D_V(X)_V = \bigvee_{Z \oplus U = V} \Sigma^{T \oplus Z} X \longrightarrow \bigvee_{Z' \oplus W = V} \Sigma^{Z'} X = D_V(X)_W.$$

By the (Σ^Z, Ω^Z) adjunction, this furnishes a prespectrum $D_V(X) = \{D_V(X)_U\}_{U \in \mathcal{C}(\mathcal{U})}$. We define $\Sigma_V^\mathcal{U} X := L D_V(X)$.

It remains to show that $\Sigma_V^\mathcal{U}$ is left adjoint to $\Omega_V^\mathcal{U}$. It suffices to work on the level of prespectra. Let $f: D_V(X) \longrightarrow E$ be a map of prespectra (with $D_V(X)$ as defined above), i.e. a collection of compatible maps $f_U: D_V(X)_U \longrightarrow E_U$. The V -th map f_V is then just a map $X \longrightarrow E_V$. Conversely, let $g: X \longrightarrow E_V$ be a map of spaces. Let U be a cofinite subspace of V . For each map $Z: V \longrightarrow U$ (that is, a finite subspace $Z \subset V$ such that $Z \oplus U = V$), we obtain the map $\Sigma^Z X \xrightarrow{\Sigma^Z g} \Sigma^Z E_V \longrightarrow E_U$, where the last map is furnished by the structure maps of E . This prescription provides compatible maps $D_V(X)_U \longrightarrow E_U$, as needed.

We record the following key characterization of $\mathbb{A}^1$ -local spectra, and an analog of Whitehead's theorem for $\mathbb{A}^1$ -weak equivalences of spectra [Hu03, Lemma 6.12 and Proposition 6.13].

Theorem 1.16. *A spectrum G is $\mathbb{A}^1$ -local iff for every $X \in \mathrm{Spc}(S)$ that is a finite colimit of smooth schemes over S , and for every $V \in \mathcal{C}(\mathcal{U})$, the map*

$$[\Sigma_V^\mathcal{U} X_+, G]_{\mathcal{H}_s(\mathcal{S})} \longrightarrow [\Sigma_V^\mathcal{U} X_+ \wedge \mathbb{A}_+^1, G]_{\mathcal{H}_s(\mathcal{S})}$$

induced by the projection $\mathbb{A}_+^1 \longrightarrow S^0$ is a bijection.

Theorem 1.17. *A map $f: G \longrightarrow G'$ in $\mathcal{S}$ is an $\mathbb{A}^1$ -weak equivalence iff for every $X \in \mathrm{Spc}(S)$ that is a finite colimit of smooth schemes over S , and $V \in \mathcal{C}(\mathcal{U})$, the induced map: $[\Sigma_V^\mathcal{U} X_+, G]_{\mathcal{H}(\mathcal{S})} \longrightarrow [\Sigma_V^\mathcal{U} X_+, G']_{\mathcal{H}(\mathcal{S})}$ is a bijection.*

For the rest of this article, we will find it customary to drop the “ $\mathbb{A}^1$ -” when referring to the distinguished classes of maps in the model structures on $\mathcal{T}$ and $\mathcal{S}$.

We denote by Σ^∞ the functor $\Sigma_{\mathcal{U}}^\infty: \mathcal{T} \rightarrow \mathcal{S}$, and by Ω^∞ the functor $\Omega_{\mathcal{U}}^\infty: \mathcal{S} \rightarrow \mathcal{T}$. We call $\Sigma^\infty X$ the *suspension spectrum* of X , and $\Omega^\infty E$ the $\mathcal{U}$ -th *space* or the *zeroth space* of E . We have just shown that the pair $(\Sigma^\infty, \Omega^\infty)$ is an adjunction between $\mathcal{T}$ and $\mathcal{S}$. This is the categorical bedrock of this article. We iron out some formal preliminaries and notation now. Denote by Γ the adjunction monad $\Omega^\infty \Sigma^\infty$. The adjunction provides a unit map $\eta: \text{id}_{\mathcal{T}} \rightarrow \Gamma$, and a counit map $\varepsilon: \Sigma^\infty \Omega^\infty \rightarrow \text{id}_{\mathcal{S}}$. We write $\Gamma[\mathcal{T}]$ for the category of Γ -algebras in $\mathcal{T}$ (i.e. spaces in $\mathcal{T}$ with prescribed Γ -actions). It is easy to note that $\Omega^\infty E$ is a Γ -algebra for every $E \in \mathcal{S}$. The Γ -action is provided by the composite $\Gamma \Omega^\infty E = \Omega^\infty \Sigma^\infty \Omega^\infty E \xrightarrow{\Omega^\infty \varepsilon_E} \Omega^\infty E$. On the other hand, Σ^∞ is a Γ -functor in the sense that there is a natural map $\beta: \Sigma^\infty \Gamma \rightarrow \Sigma^\infty$ defined by $\varepsilon_{\Sigma^\infty X}: \Sigma^\infty \Gamma X \rightarrow \Sigma^\infty X$ for every $X \in \mathcal{T}$ (such that $\beta \circ \Sigma^\infty \mu = \beta \circ \beta$ and $\beta \circ \Sigma^\infty \eta = \text{id}$).

We denote by Ω_{Γ}^∞ the functor $\mathcal{S} \rightarrow \Gamma[\mathcal{T}]$ obtained by just applying Ω^∞ . Ω_{Γ}^∞ is also a right adjoint, but its left adjoint is *not* Σ^∞ . While maps of spectra $\Sigma^\infty X \rightarrow E$ (for an algebra X) do correspond to maps of spaces $X \rightarrow E_{\mathcal{U}}$, they do not a priori correspond to *maps of algebras*. The fix for this is to coequalize with respect to the Γ -action on Σ^∞ (seen as a Γ -functor) and the Γ -action on an algebra X . For each $X \in \Gamma[\mathcal{T}]$ with Γ -action θ , define $\Sigma_{\Gamma}^\infty X$ to be the coequalizer

$$\Sigma^\infty \Gamma X \xrightarrow[\Sigma^\infty \theta]{\beta_X} \Sigma^\infty X \longrightarrow \Sigma_{\Gamma}^\infty X.$$

This construction defines a functor $\Sigma_{\Gamma}^\infty: \Gamma[\mathcal{T}] \rightarrow \mathcal{S}$ left adjoint to Ω_{Γ}^∞ , and in doing so provides a coequalized adjunction $(\Sigma_{\Gamma}^\infty, \Omega_{\Gamma}^\infty)$ between $\Gamma[\mathcal{T}]$ and $\mathcal{S}$.

We now define a subcategory of connective spectra that is especially well-behaved with respect to our adjunction. Connective spectra are considered in analogy with their namesakes from topology, where they are the spectra with trivial negative homotopy groups. Let $\mathcal{S}_c$ denote the smallest full subcategory of $\mathcal{S}$ containing all suspension spectra of smooth schemes over S that is closed under homotopy colimits and extensions. Equivalently $\mathcal{S}_c$ is generated by colimits and extensions of $\Sigma_+^\infty X$ for $X \in \text{Sm}_S$ (here $\Sigma_+^\infty X$ is used as shorthand for $\Sigma^\infty X_+$). The objects of $\mathcal{S}_c$ are referred to as very effective spectra in the literature (see [SØ12, Definition 5.5]), but we call them Ω^∞ -*connective spectra* or *connective spectra* for short. $\mathcal{S}_c$ is not a triangulated category itself, but it forms the homologically positive part of a t -structure on $\mathcal{S}$. It is symmetric monoidal however under the smash product. By Proposition 1.5, and since Σ^∞ commutes with colimits as a left adjoint, $\Sigma^\infty: \mathcal{T} \rightarrow \mathcal{S}$ takes values in $\mathcal{S}_c$. We can now begin our homotopical investigation into the $(\Sigma^\infty, \Omega^\infty)$ adjunction.

Proposition 1.18. *If $f: X \rightarrow Y$ is a weak equivalence in $\mathcal{T}$, then $\Sigma^\infty f: \Sigma^\infty X \rightarrow \Sigma^\infty Y$ is a weak equivalence in $\mathcal{S}$.*

Proof. Let $f: X \rightarrow Y$ be a weak equivalence in $\mathcal{T}$, so that for every $\mathbb{A}^1$ -local space Z , the induced map $[Y, Z]_{\mathcal{H}_s(\mathcal{T})} \rightarrow [X, Z]_{\mathcal{H}_s(\mathcal{T})}$ is an isomorphism. Let E be an arbitrary $\mathbb{A}^1$ -local spectrum in $\mathcal{S}$. Note that $\Omega^\infty E = E_{\mathcal{U}}$ is $\mathbb{A}^1$ -local in $\mathcal{T}$. This is because for every $W \in \mathcal{T}$, $[\Sigma^\infty W, E] \rightarrow [\Sigma^\infty W \wedge \mathbb{A}^1, E]$ (induced

by the projection) is an isomorphism. By the adjunction then, we can see that $[W, E_{\mathcal{U}}] \rightarrow [W \wedge \mathbb{A}^1, E_{\mathcal{U}}]$ (again induced by the projection) is an isomorphism. Now we know that $[Y, E_{\mathcal{U}}]_{\mathcal{H}_s(\mathcal{T})} \rightarrow [X, E_{\mathcal{U}}]_{\mathcal{H}_s(\mathcal{T})}$ is an isomorphism. Passing to the adjunction, we have that $[\Sigma^\infty Y, E]_{\mathcal{H}_s(\mathcal{T})} \rightarrow [\Sigma^\infty X, E]_{\mathcal{H}_s(\mathcal{T})}$ is an isomorphism as well. $\blacksquare$

The analogous result for Ω^∞ follows from Theorem 1.17 and Theorem 1.6 (and Proposition 1.5, which allows us to pass to finite colimits of smooth schemes and use 1.17).

Proposition 1.19. *If $g: E \rightarrow E'$ is a weak equivalence in $\mathcal{S}$, then $g_{\mathcal{U}}: E_{\mathcal{U}} \rightarrow E'_{\mathcal{U}}$ is one in $\mathcal{T}$.*

Moreover, a stronger result holds in the case of connective spectra— Ω^∞ reflects weak equivalences.

Proposition 1.20. *Let $g: E \rightarrow E'$ be a map between connective spectra in $\mathcal{S}$. The map g is a weak equivalence in $\mathcal{S}$ iff $g_{\mathcal{U}}: E_{\mathcal{U}} \rightarrow E'_{\mathcal{U}}$ is one in $\mathcal{T}$.*

Proof. It only remains to prove the backward direction, beginning with a map $g: E \rightarrow E'$ such that $g_{\mathcal{U}}: E_{\mathcal{U}} \rightarrow E'_{\mathcal{U}}$ is a weak equivalence in $\mathcal{T}$. To reduce the problem, we invoke [Hu03, Proposition 6.10]—a descent principle for weak equivalences of spectra which states that (small) directed colimits of spectra preserve weak equivalences. In light of this reduction, we can restrict ourselves to the case of a weak equivalence $g_{\mathcal{U}}: (\Sigma^\infty Y_+)_{\mathcal{U}} \rightarrow (\Sigma^\infty Y'_+)_{\mathcal{U}}$, where Y and Y' are finite colimits of smooth schemes.

Let X be a finite colimit of smooth schemes. By assumption,

$$[\Sigma_{\mathcal{U}}^{\mathcal{U}} X_+, \Sigma^\infty Y_+]_{\mathcal{H}(\mathcal{S})} \rightarrow [\Sigma_{\mathcal{U}}^{\mathcal{U}} X_+, \Sigma^\infty Y'_+]_{\mathcal{H}(\mathcal{S})}$$

is a bijection. Let $V \in \mathcal{C}(\mathcal{U}) \setminus \{\mathcal{U}\}$. We observe that for any finite colimit of smooth schemes $\mathcal{Y}$, $[\Sigma_V^{\mathcal{U}} X_+, \Sigma^\infty \mathcal{Y}_+]_{\mathcal{H}(\mathcal{S})}$ is trivial. Note the following chain of isomorphisms:

$$\begin{aligned} [\Sigma_V^{\mathcal{U}} X_+, \Sigma^\infty \mathcal{Y}_+]_{\mathcal{H}(\mathcal{S})} &\cong [X_+, (\Sigma^\infty \mathcal{Y}_+)_V]_{\mathcal{H}(\mathcal{S})} \\ &\cong \left[X_+, \operatorname{colim}_{(W,Z) \in V \downarrow \mathcal{C}(\mathcal{U})} \Omega^Z \bigvee_{T \oplus W = \mathcal{U}} \Sigma^T \mathcal{Y}_+ \right]_{\mathcal{H}(\mathcal{S})}, \end{aligned}$$

where $T: \mathcal{U} \rightarrow V$ in $\mathcal{C}(\mathcal{U})$. Now using Proposition 1.1 on X_+ (or rather since finite colimits of compact objects are compact), we see that:

$$\begin{aligned} [\Sigma_V^{\mathcal{U}} X_+, \Sigma^\infty \mathcal{Y}_+]_{\mathcal{H}(\mathcal{S})} &\cong \operatorname{colim}_{(W,Z) \in V \downarrow \mathcal{C}(\mathcal{U})} \left[X_+, \Omega^Z \bigvee_{T \oplus W = \mathcal{U}} \Sigma^T \mathcal{Y}_+ \right]_{\mathcal{H}(\mathcal{S})} \\ &\cong \operatorname{colim}_{(W,Z) \in V \downarrow \mathcal{C}(\mathcal{U})} \left[\Sigma^Z X_+, \bigvee_{T \oplus W = \mathcal{U}} \Sigma^T \mathcal{Y}_+ \right]_{\mathcal{H}(\mathcal{S})}. \end{aligned}$$

To finish the argument, note that $\dim Z < \dim(T) = \operatorname{codim}(W)$ for every T such that $T \oplus W = \mathcal{U}$ (since $V \neq \mathcal{U}$), and $\Sigma^T \mathcal{Y}_+$ is $(\dim(T) - 1)$ -connected⁴. We conclude from Theorem 1.17 that g is a weak equivalence in $\mathcal{S}$. $\blacksquare$

⁴See remarks following [Mor12, Theorem 18].

1.5. Simplicial axioms. The proof of the homotopical monadicity theorem, using the argument of [KMZ26], will rely on a derived homotopy theory defined by bar resolutions of Γ -algebras. The two-sided monadic bar construction used here is built simplicially, and as such we list properties of simplicial objects in $\mathcal{T}$ and $\mathcal{S}$ that will be essential for the proof. The proofs of these properties constitute Section 3.

We will write $s\mathcal{C}$ for the simplicial objects in a category $\mathcal{C}$. For a functor $F: \mathcal{C} \rightarrow \mathcal{D}$, we will write F_* for the induced functor on simplicial objects $F_*: s\mathcal{C} \rightarrow s\mathcal{D}$ given by applying F objectwise on simplices.

SA1 There exist realization functors $|-|: s\mathcal{T} \rightarrow \mathcal{T}$ and $|-|: s\mathcal{S} \rightarrow \mathcal{S}$ that are both left adjoints. The composite $|c_*|$ where c_* is the constant simplicial object functor $\mathcal{C} \rightarrow s\mathcal{C}$, is the identity for $\mathcal{C} = \mathcal{T}, \mathcal{S}$.

SA2 $|-|$ preserves homotopies on $s\mathcal{T}$ and $s\mathcal{S}$.

SA3 $|-|$ preserves weak equivalences between Reedy cofibrant objects in both $s\mathcal{T}$ and $s\mathcal{S}$.

SA4 For $K_* \in s\mathcal{T}$, there is a natural isomorphism $\Sigma^\infty |K_*| \cong |\Sigma_*^\infty K_*|$.

SA5 $|-|: s\mathcal{S} \rightarrow \mathcal{S}$ takes levelwise connective spectra to connective spectra.

SA6 For $K_* \in s\mathcal{S}$, the natural map $\gamma: |\Omega_*^\infty K_*| \rightarrow \Omega^\infty |K_*|$, given by the adjoint to the composite $\Sigma^\infty |\Omega_*^\infty E_*| \cong |\Sigma_*^\infty \Omega_*^\infty E_*| \xrightarrow{|\varepsilon_*|} |E_*|$ is, for levelwise connective E_* , a weak equivalence in $\mathcal{T}$.

We will also frequently use the following fact (which follows formally from **SA1**).

Observation 1.21. For a simplicial object $X_* \in s\mathcal{T}$ and an object $J \in \mathcal{T}$, a map $f: X_0 \rightarrow J$ such that $fd_0 = fd_1$ extends to a map $\xi_*: X_* \rightarrow c_*J$ such that $\xi_0 = f$, and therefore induces a map $\xi := |\xi_*|: |X_*| \rightarrow J$ in $\mathcal{T}$.

It is a bit awkward to list properties that we have not yet proved as “preliminaries,” but the hope is that the reader will, until Section 3, take their validity for granted. Without saying much about their proofs here, we will only say that while **SA1-SA5** have relatively quicker proofs, **SA6** requires some labor. Consequently, we will sometimes refer to the former as the *easy axioms*, and the latter as the *hard axiom*.

2. THE PROOFS OF THEOREMS 0.1 AND 0.2

2.1. Proof of the crude monadicity theorem. The proof of crude monadicity presented here is due to an upcoming article [KMZ26] and discussions with J.P. May. First, we define the two-sided bar constructions that are essential for the proof. Let us return to the monad (Γ, μ, η) , where $\Gamma = \Omega^\infty \Sigma^\infty$. We noted in Section 1.2 that Σ^∞ was a Γ -functor in the sense that there was a natural map $\beta: \Sigma^\infty \Gamma \rightarrow \Sigma^\infty$ defined by $\beta_X = \varepsilon_{\Sigma^\infty X}$. We can also consider Γ as a Γ -functor with the action map $\mu = \Omega^\infty \varepsilon: \Gamma \Gamma \rightarrow \Gamma$. For any Γ -algebra (Y, θ) , we have the two-sided bar constructions:

$$\bar{Y} = B(\Gamma, \Gamma, Y) \text{ and } \mathbb{E}Y = B(\Sigma^\infty, \Gamma, Y),$$

where $\bar{Y} \in \mathcal{T}$, and $\mathbb{E}Y \in \mathcal{S}$. Recall that $B(\Gamma, \Gamma, Y)$ is the realization of the simplicial object in $\mathcal{T}$ (denoted $B_*(\Gamma, \Gamma, Y)$) whose q -simplices are given by $\Gamma^{q+1}Y$, and $B(\Sigma^\infty, \Gamma, Y)$ is the realization of the simplicial object in $\mathcal{S}$ (denoted $B_*(\Sigma^\infty, \Gamma, Y)$ or $\mathbb{E}_*Y$) whose q -simplices are given by $\Sigma^\infty \Gamma^q Y$. The zeroth face is given by the action of Γ (which is either μ or β), the i -th face for $0 < i < q$ is given by μ , and the q -th face is given by θ . The degeneracies are all given by $\eta: \text{id} \rightarrow \Gamma$.

By Observation 1.21, applied to $\theta: \Gamma Y \rightarrow Y$, there is a natural simplicial map $\zeta_*: B_*(\Gamma, \Gamma, Y) \rightarrow c_*Y$ which induces a natural map $\zeta: \bar{Y} \rightarrow Y$. We will also sometimes find it notationally convenient to write $B_*(\Gamma, \Gamma, Y) = \Omega_*^\infty \mathbb{E}_*Y$ so that $\bar{Y} = |\Omega_*^\infty \mathbb{E}_*Y|$. An extra degeneracy argument shows that:

Lemma 2.1. *For $Y \in \Gamma[\mathcal{T}]$, the natural map $\zeta: \bar{Y} \rightarrow Y$ in $\mathcal{T}$ induced by $\mu: \Gamma^{q+1}Y \rightarrow Y$ is a homotopy equivalence with homotopy inverse $\nu: Y \rightarrow \bar{Y}$ induced by the maps $\eta: Y \rightarrow \Gamma^{q+1}Y$.*

It is in this sense that we can see $\bar{Y}$ as a derived resolution of $Y \in \Gamma[\mathcal{T}]$. However, there is a conceptual gap in the above result. Y is a Γ -algebra, and the map $\mu: \Gamma^{q+1}Y \rightarrow Y$ is a map of Γ -algebras. We would like to say that $\bar{Y}$ is a Γ -algebra as well, and ζ is a map of Γ -algebras—so that we can really think of $\bar{Y}$ as a derived resolution of Y in the category of Γ -algebras. However, Ω^∞ does not commute with $|-|$ on the nose! It is therefore not at all clear that there is a canonically induced Γ -algebra structure on $\bar{Y}$. We will soon elaborate on the kind of formal structure that $\bar{Y}$ possesses.

Now we describe a conceptually dual procedure on $\mathcal{S}$. Let $\hat{\Gamma} := \Sigma^\infty \Omega^\infty$ denote the adjunction comonad and let $E \in \mathcal{S}$. Observe that $\mathbb{E}_* \Omega_F^\infty E$ has q -simplices $\hat{\Gamma}^{q+1}E$ and all faces given by ε . We define the following as a conceptual dual to $\bar{Y}$ for $Y \in \Gamma[\mathcal{T}]$.

$$\hat{E} := B(\Sigma^\infty, \Gamma, \Omega_F^\infty E) = \mathbb{E} \Omega_F^\infty E.$$

Since $\Sigma^\infty Z$ is always connective for $Z \in \mathcal{T}$, and $|-|$ takes levelwise connective spectra to connective spectra (by **SA5**), we see that $\mathbb{E}Y$ is connective for any $Y \in \Gamma[\mathcal{T}]$. Therefore $\hat{E}$ is connective for any $E \in \mathcal{S}$. The following should be seen as the dual to Lemma 2.1.

Lemma 2.2. *For $E \in \mathcal{S}$, the face maps $\varepsilon: \hat{\Gamma}^{q+1} \rightarrow \hat{\Gamma}^q$ induce a natural map $\xi: \hat{E} \rightarrow E$ in $\mathcal{S}$. If E is connective, then ξ is a weak equivalence.*

Proof. The first part of the statement is a direct application of Observation 1.21 to $\varepsilon: \hat{\Gamma}E \rightarrow E$. Now suppose E is connective. Since $\hat{E}$ is connective, by Proposition 1.20, it suffices to show that $\Omega^\infty \xi: \Omega^\infty \hat{E} \rightarrow \Omega^\infty E$ is a weak equivalence. Consider the following diagram, which commutes by passing to the adjoint and then appealing to Observation 1.21:

$$\begin{array}{ccc} \overline{\Omega_F^\infty E} = B(\Gamma, \Gamma, \Omega^\infty E) & \xrightarrow{\gamma} & \Omega^\infty B(\Sigma^\infty, \Gamma, \Omega^\infty E) = \Omega^\infty \hat{E} \\ & \searrow \zeta & \swarrow \Omega^\infty \xi \\ & \Omega^\infty E & \end{array}$$

where γ and ζ are weak equivalences (by **SA6** and Lemma 2.1). It follows by the two-out-of-three property that ξ is a weak equivalence as well. $\blacksquare$

Just as we had thought of $\bar{Y}$ as a derived resolution of $Y \in \Gamma[\mathcal{T}]$, we should think of $\hat{E}$ as a derived connective cover of E whose $\mathcal{U}$ th space is weak equivalent to the derived resolution of the $\mathcal{U}$ th space of E .

Piecing together the derived language from above, we get our first main result—a crude step towards homotopical monadicity.

Theorem 0.1. *There is a functor $\mathbb{E}: \Gamma[\mathcal{T}] \rightarrow \mathcal{S}_c$ and a functor $\text{Bar}: \Gamma[\mathcal{T}] \rightarrow \mathcal{T}$, denoted $Y \mapsto \bar{Y}$, with the following properties.*

- (1) $\mathbb{E}: \Gamma[\mathcal{T}] \rightarrow \mathcal{S}_c$ and $\Omega_F^\infty: \mathcal{S}_c \rightarrow \Gamma[\mathcal{T}]$ preserve weak equivalences.
- (2) For $E \in \mathcal{S}_c$ and $Y \in \Gamma[\mathcal{T}]$, we have a natural homotopy equivalence ζ and weak equivalences γ and ξ in the following directions:

$$Y \xleftarrow{\zeta} \bar{Y} \xrightarrow{\gamma} \Omega_F^\infty \mathbb{E}Y$$

$$\mathbb{E}\Omega_F^\infty E \xrightarrow{\xi} E.$$

Proof. The functors $\mathbb{E}$ and Bar have already been discussed. We note that Ω_F^∞ and $\mathbb{E}$ preserve weak equivalences since Ω^∞ and Σ^∞ preserve weak equivalences, and $|-|$ preserves weak equivalences between Reedy cofibrant spectra (this is **SA3**). Lemma 2.2 provides a natural weak equivalence $\xi: \hat{E} \rightarrow E$ for every connective spectrum E . For $Y \in \Gamma[\mathcal{T}]$, the homotopy equivalence $\zeta: \bar{Y} \rightarrow Y$ is given by Lemma 2.1, and the weak equivalence $\gamma: \bar{Y} \rightarrow \Omega_F^\infty \mathbb{E}Y$ is furnished by **SA6**. ■

The conceptual gap described following Lemma 2.1 is much more apparent now. We would really like to say that $\bar{Y}$ is a Γ -algebra such that ζ and γ are Γ -algebra maps. As a result, it is a priori not clear that Theorem 0.1 provides an equivalence between the homotopy categories of Γ -algebras and connective spectra. Nonetheless, with some conceptual investigation into the relation between Γ -algebras and simplicial Γ -algebras, we will establish a sharpened result.

2.2. Op-lax maps of monads and abstract linking theory. To further understand the relation between Γ -algebras and simplicial Γ -algebras, it will be helpful to think of simplicial Γ -algebras as themselves algebras over the simplicial adjunction monad Γ_* . The natural map $\gamma: |\Gamma_*| \rightarrow \Gamma|-|$ then relates these two monads—along with the data of $|-|$, it is an instance of an *op-lax map* (see Proposition 2.12). In this section, we develop an abstract theory of *linking* to understand how a general op-lax map between two monads can link their algebras.

We first recall the definition of an op-lax map between monads.

Definition 2.3. Let $(\mathbb{C}, \mu, \eta)$ and $(\mathbb{D}, \nu, \delta)$ be monads on categories $\mathcal{V}$ and $\mathcal{W}$ respectively. We say that a pair (F, α) with a functor $F: \mathcal{V} \rightarrow \mathcal{W}$ and natural transformation $\alpha: F\mathbb{C} \rightarrow \mathbb{D}F$ is an op-lax map of monads, or simply an op-lax map, if the following commute for every X .

$$\begin{array}{ccc} F\mathbb{C}CX & \xrightarrow{\alpha CX} & \mathbb{D}F\mathbb{C}X \xrightarrow{\mathbb{D}\alpha X} \mathbb{D}\mathbb{D}FX \\ \downarrow F\mu_X & & \downarrow \nu_{FX} \\ F\mathbb{C}X & \xrightarrow{\alpha_X} & \mathbb{D}FX \end{array} \quad \begin{array}{ccc} & FX & \\ F\eta_X \swarrow & & \searrow \delta_{FX} \\ F\mathbb{C}X & \xrightarrow{\alpha_X} & \mathbb{D}FX \end{array}$$

We will find it customary to write $(F, \alpha): \mathbb{C} \rightarrow \mathbb{D}$.

The following is the central definition to our linking theory.

Definition 2.4. Let $(\mathbb{C}, \mu, \eta)$ and $(\mathbb{D}, \nu, \delta)$ be monads on $\mathcal{V}$ and $\mathcal{W}$. Let $(F, \alpha): \mathbb{C} \rightarrow \mathbb{D}$. We say that algebras $(X, \theta) \in \mathbb{C}[\mathcal{V}]$ and $(Y, \varphi) \in \mathbb{D}[\mathcal{W}]$ are F -linked by $\beta: FX \rightarrow Y$ (called an F -linking map) if the following diagram commutes.

$$\begin{array}{ccccc} F\mathbb{C}X & \xrightarrow{\alpha_X} & \mathbb{D}FX & \xrightarrow{\mathbb{D}\beta} & \mathbb{D}Y \\ F\theta \downarrow & & & & \downarrow \varphi \\ FX & \xrightarrow{\beta} & & & Y \end{array}$$

When there is no ambiguity in the op-lax map (F, α) , we will just say that X and Y are linked, and that they are linked by the map β .

Example 2.5. Consider the case where $\mathcal{V} = \mathcal{W}$, $\mathbb{C} = \mathbb{D}$, and the op-lax map of monads is (id, id) on $\mathbb{C}$. See that two algebras (X, θ) and (Y, φ) in $\mathbb{C}[\mathcal{V}]$ are linked by $\beta: X \rightarrow Y$ iff the following commutes.

$$\begin{array}{ccc} \mathbb{C}X & \xrightarrow{\mathbb{C}\beta} & \mathbb{C}Y \\ \theta \downarrow & & \downarrow \varphi \\ X & \xrightarrow{\beta} & Y \end{array}$$

That is, iff β is a map of $\mathbb{C}$ -algebras! This tells us that maps of $\mathbb{C}$ -algebras are already linking maps. And more precisely that in the simplest case, linking maps should just be algebra maps. It is in this sense that we see our linking maps as generalizing maps of algebras.

Example 2.6. We illustrate an elementary example of the idea of linking. Every group homomorphism $\lambda: M \rightarrow N$ induces an op-lax map of monads from $M \times -$ to $N \times -$ on the category of sets. This op-lax map is given by $(\text{id}, \lambda \times \text{id})$. An id-linking map $\beta: X \rightarrow Y$ from an M -set X to an N -set Y is the same thing as a λ -equivariant map, i.e. $\beta(mx) = \lambda(m)\beta(x)$ for all $m \in M$ and $x \in X$.

We now compare our definition of linking with the definitions of [KMZ26].

Remark 2.7. Consider an op-lax map (F, α) from $\mathbb{C}$ to $\mathbb{D}$. Suppose we have $(X, \theta) \in \mathbb{C}[\mathcal{V}]$, $(FX, \psi) \in \mathbb{D}[\mathcal{W}]$, and $(Y, \varphi) \in \mathbb{D}[\mathcal{W}]$. A map $\beta: FX \rightarrow Y$ is a linking map iff the following commutes.

$$\begin{array}{ccccc} F\mathbb{C}X & \xrightarrow{\alpha_X} & \mathbb{D}FX & \xrightarrow{\mathbb{D}\beta} & \mathbb{D}Y \\ F\theta \downarrow & & & & \downarrow \varphi \\ FX & \xrightarrow{\beta} & & & Y \end{array}$$

Note that if β is a map of $\mathbb{D}$ -algebras, and the following triangle commutes, then β is a linking map.

$$\begin{array}{ccc} F\mathbb{C}X & \xrightarrow{\alpha_X} & \mathbb{D}FX \\ F\theta \downarrow & \swarrow \psi & \\ FX & & \end{array}$$

Note that this triangle commutes iff $\text{id}: FX \rightarrow FX$ is a linking map linking the algebras (X, θ) and (FX, ψ) . Whenever this is the case, [KMZ26, Definition 2.31]

calls X and FX linked. As a consequence of this remark, we obtain that X and FX are linked in the sense of [KMZ26] iff X and FX are linked via $\text{id}: FX \rightarrow FX$ in the sense of Definition 2.4. Nonetheless, the more abstract definition provided in [KMZ26, Definition 4.2] is equivalent to ours.

The following visual explanation concretizes what it means for algebras to be linked in our sense. Let us fix an op-lax map $(F, \alpha): \mathbb{C} \rightarrow \mathbb{D}$, algebras $(X, \theta) \in \mathbb{C}[\mathcal{V}]$ and $(Y, \varphi) \in \mathbb{D}[\mathcal{W}]$, and a linking map $\beta: FX \rightarrow Y$. Consider the unit diagrams for the algebras (X, θ) and (Y, φ) .

$$\begin{array}{ccc} X & \xrightarrow{\eta_X} & \mathbb{C}X \\ & \searrow & \downarrow \theta \\ & & X \end{array} \quad \begin{array}{ccc} Y & \xrightarrow{\delta_Y} & \mathbb{D}Y \\ & \searrow & \downarrow \varphi \\ & & Y \end{array}$$

Applying F to the unit diagram for X , the map $\beta: FX \rightarrow Y$ links together the unit diagrams to produce the commutative diagram below.

$$\begin{array}{ccccc} & & Y & \xrightarrow{\delta_Y} & \mathbb{D}Y \\ & \nearrow \beta & & \searrow & \downarrow \varphi \\ & & & & Y \\ & \nearrow \beta & & \searrow \mathbb{D}\beta \circ \alpha_X & \\ FX & \xrightarrow{\quad} & F\mathbb{C}X & & \\ & \searrow F\eta_X & \downarrow F\theta & & \\ & & FX & \xrightarrow{\quad} & Y \end{array}$$

To see that this prism commutes, it suffices to see that its top face commutes. We can write the top face as the composite:

$$\begin{array}{ccccc} FX & \xrightarrow{\quad \beta \quad} & Y & & \\ F\eta_X \downarrow & \searrow \delta_{FX} & & \searrow \delta_Y & \\ F\mathbb{C}X & \xrightarrow{\alpha_X} & \mathbb{D}FX & \xrightarrow{\mathbb{D}\beta} & \mathbb{D}Y \end{array}$$

The triangle in this diagram commutes since it is the unit diagram of the op-lax map (F, α) , and the remaining trapezoid is a naturality square for δ .

Similarly, we have the product diagrams for X and Y :

$$\begin{array}{ccc} \mathbb{C}CX & \xrightarrow{\mathbb{C}\theta} & \mathbb{C}X \\ \mu_X \downarrow & & \downarrow \theta \\ \mathbb{C}X & \xrightarrow{\theta} & X \end{array} \quad \begin{array}{ccc} \mathbb{D}DY & \xrightarrow{\mathbb{D}\varphi} & \mathbb{D}Y \\ \nu_Y \downarrow & & \downarrow \varphi \\ \mathbb{D}Y & \xrightarrow{\varphi} & Y \end{array}$$

Once more, applying F to the product diagram for X , the map $\beta: FX \rightarrow Y$ links together the product diagrams to produce the commuting diagram below.

$$\begin{array}{ccccccc}
 & & & & \mathbb{D}\mathbb{D}Y & \xrightarrow{\mathbb{D}\varphi} & \mathbb{D}Y \\
 & & & & \downarrow \nu_Y & & \downarrow \varphi \\
 & & & \mathbb{D}\mathbb{D}FX & \xrightarrow{\mathbb{D}\beta} & \mathbb{D}Y & \xrightarrow{\varphi} & Y \\
 & & \mathbb{D}\alpha_X \circ \alpha_{\mathbb{C}X} \nearrow & & \mathbb{D}\beta \circ \alpha_X \nearrow & & & \\
 FCCX & \xrightarrow{F\mathbb{C}\theta} & FCX & \xrightarrow{F\theta} & FX & \xrightarrow{\beta} & Y \\
 \downarrow F\mu_X & & \downarrow F\theta & & & & \\
 FCX & \xrightarrow{F\theta} & FX & & & &
 \end{array}$$

The front and back faces of this cuboid commute since they are the product diagrams for X and Y . The right face and the bottom face also commute since X and Y are linked by β by assumption. We can decompose the top face into the following commutative diagrams:

$$\begin{array}{ccccccc}
 FCCX & \xrightarrow{\alpha_{\mathbb{C}X}} & \mathbb{D}FCX & \xrightarrow{\mathbb{D}\alpha_X} & \mathbb{D}\mathbb{D}FX & \xrightarrow{\mathbb{D}\mathbb{D}\beta} & \mathbb{D}\mathbb{D}Y \\
 \downarrow F\mathbb{C}\theta & & \downarrow \mathbb{D}F\theta & & & & \downarrow \mathbb{D}\varphi \\
 FCX & \xrightarrow{\alpha_X} & \mathbb{D}FX & \xrightarrow{\mathbb{D}\beta} & \mathbb{D}Y. & &
 \end{array}$$

The left square commutes as a naturality square for α , and the right square commutes since it is $\mathbb{D}$ applied to the linking diagram for β . Similarly the left face of the cuboid also decomposes as:

$$\begin{array}{ccccccc}
 FCCX & \xrightarrow{\alpha_{\mathbb{C}X}} & \mathbb{D}FCX & \xrightarrow{\mathbb{D}\alpha_X} & \mathbb{D}\mathbb{D}FX & \xrightarrow{\mathbb{D}\mathbb{D}\beta} & \mathbb{D}\mathbb{D}Y \\
 \downarrow F\mu_X & & & & \downarrow \nu_{FX} & & \downarrow \nu_Y \\
 FCX & \xrightarrow{\alpha_X} & \mathbb{D}FX & \xrightarrow{\mathbb{D}\beta} & \mathbb{D}Y. & &
 \end{array}$$

The left square commutes as the product diagram for the op-lax map (F, α) , and the right square commutes since it is the naturality square for ν .

In essence, what we have tried to motivate here is this: to say the algebras (X, θ) and (Y, φ) are linked, is to say in a precise diagrammatic sense that their algebra structures are linked together via the linking map β .

Examples 2.8 provides a general simplification of the kind that one may have hoped for in the loop space story, and indicates how it would have allowed us to evade the linking notion. The precise failure of the relevant hypothesis in the loop space story is what necessitates a general notion beyond maps of algebras.

Example 2.8. Consider again an op-lax map $(F, \alpha): \mathbb{C} \rightarrow \mathbb{D}$ and algebras $(X, \theta) \in \mathbb{C}[\mathcal{V}]$ and $(Y, \varphi) \in \mathbb{D}[\mathcal{W}]$. Suppose $\alpha_X: FCCX \cong \mathbb{D}FX$ is an isomorphism. There is then a canonical $\mathbb{D}$ -algebra structure on FX given by $\psi = F\theta \circ \alpha_X^{-1}$. This definition satisfies the triangle condition from Remark 2.7 manifestly. We infer that a map $\beta: FX \rightarrow Y$ is a linking map iff it is a map of $\mathbb{D}$ -algebras.

In what follows, we examine a general situation that is closer to our loop space story, with the idea of a *section* of an op-lax map. Remark 2.10 provides a broad class of linking maps that we will use to sharpen our monadicity result in Section 2.3.

Definition 2.9. Let $(F, \alpha): \mathbb{C} \rightarrow \mathbb{D}$ be an op-lax map of monads from $(\mathbb{C}, \mu, \eta)$ (on $\mathcal{V}$) to $(\mathbb{D}, \nu, \delta)$ (on $\mathcal{W}$). We say that (F, α) admits a section if there exists an op-lax map $(G, \text{id}): \mathbb{D} \rightarrow \mathbb{C}$ such that $FG = \text{id}$ and $\alpha_{GY} = \text{id}$ for every $Y \in \mathcal{W}$. Such an op-lax map is called a section of (F, α) .

Remark 2.10. Let $(F, \alpha): \mathbb{C} \rightarrow \mathbb{D}$ be an op-lax map with section (G, id) . If (Y, φ) is a $\mathbb{D}$ -algebra, then $(GY, G\varphi)$ is automatically a $\mathbb{C}$ -algebra. For any other $\mathbb{D}$ -algebra (Y', φ') , a map $\beta: Y \rightarrow Y'$ is a linking map iff it is a map of $\mathbb{D}$ -algebras.

$$\begin{array}{ccccc} \mathbb{D}Y & \xlongequal{\quad} & F\mathbb{C}GY & \xrightarrow{\alpha_{GY}} & \mathbb{D}FGY & \xrightarrow{\mathbb{D}\beta} & \mathbb{D}Y' \\ \varphi \downarrow & & \varphi \downarrow & & & & \downarrow \varphi' \\ Y & \xlongequal{\quad} & FGY & \xrightarrow{\quad \beta \quad} & Y' & & \end{array}$$

Now consider algebras $(X, \theta) \in \mathbb{C}[\mathcal{V}]$ and $(Y, \varphi) \in \mathbb{D}[\mathcal{W}]$. We show that if $f: X \rightarrow GY$ is a map of $\mathbb{C}$ -algebras, then $\beta := F(f): FX \rightarrow Y$ is a linking map. Since f is a map of $\mathbb{C}$ -algebras, the following commutes.

$$\begin{array}{ccc} F\mathbb{C}X & \xrightarrow{F\mathbb{C}f} & \mathbb{D}Y \\ F\theta \downarrow & & \downarrow \varphi \\ FX & \xrightarrow{Ff} & Y \end{array}$$

Moreover by the naturality of α (applied to the morphism $f: X \rightarrow GY$), we have the following commutative diagram.

$$\begin{array}{ccc} F\mathbb{C}X & \xrightarrow{F\mathbb{C}f} & \mathbb{D}Y \\ & \searrow \alpha_X & \nearrow \mathbb{D}Ff \\ & \mathbb{D}FX & \end{array}$$

We conclude that the following commutes and therefore that β is a linking map.

$$\begin{array}{ccc} F\mathbb{C}X & \xrightarrow{\alpha_X} & \mathbb{D}FY & \xrightarrow{\mathbb{D}\beta} & \mathbb{D}Y \\ F\theta \downarrow & & & & \downarrow \varphi \\ FX & \xrightarrow{\quad \beta \quad} & Y & & \end{array}$$

If furthermore α_X is an isomorphism, then by Example 2.8 there is a canonical $\mathbb{D}$ -algebra structure on FX , and β is necessarily a map of $\mathbb{D}$ -algebras. In applications, where this reduction is unavailable, the best we can say is that for $f: X \rightarrow GY$ a map of $\mathbb{C}$ -algebras, Ff is a linking map. In the loop space story, we will take G to be the constant simplicial object functor $c_*: \mathcal{T} \rightarrow s\mathcal{T}$.

Let us continue on the story of an op-lax map (F, α) from $\mathbb{C}$ to $\mathbb{D}$ with a section (G, id) . Let (Y', φ') and (Y, φ) be $\mathbb{D}$ -algebras. We are now interested in maps $Y' \rightarrow Y$ that are not algebra maps, but still retain some monadic structure.

One might hope that the notion of linking that we have discussed provides the appropriate generalization. However, we saw in Remark 2.10, that the only linking maps between $\mathbb{D}$ -algebras are the $\mathbb{D}$ -algebra maps! It seems more appropriate then to look at maps *over* an algebra that is linked to both Y and Y' . In other words, those maps that factor as spans of linking maps: $Y' \leftarrow FX \rightarrow Y$.

Definition 2.11. Let $(F, \alpha): \mathbb{C} \rightarrow \mathbb{D}$ be an op-lax map of monads with section (G, id) . We say that a map $f: Y' \rightarrow Y$ is a *linked map* if there exist linking maps $\rho: FX \rightarrow Y'$ and $\beta: FX \rightarrow Y$ such that $\beta = f \circ \rho$.

$$\begin{array}{ccc} & FX & \\ \rho \swarrow & & \searrow \beta \\ Y' & \xrightarrow{f} & Y \end{array}$$

2.3. Simplicial Γ -algebras and sharpened homotopical monadicity. Our discussion so far on abstract linking theory is instantiated here to provide a sharpened version of the homotopical monadicity theorem. From the adjunction $(\Sigma^\infty, \Omega^\infty)$ we get an induced adjunction $(\Sigma_*^\infty, \Omega_*^\infty)$ between the categories of simplicial objects such that the simplicial unit η_* and counit ε_* are just levelwise applications of η and ε . The associated simplicial adjunction monad is given by Γ_* on $\mathcal{T}$, and consequently its product map μ_* is just a levelwise application of μ . As remarked earlier, by *simplicial Γ -algebras* we will mean Γ_* -algebras. Once more, by the usual yoga of adjunction monads, $\Omega_*^\infty E_*$ has a canonical simplicial Γ -algebra structure and therefore Ω_*^∞ really lands in the category $\Gamma_*[s\mathcal{T}]$. When we want to emphasize this fact, we use the notation $\Omega_{*,\Gamma}^\infty$ for the functor landing in $\Gamma_*[s\mathcal{T}]$. Throughout this section, we will also conflate γ with the natural map $\gamma: |\Omega_*^\infty \Sigma_*^\infty| \rightarrow \Omega^\infty |\Sigma_*^\infty| \cong \Omega^\infty \Sigma^\infty |-|$.

The following proposition outlines the setting in which we will apply our linking theory.

Proposition 2.12. $(|-|, \gamma)$ is an op-lax map of monads $\Gamma_* \rightarrow \Gamma$. That is to say, $|-|: s\mathcal{T} \rightarrow \mathcal{T}$ is equipped with a natural transformation $\gamma: |\Gamma_*| \rightarrow \Gamma|-|$ such that the following commute.

$$\begin{array}{ccc} |\Gamma_* \Gamma_*| & \xrightarrow{\gamma \Gamma_*} & \Gamma |\Gamma_*| \xrightarrow{\Gamma \gamma} \Gamma \Gamma|-| \\ \downarrow |\mu_*| & & \downarrow \mu|-| \\ |\Gamma_*| & \xrightarrow{\gamma} & \Gamma|-| \end{array} \quad \begin{array}{ccc} & |-| & \\ \eta_*| \swarrow & & \searrow \eta|-| \\ |\Gamma_*| & \xrightarrow{\gamma} & \Gamma|-| \end{array}$$

Moreover for every $E_* \in s\mathcal{S}$, the algebras $\Omega_{*,\Gamma}^\infty E_*$ and $\Omega_\Gamma^\infty |E_*|$ are $|-|$ -linked by γ , meaning the following diagram commutes.

$$\begin{array}{ccc} |\Gamma_* \Omega_*^\infty E_*| & \xrightarrow{\gamma \Omega_*^\infty} & \Gamma |\Omega_*^\infty E_*| \xrightarrow{\Gamma \gamma} \Gamma \Omega^\infty |E_*| \\ \downarrow |\Omega_*^\infty \varepsilon_*| & & \downarrow \Omega^\infty \varepsilon|-| \\ |\Omega_*^\infty E_*| & \xrightarrow{\gamma} & \Omega^\infty |E_*| \end{array}$$

Proof. We prove the commutativity of the unit diagram for $(|-|, \gamma)$ (the top right diagram), and the diagram for the linking of Ω_*^∞ to Ω^∞ —the remaining diagram

follows automatically. For the unit diagram, we first pass to the adjoint:

$$\begin{array}{ccc} & |\Sigma_*^\infty| & \\ |\Sigma_*^\infty \eta_*| \swarrow & & \searrow |\Sigma_*^\infty| \\ |\Sigma_*^\infty \Omega_*^\infty \Sigma_*^\infty| & \xrightarrow{|\varepsilon_* \Sigma_*^\infty|} & |\Sigma_*^\infty|. \end{array}$$

This diagram is just the realization of a triangle identity for the adjunction $(\Sigma_*^\infty, \Omega_*^\infty)$ in $s\mathcal{T}$, and therefore it commutes. As for the product diagram, passing to the adjoint again, we have the slightly larger diagram (where $\tilde{\gamma}: \Sigma^\infty |\Omega^\infty| \rightarrow |-|$ denotes the adjoint map to γ):

$$\begin{array}{ccccccc} \Sigma^\infty |\Gamma_* \Omega_*^\infty| & \xrightarrow{\cong} & |\Sigma_*^\infty \Omega_*^\infty \Sigma_*^\infty \Omega_*^\infty| & \xrightarrow{|\varepsilon_*|} & |\Sigma_*^\infty \Omega_*^\infty| & \xrightarrow{\cong} & \Sigma^\infty |\Omega_*^\infty| \xrightarrow{\Sigma^\infty \gamma} \Sigma^\infty \Omega^\infty |-| \\ \downarrow \Sigma^\infty |\Omega_*^\infty \varepsilon| & & \downarrow |\Sigma_*^\infty \Omega_*^\infty \Sigma_*^\infty| & & \downarrow |\varepsilon_*| & \swarrow \tilde{\gamma} & \searrow \tilde{\gamma} \\ \Sigma^\infty |\Omega_*^\infty| & \xrightarrow{\cong} & |\Sigma_*^\infty \Omega_*^\infty| & \xrightarrow{|\varepsilon_*|} & |-| & \xrightarrow{=} & |-| \end{array}$$

The left two squares commute trivially. The left triangle in the rightmost square commutes by definition of γ , and the right triangle commutes since γ and $\tilde{\gamma}$ are adjoint maps related by the counit ε . $\blacksquare$

As a consequence, we see that for each $Y \in \Gamma[\mathcal{T}]$, $\bar{Y}$ and $\Omega_{\Gamma}^\infty \mathbb{E}Y$ are linked by $\gamma: \bar{Y} \rightarrow \Omega_{\Gamma}^\infty \mathbb{E}Y$. Now we can make a remark on *what could have been*, using Example 2.8. If for some simplicial Γ -algebra X_* we had $\gamma: |\Gamma_* X_*| \cong |\Gamma X_*|$, then by Example 2.8 the realization $|X_*|$ would have a canonical Γ -algebra structure, and linking maps out of $|X_*|$ would just be maps of Γ -algebras. In particular, this is maybe what one would have hoped for $\Omega_{*,\Gamma}^\infty \mathbb{E}_* Y$ and $\bar{Y}$.

Proposition 2.13. *For every $X \in \mathcal{T}$, $c_* \Gamma X = \Gamma_* c_* X$. Consequently, the pair (c_*, id) defines a section of the op-lax map $(|-|, \gamma)$.*

Proof. The first statement and the fact that (c_*, id) provides an op-lax map $\Gamma \rightarrow \Gamma_*$ both follow from the levelwise definitions of Γ_* , μ_* , and η_* . By **SA1**, for every $X \in \mathcal{T}$, $|c_* X| \cong X$. It remains to note that $\gamma_{c_* X}: |\Gamma_* c_* X| = |c_* \Gamma X| \rightarrow \Gamma X$ is the identity map $\text{id}_{\Gamma X}$. $\gamma_{c_* X}$ was defined to be the adjoint to the map $|(\varepsilon_*)_{c_* \Sigma^\infty X}|$. By definition again, $|(\varepsilon_*)_{c_* \Sigma^\infty X}| = \varepsilon_{\Sigma^\infty X}$. However $\varepsilon_{\Sigma^\infty X}$ was defined to be the adjoint to $\text{id}_{\Gamma X}$, forcing $\gamma_{c_* X} = \text{id}_{\Gamma X}$. $\blacksquare$

Remark 2.10 has several important consequences in light of the previous proposition. First, as was already intuitively clear, if X is a Γ -algebra, then there is a canonical Γ_* -algebra structure on $c_* X$ given by $\Gamma_* c_* X = c_* \Gamma X \xrightarrow{c_* \theta} c_* X$. Moreover, linking maps between Γ -algebras, $X \rightarrow X'$ (seen as maps $|c_* X| \rightarrow X'$) are exactly the same as maps of Γ -algebras. Now suppose for $X_* \in \Gamma_*[\mathcal{T}]$ and $X' \in \Gamma[\mathcal{T}]$, we had a map of Γ_* -algebras $f_*: X_* \rightarrow c_* X'$. If it had been the case that $\gamma: |\Gamma_* X_*| \cong |\Gamma X_*|$, then $|f_*|$ would have been a Γ -algebra map. Nonetheless, Remark 2.10 shows that $|f_*|: |X_*| \rightarrow X'$ is always a linking map. In particular, $\zeta := |\zeta_*|$ is a linking map! In our homotopical setting, Definition 2.11 specializes to provide the notion of *linked weak equivalences*. These are the linked maps that are also weak equivalences.

What we have put together over this section and the last is the second main result of this paper, a sharpening of Theorem 0.1.

Theorem 0.2. *Let (Y, θ) be a Γ -algebra. There are maps*

$$Y \xleftarrow{\zeta} \bar{Y} \xrightarrow{\gamma} \Omega_{\Gamma}^{\infty} \mathbb{E}Y$$

where the following hold.

- (1) *Forgetting monadic structures, ζ is a homotopy equivalence and γ is a weak equivalence.*
- (2) *ζ and γ are maps that link $\bar{Y}$ to the Γ -algebras Y and $\Omega_{\Gamma}^{\infty} \mathbb{E}Y$ respectively.*
- (3) *For ν the homotopy inverse of ζ , the composite $\gamma \circ \nu$ is a linked weak equivalence between Y and $\Omega_{\Gamma}^{\infty} \mathbb{E}Y$.*

The consequence for the homotopy categories of Γ -algebras and connective spectra that one would hope for, is provided by a direct corollary of this sharpened result.

Corollary 0.3. *The full subcategory of Γ -algebras in $\mathcal{T}$ localized at the linked weak equivalences is equivalent to the homotopy category of connective spectra.*

3. VERIFICATION OF THE SIMPLICIAL AXIOMS

3.1. The easy axioms. In this section we prove the simplicial axioms **SA1-SA5** from Section 1. We have denoted by $s\mathcal{C}$ the category of simplicial objects in a category $\mathcal{C}$. In particular, $s\mathcal{T} = \text{ssSh}((\text{Sm}_S)_{\text{Nis}})$, the category of bisimplicial sheaves on Sm_S . In order to prescribe realization functors $|-|: s\mathcal{T} \rightarrow \mathcal{T}$ and $|-|: s\mathcal{S} \rightarrow \mathcal{S}$, we must first record our choice of the cosimplicial objects in $\mathcal{T}$ and $\mathcal{S}$. There is a conventional choice in the case of $\mathcal{T}$. For each $\mathbf{n} \in \Delta$, let Δ_n denote the object in $\mathcal{T}$ representing $S \times_{\text{Spec } \mathbb{Z}} \text{Spec } \mathbb{Z}[x_0, \dots, x_n] / (\sum_i x_i = 1)$. Given $f: \mathbf{n} \rightarrow \mathbf{m}$ in Δ , there is a map $\Delta_f: \Delta_n \rightarrow \Delta_m$ (considered as schemes) given by the ring homomorphism $\Delta_f(x_i) = \sum_{j \in f^{-1}(i)} x_j$ if $f^{-1}(i) \neq \emptyset$ and 0 otherwise. Together, this defines a cosimplicial object in $\mathcal{T}$ such that each Δ_n is isomorphic to $\mathbb{A}_S^n$. The typical prescription, noting that $\mathcal{T}$ is tensored over itself, then leads us to define $|K_*| = K_* \otimes_{\Delta} \Delta_*$ for $K_* \in s\mathcal{T}$. The realization functor here however takes a much simpler form due to the bisimplicial nature of simplicial spaces. For $K_* \in s\mathcal{T}$, thought of a bisimplicial sheaf $K_{*,*}$, we have that $|K_*|$ is isomorphic to the diagonal simplicial sheaf $d(K)_n = K_{n,n}$. By the usual yoga, since $\mathcal{T}$ is cotensored over itself, $|-|: s\mathcal{T} \rightarrow \mathcal{T}$ is a left adjoint functor with right adjoint $\mathbb{S}: \mathcal{T} \rightarrow s\mathcal{T}$ given by $\mathbb{S}(X)_q = F(\Delta_q, X)$ where $F(-, -)$ is the cotensor functor $\mathcal{T}^{\text{op}} \times \mathcal{T} \rightarrow \mathcal{T}$. Since $\mathcal{S}$ is tensored and cotensored over $\mathcal{T}$, our choice of the cosimplicial objects in $\mathcal{T}$ yields the realization functor $|-|: s\mathcal{S} \rightarrow \mathcal{S}$ given by $|E_*| = E_* \otimes_{\Delta} \Delta_*$ for simplicial spectra and its right adjoint (which we also denote $\mathbb{S}: \mathcal{S} \rightarrow s\mathcal{S}$) given by $\mathbb{S}(E)_q = F(\Delta_q, E)$. As in the case of spaces, F is used to denote the cotensor functor $\mathcal{T}^{\text{op}} \times \mathcal{S} \rightarrow \mathcal{S}$. This proves **SA1**.

The following is **SA2**, where the notion of a simplicial homotopy is that of [May67, Definitions 5.1].

Proposition 3.1. *If h_* is a homotopy between simplicial maps $f_*, g_*: K_* \rightarrow L_*$ in $s\mathcal{T}$, then $|h_*|$ determines a homotopy between maps $|f_*|, |g_*|: |K_*| \rightarrow |L_*|$ in $\mathcal{T}$. The corresponding statement for $\mathcal{S}$ also holds true.*

Proof. It is important to first note that $|-|$ preserves finite products since $\mathcal{T}$ (likewise $\mathcal{S}$) is cartesian closed and since $|-|$ preserves products of representables. The proof of this fact is purely formal, starting from the co-Yoneda lemma. By $\Delta[1]$, let us denote the standard simplicial set $\mathbf{n} \mapsto \text{Hom}_\Delta(\mathbf{n}, \mathbf{1})$, considered a simplicial space using the constant scheme construction of [SGA3V1, Exposé I, 1.8] levelwise. In the case of $\mathcal{T}$, by [May67, Proposition 6.2] there exists $H_*: \Delta[1] \times K_* \rightarrow L_*$ defined as follows. Working over $U \in \text{Sm}_S$, we let $x \in K_*(U)$ and define $H_*((0), x) = g_*(x)$, $H_*((1), x) = f_*(x)$ (here (0) denotes any simplex of $\partial_1 \Delta[1]$ and (1) any simplex of $\partial_0 \Delta[1]$), and $H_q(s_{q-1} \cdots s_{i-1} s_{i+1} \cdots s_0 \mathbf{1}, x) = \partial_{i+1} h_i(x)$ for each $0 \leq i \leq q-1$. H_* constructed in this way is not only a map of simplicial sets but also one of simplicial spaces since h_i and ∂_{i+1} are morphisms in $\mathcal{T}$. Finally, it remains to note that $|\Delta[1]|$ is isomorphic to $\mathbb{A}^1$, so that the composite $\mathbb{A}^1 \times |K_*| \xrightarrow{\sim} |\Delta[1]| \times |K_*| \xrightarrow{\sim} |\Delta[1]| \times K_* \xrightarrow{|H_*|} |L_*|$ provides the required homotopy. The story for spectra is similar.⁵ ■

Since Σ^∞ is a left adjoint functor, it commutes with the colimits that build $|-|: s\mathcal{T} \rightarrow \mathcal{T}$. Thanks to this observation, we obtain **SA4**.

Proposition 3.2. *There exists a natural isomorphism $\Sigma^\infty |K_*| \cong |\Sigma_*^\infty K_*|$ for each $K_* \in s\mathcal{T}$.*

We now proceed toward **SA3**. Recall that a simplicial object $X_* \in s\mathcal{V}$ for a category $\mathcal{V}$ is called *Reedy cofibrant* if the latching maps $LX_q = \text{colim}_{\phi: \mathbf{q}+1 \twoheadrightarrow \mathbf{s}, \phi \neq \text{id}} X_s \hookrightarrow X_{q+1}$ are cofibrations for each q . LX_q here is the scheme-theoretic analog to the union of all degenerate q -simplices in the simplicial set context. As in [May09] we implicitly assume that our spaces are nondegenerately based (i.e. that for each based space X , the basepoint inclusion $S \rightarrow X$ is a cofibration). So in this case every simplicial space is Reedy cofibrant. **SA3** is derived from a general model-theoretic fact first due to [Ree73]. The presentation here is inspired from [May74].

Proposition 3.3. *$|-|$ preserves weak equivalences between Reedy cofibrant objects in $s\mathcal{T}$ (resp. in $s\mathcal{S}$).*

We define weak equivalences in $s\mathcal{T}$ and $s\mathcal{S}$ as those maps that are level-wise weak equivalences (in the appropriate model category)⁶. The proof of the above theorem relies on the motivic analog of Brown's gluing theorem [Bro06, Theorem 7.5.7].

⁵The proof here is almost identical to that of [May72, Corollary 11.10], and might be formalized to a more general context—for example to a cartesian closed concrete category with a notion of homotopy.

⁶In a sense, this result records the extent to which $|-|$ induces an equivalence between the homotopy categories of $s\mathcal{T}$ and $\mathcal{T}$ (resp. $s\mathcal{S}$ and $\mathcal{S}$).

Theorem 3.4. *Suppose given a commutative diagram in $\mathcal{T}$:*

$$\begin{array}{ccccc}
 & & X & \xrightarrow{g} & Z \\
 & \nearrow \alpha & \downarrow j & \nearrow \beta & \downarrow \bar{j} \\
 A & \xrightarrow{f} & C & & \\
 \downarrow i & & \downarrow \bar{i} & & \downarrow \\
 & \nearrow \gamma & Y & \xrightarrow{\bar{g}} & W \\
 B & \xrightarrow{\bar{f}} & D & \nearrow \delta &
 \end{array}$$

that i and j are cofibrations, and that the front and back squares are pushouts. If α , β , and γ are homotopy equivalences, then so is δ .

Proof of Proposition 3.3. As before, we provide the proof for spaces—the proof for spectra is nearly identical and is hence omitted. Let $f_*: X_* \rightarrow X'_*$ be a map of Reedy cofibrant spaces. We have the following pushout square for each q :

$$\begin{array}{ccc}
 (LX_q \times \Delta_{q+1}) \sqcup_{LX_q \times \partial \Delta_{q+1}} (X_{q+1} \times \Delta_{q+1}) & \xrightarrow{g} & F_q|X_*| \\
 \downarrow & & \downarrow \\
 X_{q+1} \times \partial \Delta_{q+1} & \longrightarrow & F_{q+1}|X_*|.
 \end{array}$$

Here the standard map $g(s_i x, u) = [x, \sigma_i u]$ and $g(x, \delta_i v) = [\partial_i x, v]$ is used. The prescription of the map here is interpreted to be given at an arbitrary object in the site $U \in \text{Sm}_S$. $F_q|X_*|$ denotes the q th piece of the natural filtration on $|X_*|$, given by capping off the tensor product defining $|X_*|$ at the q -simplices. Alternatively, one might take the above to be the definition of $F_q|X_*|$ inductively (with $F_0|X_*| = X_0$) so that $|X_*| = \varinjlim F_q|X_*|$. There is also a similar square for X'_* of course. Proceeding inductively and applying Theorem 3.4, we see it suffices to show that $f_q: LX_q \rightarrow LX'_q$ is a homotopy equivalence for each q . For each $0 \leq k \leq q$ let $L^k X_q = \text{colim}_{\phi: \mathbf{q}+1 \rightarrow \mathbf{s}, s \leq k} X_s$. We obtain the following commutative diagram with a pushout square on the right side:

$$\begin{array}{ccccc}
 L^{k-1}X_{q-1} & \xrightarrow[\sim]{L_k} & L^{k-1}X_q \times_{X_{q+1}} & L_k X_q & \longrightarrow & L^{k-1}X_q \\
 \downarrow & & \downarrow & & & \downarrow \\
 X_q & \xrightarrow[\sim]{L_k} & L_k X_q & \longrightarrow & & L^k X_q
 \end{array}$$

Here $L_k X_q$ denotes the image of $X_k \hookrightarrow X_{q+1}$, the map seen in the colimit definition of LX_q . Proceeding by induction on q , assuming that each $L^{k-1}X_{q-1} \rightarrow L^k X_{q-1}$ is a cofibration for each $0 < k < q$, we conclude that $L^{k-1}X_q \rightarrow L^k X_q$ is a cofibration. The same is true of X' . Since we know $L_0: X_q \rightarrow L_0 X_q$ is an isomorphism, $f_{q+1}: L^0 X_q \rightarrow L^0 X'_q$ must be a homotopy equivalence. By induction on q , and induction on k having fixed q , and Theorem 3.4, we know that $f_{q+1}: L^k X_q \rightarrow L^k X'_q$ is a homotopy equivalence for all k and q . $\blacksquare$

SA5 is clear from Proposition 3.2, our definition of connective spectra, and the fact that $|-|$ is a left adjoint. Alternatively, if one defines homotopy sheaves of

spectra as being certain direct limits of homotopy sheaves of spaces, then since $|-|$ is a left adjoint, it commutes with those direct limits. The statement is clear once again if one takes connective spectra to be spectra with certain vanishing homotopy sheaves.

Proposition 3.5. $|-|: s\mathcal{T} \longrightarrow \mathcal{T}$ takes levelwise connective spectra to connective spectra.

3.2. The hard axiom. We conclude with the proof of **SA6**, the most significant assumption in this paper.

Proposition 3.6. The map $\gamma: |\Omega_*^\infty E_*| \longrightarrow \Omega^\infty |E_*|$ given by the adjoint to the composite $\Sigma^\infty |\Omega_*^\infty E_*| \cong |\Sigma_*^\infty \Omega_*^\infty E_*| \xrightarrow{|\varepsilon|} |E_*|$ is, for levelwise connective E_* , a weak equivalence in $\mathcal{T}$.

An important distinction from the theory of operadic monads must be made here. Generally speaking, a monad $\mathbb{C}$ coming from an operad commutes with $|-|$ up to isomorphism by the categorical Fubini theorem. For the adjunction monad however, as we will show now, a priori Ω^∞ commutes with $|-|$ only up to weak equivalence—and therefore Γ commutes with $|-|$ only up to weak equivalence.

We prove Proposition 3.6 by reducing to n -fold and singlefold loop spaces first, and then passing to the colimit. For $X \in \mathcal{T}$ denote by ΩX the hom-space $\underline{\mathrm{Hom}}_{\mathcal{T}}(S^{\mathbb{A}^1}, X)$ (where $S^{\mathbb{A}^1} = \mathbb{A}^1/(\mathbb{A}^1 - \{0\})$). Denote by $\Omega^n X$ the hom-space $\underline{\mathrm{Hom}}_{\mathcal{T}}(S^{\mathbb{A}^n}, X)$ (where $S^{\mathbb{A}^n} = \mathbb{A}^n/(\mathbb{A}^n - \{0\})$). Ω^n (as seen by the loop-suspension adjunction) is the n -fold composition of Ω with itself. There is a map $|\Omega_*^n X_*| \longrightarrow \Omega^n |X_*|$ again given by the adjoint to $\Sigma^n |\Omega_*^n X_*| \cong |\Sigma_*^n \Omega_*^n X_*| \xrightarrow{|\varepsilon|} |X_*|$ (here $\Sigma^n(-) = - \wedge S^{\mathbb{A}^n}$) which we also call γ .

Lemma 3.7. If $X_* \in s\mathcal{T}$ is Reedy cofibrant and each X_q is connected, then $\gamma: |\Omega_* X_*| \longrightarrow \Omega |X_*|$ is a weak equivalence in $\mathcal{T}$.

We first note that Lemma 3.7 allows us to conclude Proposition 3.6. Of course, iteratively applying the above lemma tells us that $\gamma^n: |\Omega_*^n X_*| \longrightarrow \Omega^n |X_*|$ is a weak equivalence for Reedy cofibrant X_* with connected X_q . The idea now is that (under the assumptions on E_*) the map $\gamma: |\Omega_*^\infty E_*| \longrightarrow \Omega^\infty |E_*|$ can be identified with the colimit of the maps $\gamma^n: |\Omega_*^n (E_n)_*| \longrightarrow \Omega^n |(E_n)_*|$, so that Proposition 3.6 follows by passing to colimits. We must clarify the notation used. We have written E_n to mean $E_{\mathcal{U}/\mathbb{A}^n}$ —the *sequential* coordinatized indexing here is chosen purely for simplicity. Note that each $(E_n)_*$ is a simplicial space, and $\Omega_*^n (E_n)_*$ is compatibly isomorphic to $(E_0)_*$, so that the colimit of $|\Omega_*^n (E_n)_*|$ is identified with $|\Omega_*^\infty E_*|$. By a remark made after Proposition 1.10, for $E_* = LT_*$, we can compute the colimit of $\Omega^n |(E_n)_*|$ as $\Omega^\infty |LT_*| \cong \Omega^\infty L|T_*|$, where the realization of a simplicial prespectrum is defined as the levelwise realization—so that the needed observation follows by working at the prespectrum level.

There is, however, a caveat. The argument we use moving forward only works on fibrant objects—this requires us to pass to fibrant replacements. We introduce fibrant replacements in $s\mathcal{T}$ after proving Lemma 3.9 thus proving Lemma 3.7. The following is a precise reformulation of the claim that emphasizes what is needed.

Lemma 3.8. *Let $X_* \in s\mathcal{T}$, then $|P_*X_*|$ is contractible, and there are natural morphisms γ and δ making the following diagram commute:*

$$\begin{array}{ccccc} |\Omega_*X_*| & \xrightarrow{\subset} & |P_*X_*| & \xrightarrow{|p_*|} & |X_*| \\ \gamma \downarrow & & \delta \downarrow & & \parallel \\ \Omega|X_*| & \xrightarrow{\subset} & P|X_*| & \xrightarrow{p} & |X_*|. \end{array}$$

Moreover, when X_* is fibrant, Reedy cofibrant, and level-wise connected, the top row of the above diagram $|\Omega_*X_*| \longrightarrow |P_*X_*| \xrightarrow{|p_*|} |X_*|$ is a homotopy fiber sequence, and therefore $\gamma: |\Omega_*X_*| \longrightarrow \Omega|X_*|$ is a weak equivalence.

We have used the standard notation PY to denote the path-space of $Y \in \mathcal{T}$, defined by $\underline{\text{Hom}}_{\mathcal{T}}(\mathbb{A}_+^1, Y)$. The simplicial version P_*Y_* for $Y_* \in s\mathcal{T}$ is defined by taking the levelwise path-space. We outline the proof of the first part here. Firstly, we note that there is a natural contracting homotopy equivalence $h: PY \longrightarrow *$ given by $\varphi \mapsto \varphi \circ \iota_0$ (here $\iota_0: S \longrightarrow \mathbb{A}_S^1$ denotes the inclusion of S as the basepoint 0 of $\mathbb{A}_+^1$ and we identify the composite $S \longrightarrow Y$ with the basepoint of Y). This homotopy, applied to each PX_q , descends to a simplicial contracting homotopy $\Delta[1] \times P_*X_* \longrightarrow P_*X_*$. By Proposition 3.1, we conclude that $|P_*X_*|$ is indeed contractible. We are left with defining $\delta: |P_*X_*| \longrightarrow P|X_*|$. Everything that follows in this paragraph fixes an arbitrary $U \in \text{Sm}_S$, and we work over this U implicitly. For $f \in PX_q$, $u \in \Delta_q$, and $t \in \mathbb{A}^1$, define $\delta[f, u](t) = [f(t), u]$. δ is a well-defined morphism in $\mathcal{T}$, and it indeed makes the diagram in question commute. In some sense, this definition of δ was the only sensible choice we had. There is categorical reason for this. The functors $-\wedge \mathbb{A}_+^1$ and $P(-)$ are adjoint to each other. The map $\delta: |P_*X_*| \longrightarrow P|X_*|$ is the adjoint to $|P_*X_*| \wedge \mathbb{A}_+^1 \cong |(PX \wedge \mathbb{A}_+^1)_*| \xrightarrow{|\epsilon_*|} |X_*|$ where $\epsilon: PX \wedge \mathbb{A}_+^1 \longrightarrow X$ is the counit map.

What remains is the following, an analog of J.P. May's result in [May72], and a specialization of Anderson's result in [And78], for the geometric realization of fibrations.

Lemma 3.9. *If $p_*: E_* \longrightarrow B_*$ is a fibration in $s\mathcal{T}$ with fiber F_* and B_* is Reedy cofibrant and level-wise connected then $|F_*| \longrightarrow |E_*| \xrightarrow{|p_*|} |B_*|$ is a homotopy fiber sequence. Moreover, when X_* is fibrant, $p_*: P_*X_* \longrightarrow X_*$ is a fibration in $s\mathcal{T}$ with fiber Ω_*X_* , leading us to conclude that $|\Omega_*X_*| \longrightarrow |P_*X_*| \xrightarrow{|p_*|} |X_*|$ is a homotopy fiber sequence (when X_* is Reedy cofibrant and levelwise connected).*

We must first define what we mean by a fibration $p_*: E_* \longrightarrow B_*$ in $s\mathcal{T}$. This definition follows that of Reedy for the model structure on $s\mathcal{C}$ given a model category $\mathcal{C}$, using coskeletons. Namely a map $p_*: E_* \longrightarrow B_*$ is a fibration in $s\mathcal{T}$ if for each m , the induced map $E_{m+1} \longrightarrow \text{cosk}_m(E)_{m+1} \times_{\text{cosk}_m(B)_{m+1}} B_{m+1}$ is a fibration in $\mathcal{T}$. Together with the dually defined cofibrations in $s\mathcal{T}$, we obtain a model structure on $s\mathcal{T}$ that is in fact simplicially enriched. There is a pairing $-\otimes -: \mathbf{sSet} \times s\mathcal{T} \longrightarrow s\mathcal{T}$ given by $(A_* \otimes X_*)_k = X_k[A_k] := \coprod_{a \in A_k} X_k$. Appealing to the dual provides a pairing $\text{Hom}(-, -): \mathbf{sSet}^{\text{op}} \times s\mathcal{T} \longrightarrow s\mathcal{T}$. Finally, there is the standard $\mathbf{sSet}$ -enrichment on $s\mathcal{T}$, $\text{Hom}(-, -): s\mathcal{T}^{\text{op}} \times s\mathcal{T} \longrightarrow \mathbf{sSet}$, given by $\text{Hom}(X_*, Y_*) =$

$\mathrm{Hom}_{s\mathcal{T}}(\Delta[-] \otimes X_*, Y_*) \in \mathbf{sSet}$. Moreover, these pairings satisfy the pullback-power axiom⁷:

- If $i_*: A_* \rightarrow B_*$ is a cofibration in $\mathbf{sSet}$ and $p_*: X_* \rightarrow Y_*$ is a fibration in $s\mathcal{T}$, then the following is a fibration in $\mathbf{sSet}$.

$$\mathrm{Hom}(B_*, X_*) \rightarrow \mathrm{Hom}(B_*, Y_*) \times_{\mathrm{Hom}(A_*, Y_*)} \mathrm{Hom}(A_*, X_*)$$

Moreover, the above map is an acyclic fibration if p_* is also a weak equivalence.

With this new enriched model structure in our toolbox, we now handle the simplicial path space fibration on fibrant objects. That $p_*: P_*X_* \rightarrow X_*$ is a fibration in $s\mathcal{T}$ when X_* is fibrant, is an instance of a well-known general fact (see [GJ99, Lemma 7.5] for example). The idea is to begin with the cofiber sequence of pointed simplicial sets⁸ $\partial\Delta[1] \hookrightarrow \Delta[1] \rightarrow \Delta[1]/\partial\Delta[1]$. The pullback-power axiom for $s\mathcal{T}$ discussed above then ensures that when we power this sequence by a fibrant X_* , we obtain a fiber sequence $\mathrm{Hom}(\Delta[1]/\partial\Delta[1], X_*) \rightarrow \mathrm{Hom}(\Delta[1], X_*) \rightarrow \mathrm{Hom}(\partial\Delta[1], X_*)$. Noting that $\Omega_*X_* \simeq \mathrm{Hom}(\Delta[1]/\partial\Delta[1], X_*)$, $P_*X_* \simeq \mathrm{Hom}(\Delta[1], X_*)$, and $\mathrm{Hom}(\partial\Delta[1], X_*) \simeq X_*$, this yields a fiber sequence $\Omega X_* \rightarrow P X_* \rightarrow X_*$.

It remains to show that if $p_*: E_* \rightarrow B_*$ is a fibration in $s\mathcal{T}$ with fiber F_* and B_* is Reedy cofibrant and level-wise connected, then $|F_*| \rightarrow |E_*| \rightarrow |B_*|$ is a homotopy fiber sequence. This is the real meat of Lemma 3.9. In [May72], the author employs the Dold-Thom criteria for quasifibrations to achieve the required analogous result. Anderson, in [And78], provides a conceptually clearer treatment of the result by passing to bisimplicial sets and solving a lifting problem with respect to the image of the left adjoint of the realization functor. The initial idea here was to make use of Anderson's argument, or the π_* -Kan condition for bisimplicial sets (due to Bousfield and Friedlander [BF78])—both of which, in a sense, rely on the construction of the simplicial groupoid $\Pi_\infty(-)$ ⁹. However, the conceptual reformulation, and the strengthened result pertaining to geometric realizations of homotopy pullbacks, in [Rez14] is more amenable to our purposes—and is easier to approach. A specialized application of [Lur17, Proposition 5.5.6.17] to the ∞ -topos $L_W\mathcal{T}$ also yields the required result, but we expect that the generality of Lurie's statement will be more relevant in future studies where Rezk's argument is not readily accessible. We state the goal now:

Theorem 3.10. *Let $B_* \in s\mathcal{T}$ be a simplicial space such that the simplicial sheaf $\pi_0^{\mathbb{A}^1}(B_*)$, defined levelwise as $\pi_0^{\mathbb{A}^1}(B_*)_n = \pi_0^{\mathbb{A}^1}(B_n)$, is discrete (as a simplicial object). Let $f_*: Y_* \rightarrow B_*$ be a map in $s\mathcal{T}$. Then for every homotopy pullback square,*

$$\begin{array}{ccc} F_* & \longrightarrow & Y_* \\ \downarrow & & \downarrow f_* \\ E_* & \xrightarrow{p_*} & B_* \end{array}$$

⁷They also satisfy an equivalent dual *pushout-copower* axiom.

⁸We consider $\Delta[1]$ to be pointed at 1 here.

⁹For completeness, the groupoid $\Pi_\infty(Z)$ is the wreath product of the fundamental groupoid with the product of the higher homotopy groups of Z .

the following square,

$$\begin{array}{ccc} |F_*| & \longrightarrow & |Y_*| \\ \downarrow & & \downarrow |f_*| \\ |E_*| & \xrightarrow{|p_*|} & |B_*| \end{array}$$

obtained by applying $|-|$ to each corner, is still a homotopy pullback.

3.3. Realization-fibrations and the proof of Theorem 3.10. The notation used in Theorem 3.10 is of course suggestive of the fact that in our applications, we will take Y_* to be the terminal simplicial space—so that the level-wise connectedness of B_* will give us Lemma 3.9. But one must take brief respite to make a conceptual remark about Theorem 3.10 to put us directly in the framework of [Rez14]. Implicitly restricting down to Reedy cofibrant objects, a routine check using the Bousfield-Kan map tells us that there is a weak equivalence $\mathrm{hocolim}_{\Delta^{\mathrm{op}}} X_* \longrightarrow |X_*|$ for each $X_* \in s\mathcal{T}$. Under this lens, Theorem 3.10 is manifestly a question of the preservation of homotopy pullbacks under certain kinds of homotopy colimits. For the rest of this section, we will use $|X_*|$ and $\mathrm{hocolim}_{\Delta^{\mathrm{op}}} X_*$ interchangeably (and by one, we may as well mean the other). By a *realization-fibration* (sometimes denoted RF), we will mean a map $f_*: Y_* \longrightarrow B_*$ such that the output of Theorem 3.10 holds true (i.e. a realization-fibration is a map such that homotopy colimits commute with homotopy pullbacks of the map). Our task then is to ascertain conditions under which a map of simplicial spaces is a realization-fibration.

The perspective due to Rezk allows us to deduce Theorem 3.10 from a local-to-global principle for realization-fibrations, applied to the homotopy colimit of representable sifted presheaves. Our starting point for the local-to-global point of view is the principle of descent, as presented in [Rez10, Section 6.5 and Proposition 6.6] for model toposes. We say that a natural transformation $f: U \longrightarrow V$ of functors $U, V: \mathcal{J} \longrightarrow \mathcal{T}$ is equifibered if for every $J \longrightarrow J'$ in $\mathcal{J}$, the associated naturality square is a homotopy pullback square. We will use the notation $(\mathcal{C}, \mathcal{D})$ to mean the functor category whose objects are functors $\mathcal{C} \longrightarrow \mathcal{D}$ and morphisms are natural transformations.

Lemma 3.11 (Descent). *Let $\mathcal{J}$ be a (small) category.*

- (1) *Let $p: E \longrightarrow B$ be a fibration in $\mathcal{T}$ and $V: \mathcal{J} \longrightarrow \mathcal{T}$ be a diagram of spaces such that $h: \mathrm{hocolim}_{\mathcal{J}} V \longrightarrow B$ is a weak equivalence in $\mathcal{T}$. Let $U: \mathcal{J} \longrightarrow \mathcal{T}$ be the diagram of spaces defined by $U(J) = V(J) \times_B E$. The map $\mathrm{hocolim}_{\mathcal{J}} U \longrightarrow E$ is a weak equivalence.*
- (2) *Let U and V be diagrams of spaces $\mathcal{J} \longrightarrow \mathcal{T}$, and $f: U \longrightarrow V$ an equifibered map in $(\mathcal{J}, \mathcal{T})$. For each $J \in \mathcal{J}$, the square:*

$$\begin{array}{ccc} U(J) & \longrightarrow & \mathrm{hocolim}_{\mathcal{J}} U \\ \downarrow & & \downarrow \\ V(J) & \longrightarrow & \mathrm{hocolim}_{\mathcal{J}} V \end{array}$$

is a homotopy pullback.

(3) Let $f: X \rightarrow Y$ be an equifibered map, and

$$\begin{array}{ccc} X' & \longrightarrow & X \\ \downarrow & & \downarrow f \\ Y' & \longrightarrow & Y \end{array}$$

a homotopy pullback square in $(\mathcal{J}, \mathcal{T})$. Applying $\mathrm{hocolim}_{\mathcal{J}}$ at each corner of the above square still yields a homotopy pullback.

The central local-to-global statement we seek states that under certain hypotheses (i.e. equifiberedness) realization-fibrations can be glued together to get a realization-fibration.

Lemma 3.12. *Let $\mathcal{J}$ be a small category, $h: W \rightarrow V$ an equifibered map in $(\mathcal{J}, s\mathcal{T})$ that is an object-wise realization-fibration. Then the map obtained by passing to the homotopy colimit $\mathrm{hocolim}_{\mathcal{J}} h: \mathrm{hocolim}_{\mathcal{J}} W \rightarrow \mathrm{hocolim}_{\mathcal{J}} V$ is also a realization-fibration.*

Proof. The argument in this proof is due to [Rez14]. Define $B_* := \mathrm{hocolim}_{\mathcal{J}} V$ and pick a factorization of $\mathrm{hocolim}_{\mathcal{J}} h: \mathrm{hocolim}_{\mathcal{J}} W \rightarrow B_*$ as the composite $\mathrm{hocolim}_{\mathcal{J}} W \xrightarrow{\iota_*} E_* \xrightarrow{p_*} B_*$, where ι_* is a weak equivalence and p_* is a fibration. Note that to show $\mathrm{hocolim}_{\mathcal{J}} h$ is an RF, it suffices to show p_* is an RF. Toward that end, we make some initial remarks. For each $J \in \mathcal{J}$, we have a map $V(J)_* \rightarrow B_*$. Denote by $U(J)_*$ the fiber product $E_* \times_{B_*} V(J)_*$.

$$\begin{array}{ccc} U(J)_* & \longrightarrow & E_* \\ \downarrow & & \downarrow p \\ V(J)_* & \longrightarrow & B_* \end{array}$$

$h: W \rightarrow V$ is equifibered, so Lemma 3.11 tells us that there is a homotopy pullback square.

$$\begin{array}{ccc} W(J)_* & \longrightarrow & \mathrm{hocolim}_{\mathcal{J}} W \\ \downarrow & & \downarrow \mathrm{hocolim}_{\mathcal{J}} h \\ V(J)_* & \longrightarrow & B \end{array}$$

Since $\iota_*: \mathrm{hocolim}_{\mathcal{J}} W \rightarrow E_*$ is a weak equivalence, we see that there is a weak equivalence $W(J)_* \rightarrow U(J)_*$ for each $J \in \mathcal{J}$. In particular, the maps $U(J)_* \rightarrow V(J)_*$ are RF too. Lemma 3.11 also provides a chain of weak equivalences $\mathrm{hocolim}_{\mathcal{J}} W \rightarrow \mathrm{hocolim}_{\mathcal{J}} U \rightarrow E_*$. We are ready to show that p_* is an RF now.

Let $\phi: B'_* \rightarrow B_*$ be a map in $s\mathcal{T}$. Define E'_* , U' , and V' by the following homotopy pullbacks.

$$\begin{array}{ccc} E'_* & \longrightarrow & E_* \\ \downarrow & & \downarrow p \\ B'_* & \xrightarrow{\phi} & B_* \end{array} \quad \begin{array}{ccc} V'(J)_* & \longrightarrow & V(J)_* \\ \downarrow & & \downarrow \\ B'_* & \xrightarrow{\phi} & B_* \end{array} \quad \begin{array}{ccc} U'(J)_* & \longrightarrow & U(J)_* \\ \downarrow & & \downarrow \\ B'_* & \xrightarrow{\phi} & B_* \end{array}$$

By observing the middle and the right homotopy pullbacks, and using the homotopy pullback lemma, we get the following homotopy pullback square.

$$\begin{array}{ccc} U' & \longrightarrow & U \\ \downarrow & & \downarrow \\ V' & \longrightarrow & V \end{array}$$

We already know that there is a weak equivalence $\mathrm{hocolim}_{\mathcal{J}} U \rightarrow E_*$, and an equality $B_* = \mathrm{hocolim}_{\mathcal{J}} V$. But by Lemma 3.11, there are also weak equivalences $\mathrm{hocolim}_{\mathcal{J}} V' \rightarrow B'_*$ and $\mathrm{hocolim}_{\mathcal{J}} U' \rightarrow E'_*$. This tells us that applying $\mathrm{hocolim}_{\mathcal{J}}$ to each corner of the above square, we get a homotopy pullback

$$\begin{array}{ccc} E'_* & \longrightarrow & E_* \\ \downarrow & & \downarrow p \\ B'_* & \xrightarrow{\phi} & B_* \end{array}$$

It suffices to know that applying $|-|$ at each corner, we still get a homotopy pullback square. But notice that $|\mathrm{hocolim}_{\mathcal{J}}(-)| \simeq \mathrm{hocolim}_{\mathcal{J}} |-|$ since colimits commute with colimits. So we might as well have realized down in our earlier homotopy pullback, and then applied $\mathrm{hocolim}_{\mathcal{J}}$. Note that since each $U(J)_* \rightarrow V(J)_*$ is an RF, upon realization we get a homotopy pullback.

$$\begin{array}{ccc} |U'| & \longrightarrow & |U| \\ \downarrow & & \downarrow \\ |V'| & \longrightarrow & |V| \end{array}$$

Now we notice that $|U| \rightarrow |V|$ is equifibered since each $U(J)_* \rightarrow V(J)_*$ is an RF. Therefore by descent, applying $\mathrm{hocolim}_{\mathcal{J}}$ to each corner will yield yet another homotopy pullback square, which is what we needed. $\blacksquare$

We now make use of the siftedness of Δ^{op} . When we say sifted, we implicitly mean the kind of homotopy siftedness (after a routine check in homotopy coend calculus using the Bousfield-Kan formula) that results in a canonical weak equivalence $|X_* \times Y_*| \rightarrow |X_*| \times |Y_*|$ for any X_*, Y_* in $s\mathcal{T}$. In simple terms, siftedness tells us that realization commutes with finite products in $s\mathcal{T}$. The primary utility of working over a sifted category is demonstrated in the following lemma.

Lemma 3.13. *All maps of the form $p_*: B_* \times C_* \rightarrow B_*$ in $s\mathcal{T}$ are realization-fibrations.*

Proof. Let $\phi_*: B'_* \rightarrow B_*$ be a map in $s\mathcal{T}$. Any homotopy pullback of p_* along ϕ_* must look like the following diagram.

$$\begin{array}{ccc} B'_* \times C_* & \xrightarrow{\phi_* \times \mathrm{id}} & B_* \times C_* \\ \pi_{B'_*} \downarrow & & \downarrow p_* \\ B'_* & \xrightarrow{\phi_*} & B_* \end{array}$$

where $\pi_{B'_*}$ is the projection map onto B'_* . Applying $|-|$ on each corner, and using the canonical weak equivalences $|B_* \times C_*| \xrightarrow{\sim} |B_*| \times |C_*|$ and $|B'_* \times C_*| \xrightarrow{\sim} |B'_*| \times |C_*|$, we get a diagram.

$$\begin{array}{ccc} |B'_*| \times |C_*| & \xrightarrow{|\phi_*| \times \text{id}} & |B_*| \times |C_*| \\ \pi_{|B'_*|} \downarrow & & \downarrow |p_*| \\ |B'_*| & \xrightarrow{|\phi_*|} & |B_*| \end{array}$$

where $\pi_{|B'_*|}$ is the projection map onto $|B'_*|$. This is evidently another homotopy pullback square, and we are done. $\blacksquare$

Now we introduce the notions of weak and local projection maps. We will soon see that every local projection map is in fact a realization-fibration, and that the hypotheses in Theorem 3.10 ensure that $f_*: Y_* \rightarrow B_*$ will be a local projection. A map $p_*: E_* \rightarrow B_*$ in $s\mathcal{T}$ is called a *weak projection* if it is weakly equivalent over B_* to a projection map $\tilde{p}_*: B_* \times C_* \rightarrow B_*$ in $s\mathcal{T}$. We will say that a map $p_*: E_* \rightarrow B_*$ in $s\mathcal{T}$ is a *local projection* if for every n , and every homotopy pullback square of the form:

$$\begin{array}{ccc} E'_* & \longrightarrow & E_* \\ q_* \downarrow & & \downarrow p_* \\ \Delta[n] & \xrightarrow{b} & B_* \end{array}$$

where $\Delta[n]$ denotes the standard simplicial set $\mathbf{m} \mapsto \text{Hom}_\Delta(\mathbf{m}, \mathbf{n})$ (thought of as a simplicial space by the constant scheme construction), the map q_* is a weak projection map. Given $X_* \in s\mathcal{T}$, there is a simplicial presheaf $\pi_0^{\mathbb{A}^1}(X_*)$ given by $(\pi_0^{\mathbb{A}^1}(X_*))_n = \pi_0^{\mathbb{A}^1}(X_n)$. We can think of $\pi_0^{\mathbb{A}^1}(X_*)$ as a levelwise discrete object in $s\mathcal{T}$. Given a map $p_*: E_* \rightarrow B_*$, let $\text{lproj}(p)_n \subseteq (\pi_0^{\mathbb{A}^1}(B_*))_n$ denote the presheaf prescribed locally as the collection of sections of $(\pi_0^{\mathbb{A}^1}(B_*))_n$ represented by maps $b_*: \Delta[n] \rightarrow B_*$ such that the pullback of p_* along b_* is a weak projection. This prescription defines a simplicial subpresheaf $\text{lproj}(p_*) \subseteq \pi_0^{\mathbb{A}^1}(B_*)$.

Suppose $p_*: E_* \rightarrow B_*$ and $\phi_*: B'_* \rightarrow B_*$ are maps in $s\mathcal{T}$. Note that by the pullback lemma, the pullback of p_* along ϕ_* is a local projection iff $\phi(\pi_0^{\mathbb{A}^1} B'_*) \subseteq \text{lproj}(p_*)$. Taking $B'_* = B_*$ and $\phi_* = \text{id}_{B_*}$, this means in particular that p_* is a local projection iff $\text{lproj}(p_*) = \pi_0^{\mathbb{A}^1}(B_*)$. In applications, we typically demonstrate that p_* is a local projection by showing that $\pi_0^{\mathbb{A}^1}(B_*) \subseteq \text{lproj}(p_*)$.

Lemma 3.14. *All local projections in $s\mathcal{T}$ are realization-fibrations.*

Proof. Let $p_*: E_* \rightarrow B_*$ be a local projection in $s\mathcal{T}$. Note that we can write B_* up to weak equivalence as the colimit of representable objects in $s\mathcal{T}$ (by which we mean $\Delta[n]$ for $n \geq 0$, considered as simplicial spaces). For each map $\Delta[n] \rightarrow B_*$ forming the appropriate colimit, since p_* is a local projection, we can form the

homotopy pullback square below.

$$\begin{array}{ccc} \Delta[n] \times C_* & \longrightarrow & E_* \\ \downarrow & & \downarrow p \\ \Delta[n] & \longrightarrow & B_* \end{array}$$

Descent tells us that E_* then is the homotopy colimit of objects of the form $\Delta[n] \times C_*$. In sum, p_* can be written as the homotopy colimit of maps that each look like $\Delta[n] \times C_* \longrightarrow \Delta[n]$. By Lemma 3.13, we know that each of these maps is a realization-fibration. Lemma 3.12 then tells us that p_* is a realization-fibration too. $\blacksquare$

Proof of Theorem 3.10. We return to the notation set up in the statement of Theorem 3.10. Namely, let $f_*: Y_* \longrightarrow B_*$ be a map in $s\mathcal{T}$. Note first that since $\Delta[0]$, as a simplicial space, is terminal, every map into $\Delta[0]$ in $s\mathcal{T}$ is a weak projection. As a result, any 0-simplex of $\pi_0^{\mathbb{A}^1}(B_*)$, thought of as being represented by a map out of $\Delta[0]$, is contained in $\text{lproj}(f_*)$. Since $\pi_0^{\mathbb{A}^1}(B_*)$ is discrete by assumption, meaning $\pi_0^{\mathbb{A}^1}(B_*) \simeq \text{const}(\pi_0^{\mathbb{A}^1}(B_0))$, we see in fact that $\pi_0^{\mathbb{A}^1}(B_*) \subseteq \text{lproj}(f_*)$. We conclude that f_* is a local projection, and therefore a realization-fibration. $\blacksquare$

We complete the proof of Lemma 3.7 by introducing, and passing to, fibrant replacements. Recall that $\mathcal{T}$ admits functorial factorization. It follows from a result on Reedy model structures that this property lifts to simplicial spaces as well, so that $s\mathcal{T}$ admits functorial factorization too (see [Hov99, Theorem 5.2.5] for the general statement due to transfinite induction). The existence of a fibrant replacement functor $\text{Ex}: s\mathcal{T} \longrightarrow s\mathcal{T}$ with the following properties is now immediate.

- (1) For every $X_* \in s\mathcal{T}$, $\text{Ex}(X_*)$ is fibrant in $s\mathcal{T}$.
- (2) For every $X_* \in s\mathcal{T}$, there is an acyclic cofibration $\mu_*: X_* \longrightarrow \text{Ex}(X_*)$.
- (3) Ex preserves (and in fact reflects) weak equivalences¹⁰.

Let X_* be an arbitrary simplicial space that is Reedy cofibrant and level-wise connected. Consider the following naturality square.

$$\begin{array}{ccc} |\Omega_* X_*| & \xrightarrow{\gamma} & \Omega|X_*| \\ \downarrow |\Omega_* \mu_*| & & \downarrow \Omega|\mu_*| \\ |\Omega_* \text{Ex } X_*| & \xrightarrow{\gamma} & \Omega|\text{Ex } X_*| \end{array}$$

The bottom arrow of this square (by Lemma 3.8) is a weak equivalence since $\text{Ex } X_*$ is fibrant. The left and right arrows of the square are weak equivalences since Ω preserves weak equivalences and $|-|$ preserves weak equivalences between Reedy cofibrant objects. By the two-out-of-three property for weak equivalences, it follows that the top arrow $\gamma: |\Omega_* X_*| \longrightarrow \Omega|X_*|$ is a weak equivalence.

REFERENCES

- [And78] D. W. Anderson. “Fibrations and geometric realizations”. In: *Bull. Amer. Math. Soc.* 84.5 (1978), pp. 765–788. ISSN: 0002-9904. DOI: 10.1090/S0002-9904-1978-14512-1. URL: <https://doi.org/10.1090/S0002-9904-1978-14512-1>.

¹⁰This follows from applications of the two-out-of-three property for weak equivalences.

- [Bec25] Jonathan Mock Beck. “The tripleability theorem”. In: *Repr. Theory Appl. Categ.* 31 (2025). With a preface by Robert Paré, p. 10.
- [BF78] A. K. Bousfield and E. M. Friedlander. “Homotopy theory of Γ -spaces, spectra, and bisimplicial sets”. In: *Geometric applications of homotopy theory (Proc. Conf., Evanston, Ill., 1977), II*. Vol. 658. Lecture Notes in Math. Springer, Berlin-New York, 1978, pp. 80–130. ISBN: 3-540-08859-8.
- [Bro06] Ronald Brown. *Topology and groupoids*. Third. With 1 CD-ROM (Windows, Macintosh and UNIX). BookSurge, LLC, Charleston, SC, 2006, pp. xxvi+512. ISBN: 1-4196-2722-8.
- [GJ99] Paul G. Goerss and John F. Jardine. *Simplicial homotopy theory*. Vol. 174. Progress in Mathematics. Birkhäuser Verlag, Basel, 1999, pp. xvi+510. ISBN: 3-7643-6064-X. DOI: 10.1007/978-3-0348-8707-6. URL: <https://doi.org/10.1007/978-3-0348-8707-6>.
- [Hov99] Mark Hovey. *Model categories*. Vol. 63. Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 1999, pp. xii+209. ISBN: 0-8218-1359-5.
- [Hu03] Po Hu. “ S -modules in the category of schemes”. In: *Mem. Amer. Math. Soc.* 161.767 (2003), pp. viii+125. ISSN: 0065-9266,1947-6221. DOI: 10.1090/memo/0767. URL: <https://doi.org/10.1090/memo/0767>.
- [Jar87] J. F. Jardine. “Simplicial presheaves”. In: *J. Pure Appl. Algebra* 47.1 (1987), pp. 35–87. ISSN: 0022-4049,1873-1376. DOI: 10.1016/0022-4049(87)90100-9. URL: [https://doi.org/10.1016/0022-4049\(87\)90100-9](https://doi.org/10.1016/0022-4049(87)90100-9).
- [KMZ26] Hana Jia Kong, J. Peter May, and Foling Zou. *The homotopical monadicity theorem*. To appear. 2026.
- [Lur17] Jacob Lurie. *Higher Algebra*. 2017. URL: <https://people.math.harvard.edu/~lurie/papers/HA.pdf>.
- [May09] J. P. May. “What precisely are E_∞ ring spaces and E_∞ ring spectra?” In: *New topological contexts for Galois theory and algebraic geometry (BIRS 2008)*. Vol. 16. Geom. Topol. Monogr. Geom. Topol. Publ., Coventry, 2009, pp. 215–282. DOI: 10.2140/gtm.2009.16.215. URL: <https://doi.org/10.2140/gtm.2009.16.215>.
- [May67] J. Peter May. *Simplicial objects in algebraic topology*. Vol. No. 11. Van Nostrand Mathematical Studies. D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto, Ont.-London, 1967, pp. vi+161.
- [May72] J. P. May. *The geometry of iterated loop spaces*. Vol. Vol. 271. Lecture Notes in Mathematics. Springer-Verlag, Berlin-New York, 1972, pp. viii+175.
- [May74] J. P. May. “ E_∞ spaces, group completions, and permutative categories”. In: *New developments in topology (Proc. Sympos. Algebraic Topology, Oxford, 1972)*. Vol. No. 11. London Math. Soc. Lecture Note Ser. Cambridge Univ. Press, London-New York, 1974, pp. 61–93.
- [Mor12] Fabien Morel. *$\mathbb{A}^1$ -algebraic topology over a field*. Vol. 2052. Lecture Notes in Mathematics. Springer, Heidelberg, 2012, pp. x+259. ISBN: 978-3-642-29513-3. DOI: 10.1007/978-3-642-29514-0. URL: <https://doi.org/10.1007/978-3-642-29514-0>.
- [MV99] Fabien Morel and Vladimir Voevodsky. “ $\mathbb{A}^1$ -homotopy theory of schemes”. In: *Inst. Hautes Études Sci. Publ. Math.* 90 (1999), pp. 45–143. ISSN: 0073-8301,1618-1913. URL: http://www.numdam.org/item?id=PMIHES_1999__90__45_0.
- [Ree73] C. L. Reedy. *Homotopy Theory of Model Categories*. Preprint. 1973. URL: <https://math.mit.edu/~psh/reedy.pdf>.
- [Rez10] Charles Rezk. *Toposes and Homotopy Toposes*. 2010. URL: https://ncatlab.org/nlab/files/Rezk_HomotopyToposes.pdf.
- [Rez14] C. Rezk. *When are homotopy limits compatible with homotopy base change?* Preprint. 2014. URL: <https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=7fa39aaddb0b1f169698696761e0add77dc64aa>.
- [SGA3V1] Philippe Gille and Patrick Polo, eds. *Schémas en groupes (SGA 3). Tome I. Propriétés générales des schémas en groupes*. annotated. Vol. 7. Documents Mathématiques (Paris) [Mathematical Documents (Paris)]. Séminaire de Géométrie Algébrique du Bois Marie 1962–64. [Algebraic Geometry Seminar of Bois Marie 1962–64], A seminar directed by M. Demazure and A. Grothendieck with the collaboration of M.

- Artin, J.-E. Bertin, P. Gabriel, M. Raynaud and J-P. Serre. Société Mathématique de France, Paris, 2011, pp. xxviii+610. ISBN: 978-2-85629-323-2.
- [SGA4V2] *Théorie des topos et cohomologie étale des schémas. Tome 2.* Vol. 270. Lecture Notes in Mathematics. Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4), Dirigé par M. Artin, A. Grothendieck et J. L. Verdier. Avec la collaboration de N. Bourbaki, P. Deligne et B. Saint-Donat. Springer-Verlag, Berlin-New York, 1972, pp. iv+418.
- [SØ12] Markus Spitzweck and Paul Arne Østvær. “Motivic twisted K -theory”. In: *Algebr. Geom. Topol.* 12.1 (2012), pp. 565–599. ISSN: 1472-2747,1472-2739. DOI: 10.2140/agt.2012.12.565. URL: <https://doi.org/10.2140/agt.2012.12.565>.
- [Voe00] Vladimir Voevodsky. *Some open problems in the motivic homotopy theory*. 2000. URL: https://www.math.ias.edu/Voevodsky/files/files-original/Dropbox/Published_papers/Motives/Open/Stage2/open.pdf.
- [Voe96] Vladimir Voevodsky. *The Milnor Conjecture*. Preprint. 1996. URL: <http://k-theory.org/0170/>.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109
Email address: avsriniv@umich.edu