RefID: 1, There and Back Again: A Review of Residency and Return Migrations in Sharks, with Implications for Population Structure and Management. Chapman DD, Feldheim KA, Papastamatiou Y, Hueter RE

The overexploitation of sharks has become a global environmental issue in need of a comprehensive and multifaceted management response. Tracking studies are beginning to elucidate how shark movements shape the internal dynamics and structure of populations, which determine the most appropriate scale of these management efforts.

Tracked sharks frequently either remain in a restricted geographic area for an extended period of time (residency) or return to a previously resided-in area after making long-distance movements (site fidelity). Genetic studies have shown that some individuals of certain species preferentially return to their exact birthplaces (natal philopatry) or birth regions (regional philopatry) for either parturition or mating, even though they make long-distance movements that would allow them to breed elsewhere. More than 80 peer-reviewed articles, constituting the majority of published shark tracking and population genetic studies, provide evidence of at least one of these behaviors in a combined 31 shark species from six of the eight extant orders.

Residency, site fidelity, and philopatry can alone or in combination structure many coastal shark populations on finer geographic scales than expected based on their potential for dispersal. This information should therefore be used to scale and inform assessment, management, and conservation activities intended to restore depleted shark populations. Expected final online publication date for the Annual Review of Marine Science Volume 7 is January 03, 2015.

1. **Article ID**
   formula: R + 2022 + REFID
   ...

Strata formula

Ex: Article reported COD data for
1) children aged 0-1m and 1-59m for 3 regions; and
2) mortality rates for 2000, 2010, and 2015

= [2(age group)*3(regions)] + 3(mortality rates)

Note on sex strata: we are interested in the sex split for 5-19 year olds but not under-5's. (i.e. if male, female, and total is available, extracte 5 strata and "male" "female" for 5-19 strata. If only "male" "female" is available, extract for all ages exactly as presented in article (DO NOT create a "total" strata for under-5s)

...

Total number of strata ...

Ex: from the formula above = 9

2. Study design

- ◯ Epidemiologic
- ◯ National representative survey
- ◯ Sub-national survey
- ◯ Vital statistics registration system / Sample registration system (nationwide)
- ◯ Control arm of intervention (includes control time period or control intervention)
- ◯ Demographic Surveillance System (DSS) (not nationwide)
- ◯ Other

3. Reference type

- ◯ Internet
- ◯ Journal
- ◯ Monograph
- ◯ PhD thesis
- ◯ DSS (not in a Journal)
- ◯ Report
- ◯ Unpublished
- ◯ Other

Clear Response ()

4. Country Select an Answer

5. Describe the location of the study area

(General description of the study area including city, state, etc.)

6. Place of residence of study population ◯ Rural ◯ Urban ◯ Mixed rural and urban ◯ Not specified **Clear Res**

7. Risk of malaria among study children ◯ Negligible ◯ Endemic ◯ Epidemic ◯ Not specified **Clear Response**

8. Surveillance years for mortality study:

Start (yyyy): ...

End (yyyy): ...

9. Age range of follow-up for mortality study:

Unit: Select an Answer

Youngest: ...

Oldest: ...

10. Methods for gathering Cause of Death information

- ◯ Verbal Autopsy
- ◯ Verbal Autopsy+Medical Records
- ◯ Medical Records
- ◯ Vital Registry
- ◯ Vital Registry+Medical Records
- ◯ Other
- ◯ Cannot be determined

Clear Response ()

11. Methods for assigning Cause of Death

- ◯ Single expert
- ◯ Two independent experts, one expert selected in case of discrepency
- ◯ Two independent experts, achieve consensus among 2 experts in case of discrepency
- ◯ Two independent experts, 3rd independent expert in case of discrepency
- ◯ Two independent experts, panel in case of discrepency
- ◯ Three independent experts, one expert selected in case of discrepency
- ◯ Three independent experts, achieve consensus among 3 experts in case of discrepency
- ◯ Three independent experts, 4th independent expert in case of discrepency
- ◯ Panel
- ◯ Computer Algorithm
- ◯ Other standarized procedure (specify)
- ◯ Unstandardized procedure
- ◯ Method not reported in article

Clear Response ()

12. Comments

Copyright © 2008-2026 DistillerSR Inc., All Rights Reserved. Patent Pending.