RefID: 1, There and Back Again: A Review of Residency and Return Migrations in Sharks, with Implications for Population Structure and Management. Chapman DD, Feldheim KA, Papastamatiou Y, Hueter RE

The overexploitation of sharks has become a global environmental issue in need of a comprehensive and multifaceted management response. Tracking studies are beginning to elucidate how shark movements shape the internal dynamics and structure of populations, which determine the most appropriate scale of these management efforts.

Tracked sharks frequently either remain in a restricted geographic area for an extended period of time (residency) or return to a previously resided-in area after making long-distance movements (site fidelity). Genetic studies have shown that some individuals of certain species preferentially return to their exact birthplaces (natal philopatry) or birth regions (regional philopatry) for either parturition or mating, even though they make long-distance movements that would allow them to breed elsewhere. More than 80 peer-reviewed articles, constituting the majority of published shark tracking and population genetic studies, provide evidence of at least one of these behaviors in a combined 31 shark species from six of the eight extant orders.

Residency, site fidelity, and philopatry can alone or in combination structure many coastal shark populations on finer geographic scales than expected based on their potential for dispersal. This information should therefore be used to scale and inform assessment, management, and conservation activities intended to restore depleted shark populations. Expected final online publication date for the Annual Review of Marine Science Volume 7 is January 03, 2015.

Risk Factors / Covariates Form

Article reported risk factor data separately by strata: ...

If yes, enter strata_id

If no, enter "no"

Risk Factor	Value reported in article	Value reported elsewhere	Source of data elsewhere	Comment
GNI per capita	...	...	...	...
Proportion of population living in urban area	(%) ...	(%) ...	...	...
Proportion of adult females in the study population who are literate	(%) ...	(%) ...	...	...
Proportion of study children using improved water sources (i.e. piped water into dwelling, plot or yard, public tap/standpipe, tube well/borehole, protected dug well, protected spring, rainwater)	(%) ...	(%) ...	...	...

Proportion of study children using improved sanitation facilities (i.e. flush/pour-flush to: piped water system, septic tank, pit latrine; ventilated improved pit latrine, pit latrine with slap, composting toilet)	(%) ...	(%) ...	...	...
Proportion of birth in the study area attended by a skilled attendant (i.e. doctors, nurses, or midwives)	(%) ...	(%) ...	...	...
Exclusive breastfeeding	(%) ...	(%) ...	...	...
Proportion of study children who are moderate/ severe underweight: (<-2 SDs from median WA of the NCHS/WHO reference population)	(%) ...	(%) ...	...	...
Proportion of study children who are moderate/ severe wasted: (<-2 SDs from median WH of the NCHS/WHO reference population)	(%) ...	(%) ...	...	...

Proportion of study children who are moderate/ severe stunted: (<-2 SDs from median HA of the NCHS/WHO reference population)	(%) ...	(%) ...	...	...
Proportion of study children with diarrhea receiving oral rehydration	Type: Select an Answer (%) ...	(%) ...	...	...
Meningitis epidemic occurred during study period	(Y/N) ◯ Yes ◯ No **Clear Response ()**	(Y/N) ◯ Yes ◯ No **Clear Response ()**	...	...
Measles vaccine coverage	(%) ...	(%) ...	...	...
Hib3 vaccine coverage	(%) ...	(%) ...	...	...
Pneumococcal vaccine coverage	(%) ...	(%) ...	...	...
Rotavirus vaccine coverage	(%) ...	(%) ...	...	...

Comments

Copyright © 2008-2026 DistillerSR Inc., All Rights Reserved. Patent Pending.