

# Finding Order in Metric Structures

Aaron Anderson (joint with Diego Bejarano)

UPenn

February 3, 2026



# Metric Structures

## Definition

A *metric language* is just like a regular first-order language, consisting of functions and relations.

## Definition

A *metric structure* consists of:

- A complete metric space of diameter  $\leq 1$
- For each  $n$ -ary function symbol, a uniformly continuous function  $M^n \rightarrow M$
- For each  $n$ -ary relation symbol, a uniformly continuous function  $M^n \rightarrow [0, 1]$

# Formulas

## Definition

An *atomic formula* is defined as usual, except instead of  $=$ , the basic relation is  $d(x, y)$ .

## Definition

A *formula* is

- An atomic formula
- $u(\phi_1, \dots, \phi_n)$  where  $\phi_i$ s are formulas and  $u : [0, 1]^n \rightarrow [0, 1]$  is continuous
- $\sup_x \phi$  or  $\inf_x \phi$

# Making Linear Orders Metric

- Most famous examples of metric structures are stable or not NIP - how do we put an order on one?

# Making Linear Orders Metric

- Most famous examples of metric structures are stable or not NIP - how do we put an order on one?
- Itai Ben Yaacov has described Ordered Real Closed Metric Valued Fields, but making the metric space bounded complicated things.

# Making Linear Orders Metric

- Most famous examples of metric structures are stable or not NIP - how do we put an order on one?
- Itai Ben Yaacov has described Ordered Real Closed Metric Valued Fields, but making the metric space bounded complicated things.
- Diego Bejarano and I have simplified this approach.

# Metric Linear Orders

- Call  $M$  a *metric linear order* if
  - $M$  has a complete metric of diameter  $\leq 1$
  - $M$  has a linear order
  - open balls are order-convex.
- $M$  is a metric structure in the language  $\{r\}$ , with

$$r(x, y) = \begin{cases} 0 & x \leq y \\ d(x, y) & y \leq x \end{cases}$$

- Think of  $r(x, y)$  as “the amount  $x$  is greater than  $y$ ,” or

$$r(x, y) = d(x, (\infty, y]).$$

# Axiomatizing Metric Linear Orders

## Definition

A *theory* is a set of *conditions*  $\phi(x) = 0$ .



# Axiomatizing Metric Linear Orders

## Definition

A *theory* is a set of *conditions*  $\phi(x) = 0$ .

## Theorem (A., Bejarano)

*Metric linear orders are axiomatized in  $\{r\}$  by*

- $\sup_{x,y} |(r(x,y) + r(y,x)) - d(x,y)| = 0$

# Axiomatizing Metric Linear Orders

## Definition

A *theory* is a set of *conditions*  $\phi(x) = 0$ .

## Theorem (A., Bejarano)

*Metric linear orders are axiomatized in  $\{r\}$  by*

- $\sup_{x,y} |(r(x,y) + r(y,x)) - d(x,y)| = 0$
- $\sup_{x,y} \min\{r(x,y), r(y,x)\} = 0$

# Axiomatizing Metric Linear Orders

## Definition

A *theory* is a set of *conditions*  $\phi(x) = 0$ .

## Theorem (A., Bejarano)

*Metric linear orders are axiomatized in  $\{r\}$  by*

- $\sup_{x,y} |(r(x,y) + r(y,x)) - d(x,y)| = 0$
- $\sup_{x,y} \min\{r(x,y), r(y,x)\} = 0$
- $\sup_{x,y,z} r(x,z) \dot{-} (r(x,y) + r(y,z)) = 0$

# Ultrametric Dense Linear Orders

- DLO and  $\text{Th}(\mathbb{Z}, <)$  are two useful completions of the theory of linear orders

# Ultrametric Dense Linear Orders

- DLO and  $\text{Th}(\mathbb{Z}, <)$  are two useful completions of the theory of linear orders
- We seek analogous completions of MLO

# Ultrametric Dense Linear Orders

- DLO and  $\text{Th}(\mathbb{Z}, <)$  are two useful completions of the theory of linear orders
- We seek analogous completions of MLO
- For simplicity, assume the metric is an ultrametric.

# Axiomatizing Ultrametric Dense Linear Orders

## Definition

Let UDLO be the theory of *ultrametric-dense linear orders*, consisting of MLO with the following axioms:

- $d(x, z) \leq \max(d(x, y), d(y, z))$
- For any rational  $p \in \mathbb{Q} \cap [0, 1]$ ,  $\sup_x \inf_y |r(x, y) - p| = 0$
- For any rational  $p \in \mathbb{Q} \cap [0, 1]$ ,  $\sup_x \inf_y |r(y, x) - p| = 0$ .

Basically, the distances from  $x$  to  $y > x$  are dense in  $[0, 1]$ .

# Stable Diversion: Dense Ultrametrics

Call an ultrametric space *dense* if the set of distances to any point is dense in  $[0, 1]$ .

## Fact (Conant)

*The theory of dense ultrametrics is complete and has QE, but is not  $\aleph_0$ -categorical.*



# Stable Diversion: Dense Ultrametrics

Call an ultrametric space *dense* if the set of distances to any point is dense in  $[0, 1]$ .

## Fact (Conant)

*The theory of dense ultrametrics is complete and has QE, but is not  $\aleph_0$ -categorical.*

(A separable ultrametric space has countably many distances!)

# Stable Diversion: Dense Ultrametrics

Call an ultrametric space *dense* if the set of distances to any point is dense in  $[0, 1]$ .

## Fact (Conant)

*The theory of dense ultrametrics is complete and has QE, but is not  $\aleph_0$ -categorical.*

(A separable ultrametric space has countably many distances!)

## Lemma (A., Bejarano)

*The theory of dense ultrametrics is the model companion of the theory of ultrametrics, and is approximately  $\aleph_0$ -categorical.*

# Ultrametric Dense Linear Orders

## Theorem (A., Bejarano)

- UDLO *is complete*
- UDLO *eliminates quantifiers*
- UDLO *is the model companion of the theory of ultrametric linear orders*
- UDLO *is approximately  $\aleph_0$ -categorical*

# Ultrametric Dense Linear Orders

## Theorem (A., Bejarano)

- UDLO *is complete*
- UDLO *eliminates quantifiers*
- UDLO *is the model companion of the theory of ultrametric linear orders*
- UDLO *is approximately  $\aleph_0$ -categorical*

## Theorem (A., Bejarano)

*In a model of UDLO,*

- *the metric and order topologies agree*
- *dcl is closure*

# A Separable UDLO

- Let  $S \subseteq (0, 1]$  be countable, dense.

# A Separable UDLO

- Let  $S \subseteq (0, 1]$  be countable, dense.
- Let  $U_S = \{f : S \rightarrow \mathbb{Q} : \forall r > 0, \{s > r : f(s) \neq 0\} \text{ is finite.}\}$ .

# A Separable UDLO

- Let  $S \subseteq (0, 1]$  be countable, dense.
- Let  $U_S = \{f : S \rightarrow \mathbb{Q} : \forall r > 0, \{s > r : f(s) \neq 0\} \text{ is finite.}\}$ .
- Let  $d(f, g) = \max\{x : f(x) \neq g(x)\}$ .

# A Separable UDLO

- Let  $S \subseteq (0, 1]$  be countable, dense.
- Let  $U_S = \{f : S \rightarrow \mathbb{Q} : \forall r > 0, \{s > r : f(s) \neq 0\} \text{ is finite.}\}$ .
- Let  $d(f, g) = \max\{x : f(x) \neq g(x)\}$ .
- Let  $f(x) < g(x)$  when  $f(d(f, g)) < g(d(f, g))$ .



# A Separable UDLO

- Let  $S \subseteq (0, 1]$  be countable, dense.
- Let  $U_S = \{f : S \rightarrow \mathbb{Q} : \forall r > 0, \{s > r : f(s) \neq 0\} \text{ is finite.}\}$ .
- Let  $d(f, g) = \max\{x : f(x) \neq g(x)\}$ .
- Let  $f(x) < g(x)$  when  $f(d(f, g)) < g(d(f, g))$ .

Lemma (A., Bejarano)

$U_S \models \text{UDLO}$ .

# A Separable UDLO

- Let  $S \subseteq (0, 1]$  be countable, dense.
- Let  $U_S = \{f : S \rightarrow \mathbb{Q} : \forall r > 0, \{s > r : f(s) \neq 0\} \text{ is finite.}\}$ .
- Let  $d(f, g) = \max\{x : f(x) \neq g(x)\}$ .
- Let  $f(x) < g(x)$  when  $f(d(f, g)) < g(d(f, g))$ .

Lemma (A., Bejarano)

$U_S \models \text{UDLO}$ .

Theorem (Van Thé)

$U_S$  has extremely amenable automorphism group, following from Fraïssé theory in the **discrete-logic** language of ordered  $S$ -valued metric spaces.

# o-Minimality in Discrete Logic

## Fact

*If  $M$  expands a linear order, TFAE:*

- *every formula  $\phi(x)$  in one variable is qf-definable in  $\{<\}$*
- *every formula  $\phi(x)$  in one variable is a finite union of intervals.*
- *If these happen,  $M$  is o-minimal.*

# *o*-Minimality in Discrete Logic

## Fact

*If  $M$  expands a linear order, TFAE:*

- *every formula  $\phi(x)$  in one variable is qf-definable in  $\{<\}$*
- *every formula  $\phi(x)$  in one variable is a finite union of intervals.*
- If these happen,  $M$  is *o-minimal*.
- How do we describe these properties for MLOs?

# Metric $\mathcal{o}$ -Minimality

## Theorem (A., Bejarano)

*If  $M$  expands a metric linear order, TFAE:*

- *every formula  $\phi(x)$  in one variable is qf-definable in  $\{r\}$*
- *every formula  $\phi(x)$  in one variable is regulated (a uniform limit of step functions).*

# Metric *o*-Minimality

## Theorem (A., Bejarano)

*If  $M$  expands a metric linear order, TFAE:*

- *every formula  $\phi(x)$  in one variable is qf-definable in  $\{r\}$*
- *every formula  $\phi(x)$  in one variable is regulated (a uniform limit of step functions).*

## Definition

If  $M$  expands a metric linear order, call  $M$  *o-minimal* if every formula  $\phi(x)$  in one variable satisfies these equivalent properties.

# Metric *o*-Minimality

## Theorem (A., Bejarano)

*If  $M$  expands a metric linear order, TFAE:*

- *every formula  $\phi(x)$  in one variable is qf-definable in  $\{r\}$*
- *every formula  $\phi(x)$  in one variable is regulated (a uniform limit of step functions).*

## Definition

If  $M$  expands a metric linear order, call  $M$  *o-minimal* if every formula  $\phi(x)$  in one variable satisfies these equivalent properties.

By QE, a model of UDLO is *o*-minimal.

# Regulated Functions

Regulated functions  $[a, b] \rightarrow \mathbb{R}$  were defined by Bourbaki.

## Lemma (A. Bejarano)

If  $M$  is a linear order,  $f : M \rightarrow [0, 1]$ , TFAE:

- ①  $f$  is a uniform limit of step functions
- ② For any  $a < b$ ,  $M$  can be partitioned into finitely many intervals on which either  $f(x) > a$  or  $f(x) < b$ .

If  $M \models \text{UDLO}$ , and  $f : M \rightarrow M$  is definable ( $d(f(x), y)$  is a formula), then  $f$  is continuous and satisfies (2).



# Metric Valued Fields

## Definition

A *metric valued field* is a field  $K$  with an absolute value  $|\cdot| : K \rightarrow \mathbb{R}^{\geq 0}$  satisfying valuation axioms:

- $|x| = 0 \iff x = 0$
- $|xy| = |x||y|$
- $|x + y| \leq \max(|x|, |y|)$

which gives rise to an ultrametric  $d(x, y) = |x - y|$ .

# Real Closed Metric Valued Fields

## Definition

If  $K$  is a metric valued field equipped with an order,  $K \equiv \mathbb{R}$  as ordered fields in classical logic, and  $|\cdot|$  takes values outside  $\{0, 1\}$ , call  $K$  a *real closed metric valued field*.

These are almost metric structures, but the metric is unbounded, a problem for continuous logic.

# Metric Valued Fields, à Rideau-Kikuchi, Scanlon, Simon

- Typically we just deal with the ball  $B_{\leq 1}(0)$ .
- For a metric valued field  $K$ ,  $\{x \in K : |x| \leq 1\}$  is a subring - the *valuation ring*.
- This is a metric structure in the ring language.

# Metric Valued Fields, à Rideau-Kikuchi, Scanlon, Simon

- Typically we just deal with the ball  $B_{\leq 1}(0)$ .
- For a metric valued field  $K$ ,  $\{x \in K : |x| \leq 1\}$  is a subring - the *valuation ring*.
- This is a metric structure in the ring language.
- Unfortunately, these are elementarily equivalent to other convex subrings of metric valued fields.

# Ordered Real Closed Metric Valuation Rings

## Theorem (A., Bejarano)

- *Models of our theory ORCMVR are convex subrings of real closed metric valued fields.*
- *These have quantifier-elimination once we add a divisibility relation.*
- *These can only be weakly o-minimal.*

# Real Closed Metric Valued Fields à Ben Yaacov

- Itai Ben Yaacov studied the projective line instead

# Real Closed Metric Valued Fields à Ben Yaacov

- Itai Ben Yaacov studied the projective line instead
- Artem was right - we need to view these as *cyclic* orders

# Real Closed Metric Valued Fields à Ben Yaacov

- Itai Ben Yaacov studied the projective line instead
- Artem was right - we need to view these as *cyclic* orders

## Theorem (A., Bejarano)

*Any model of ORCMVF (the projective line) is an o-minimal expansion of a model of UDCO, the complete theory of ultrametric dense cyclic orders.*



Thank you, CMU!