

# Finding Order in Metric Structures

Aaron Anderson (joint with Diego Bejarano)

UPenn

May 13, 2025

# Metric Structures

## Definition

A *metric language* is just like a regular first-order language, consisting of functions and relations.

## Definition

A *metric structure* consists of:

- A complete metric space of diameter 1
- For each  $n$ -ary function symbol, a uniformly continuous function  $M^n \rightarrow M$
- For each  $n$ -ary relation symbol, a uniformly continuous function  $M^n \rightarrow [0, 1]$

# Formulas

## Definition

An *atomic formula* is defined as usual, except instead of  $=$ , the basic relation is  $d(x, y)$ .

## Definition

A *formula* is

- An atomic formula
- $u(\phi_1, \dots, \phi_n)$  where  $\phi_i$ s are formulas and  $u : [0, 1]^n \rightarrow [0, 1]$  is continuous
- $\sup_x \phi$  or  $\inf_x \phi$

## Definition

A *definable predicate* is a uniform limit of formulas.

# Making Linear Orders Metric

- Most famous examples of metric structures are stable or not NIP - how do we put an order on one?

# Making Linear Orders Metric

- Most famous examples of metric structures are stable or not NIP - how do we put an order on one?
- Itai Ben Yaacov has described Ordered Real Closed Metric Valued Fields, but making the metric space bounded complicated things.

# Making Linear Orders Metric

- Most famous examples of metric structures are stable or not NIP - how do we put an order on one?
- Itai Ben Yaacov has described Ordered Real Closed Metric Valued Fields, but making the metric space bounded complicated things.
- Diego Bejarano and I are working to simplify this approach.

# Metric Linear Orders

- Call  $M$  a *metric linear order* if
  - $M$  has a bounded complete metric
  - $M$  has a linear order
  - open balls are order-convex.
- $M$  is a metric structure in the language  $\{r\}$ , with

$$r(x, y) = \begin{cases} 0 & x \leq y \\ d(x, y) & y \leq x \end{cases}$$

- Think of  $r(x, y)$  as “the amount  $x$  is greater than  $y$ .”

# Axiomatizing Metric Linear Orders

## Definition

A *theory* is a set of *conditions*  $\phi(x) = 0$ .



# Axiomatizing Metric Linear Orders

## Definition

A *theory* is a set of *conditions*  $\phi(x) = 0$ .

## Theorem (A., Bejarano)

*Metric linear orders are axiomatized in  $\{r\}$  by*

- $\sup_{x,y} |(r(x,y) + r(y,x)) - d(x,y)| = 0$

# Axiomatizing Metric Linear Orders

## Definition

A *theory* is a set of *conditions*  $\phi(x) = 0$ .

## Theorem (A., Bejarano)

*Metric linear orders are axiomatized in  $\{r\}$  by*

- $\sup_{x,y} |(r(x,y) + r(y,x)) - d(x,y)| = 0$
- $\sup_{x,y} \min\{r(x,y), r(y,x)\} = 0$

# Axiomatizing Metric Linear Orders

## Definition

A *theory* is a set of *conditions*  $\phi(x) = 0$ .

## Theorem (A., Bejarano)

*Metric linear orders are axiomatized in  $\{r\}$  by*

- $\sup_{x,y} |(r(x,y) + r(y,x)) - d(x,y)| = 0$
- $\sup_{x,y} \min\{r(x,y), r(y,x)\} = 0$
- $\sup_{x,y,z} r(x,z) - (r(x,y) + r(y,z)) = 0$

# Ultrametric Dense Linear Orders

- DLO and  $\text{Th}(\mathbb{Z}, <)$  are two useful completions of the theory of linear orders

# Ultrametric Dense Linear Orders

- DLO and  $\text{Th}(\mathbb{Z}, <)$  are two useful completions of the theory of linear orders
- We seek an analogous completion of MLO

# Ultrametric Dense Linear Orders

- DLO and  $\text{Th}(\mathbb{Z}, <)$  are two useful completions of the theory of linear orders
- We seek an analogous completion of MLO
- For simplicity, assume the metric is an ultrametric

# Ultrametric Dense Linear Orders

- DLO and  $\text{Th}(\mathbb{Z}, <)$  are two useful completions of the theory of linear orders
- We seek an analogous completion of MLO
- For simplicity, assume the metric is an ultrametric

## Definition

Let UDLO be the theory of *ultrametric-dense linear orders*, consisting of MLO with the following axioms:

- $d(x, z) \leq \max(d(x, y), d(y, z))$
- For any rational  $p \in \mathbb{Q} \cap [0, 1]$ ,  $\sup_x \inf_y |r(x, y) - p| = 0$
- For any rational  $p \in \mathbb{Q} \cap [0, 1]$ ,  $\sup_x \inf_y |r(y, x) - p| = 0$ .

# Ultrametric Dense Linear Orders

## Theorem (A., Bejarano)

- *UDLO is complete*
- *UDLO eliminates quantifiers*
- *UDLO is the model companion of the theory of ultrametric linear orders*



# Ultrametric Dense Linear Orders

## Theorem (A., Bejarano)

- *UDLO is complete*
- *UDLO eliminates quantifiers*
- *UDLO is the model companion of the theory of ultrametric linear orders*
- *The metric and order topologies agree in a model of UDLO*

# Ultrametric Dense Linear Orders

## Theorem (A., Bejarano)

- *UDLO is complete*
- *UDLO eliminates quantifiers*
- *UDLO is the model companion of the theory of ultrametric linear orders*
- *The metric and order topologies agree in a model of UDLO*
- *dcl in UDLO is metric closure*

# Ultrametric Dense Linear Orders

## Theorem (A., Bejarano)

- *UDLO is complete*
- *UDLO eliminates quantifiers*
- *UDLO is the model companion of the theory of ultrametric linear orders*
- *The metric and order topologies agree in a model of UDLO*
- *dcl in UDLO is metric closure*
- *Orders Van Thé put on Urysohn ultrametric spaces model UDLO*

# Constructing a Separable UDLO

- Let  $S \subseteq (0, 1]$  be countable, dense.

# Constructing a Separable UDLO

- Let  $S \subseteq (0, 1]$  be countable, dense.
- Let  $U_S = \{f : S \rightarrow \mathbb{Q} : \forall r > 0, \{s > r : f(s) \neq 0\} \text{ is finite.}\}$ .

# Constructing a Separable UDLO

- Let  $S \subseteq (0, 1]$  be countable, dense.
- Let  $U_S = \{f : S \rightarrow \mathbb{Q} : \forall r > 0, \{s > r : f(s) \neq 0\} \text{ is finite.}\}$ .
- Let  $d(f, g) = \max\{x : f(x) \neq g(x)\}$ .

# Constructing a Separable UDLO

- Let  $S \subseteq (0, 1]$  be countable, dense.
- Let  $U_S = \{f : S \rightarrow \mathbb{Q} : \forall r > 0, \{s > r : f(s) \neq 0\} \text{ is finite.}\}$ .
- Let  $d(f, g) = \max\{x : f(x) \neq g(x)\}$ .
- Let  $f(x) < g(x)$  when  $f(d(f, g)) < g(d(f, g))$ .

# Constructing a Separable UDLO

- Let  $S \subseteq (0, 1]$  be countable, dense.
- Let  $U_S = \{f : S \rightarrow \mathbb{Q} : \forall r > 0, \{s > r : f(s) \neq 0\} \text{ is finite.}\}$ .
- Let  $d(f, g) = \max\{x : f(x) \neq g(x)\}$ .
- Let  $f(x) < g(x)$  when  $f(d(f, g)) < g(d(f, g))$ .

Lemma (A., Bejarano)

$U_S \models \text{UDLO}$ .



# Constructing a Separable UDLO

- Let  $S \subseteq (0, 1]$  be countable, dense.
- Let  $U_S = \{f : S \rightarrow \mathbb{Q} : \forall r > 0, \{s > r : f(s) \neq 0\} \text{ is finite.}\}$ .
- Let  $d(f, g) = \max\{x : f(x) \neq g(x)\}$ .
- Let  $f(x) < g(x)$  when  $f(d(f, g)) < g(d(f, g))$ .

Lemma (A., Bejarano)

$U_S \models \text{UDLO}$ .

Theorem (Van Thé)

*$U_S$  has extremely amenable automorphism group, following from Fraïssé theory in the discrete-logic language of ordered  $S$ -valued metric spaces.*

# o-Minimality in Discrete Logic

## Fact

*If  $M$  expands a linear order, TFAE:*

- *every formula  $\phi(x)$  in one variable is qf-definable in  $\{<\}$*
- *every formula  $\phi(x)$  in one variable is a finite union of intervals.*
- *If these happen,  $M$  is o-minimal.*

# o-Minimality in Discrete Logic

## Fact

*If  $M$  expands a linear order, TFAE:*

- *every formula  $\phi(x)$  in one variable is qf-definable in  $\{<\}$*
- *every formula  $\phi(x)$  in one variable is a finite union of intervals.*
- If these happen,  $M$  is *o-minimal*.
- How do we describe these properties for MLOs?

# Metric o-Minimality

## Theorem (A., Bejarano)

*If  $M$  expands a metric linear order, TFAE:*

- *every predicate  $\phi(x)$  in one variable is qf-definable in  $\{r\}$*
- *every predicate  $\phi(x)$  in one variable is regulated (a uniform limit of step functions).*

# Metric o-Minimality

## Theorem (A., Bejarano)

*If  $M$  expands a metric linear order, TFAE:*

- *every predicate  $\phi(x)$  in one variable is qf-definable in  $\{r\}$*
- *every predicate  $\phi(x)$  in one variable is regulated (a uniform limit of step functions).*

## Definition

If  $M$  expands a metric linear order, call  $M$  *o-minimal* if every predicate  $\phi(x)$  in one variable satisfies these equivalent properties.

# Metric o-Minimality

## Theorem (A., Bejarano)

*If  $M$  expands a metric linear order, TFAE:*

- *every predicate  $\phi(x)$  in one variable is qf-definable in  $\{r\}$*
- *every predicate  $\phi(x)$  in one variable is regulated (a uniform limit of step functions).*

## Definition

If  $M$  expands a metric linear order, call  $M$  *o-minimal* if every predicate  $\phi(x)$  in one variable satisfies these equivalent properties.

By QE, a model of UDLO is o-minimal.

# Definable Functions

## Definition

A function  $f : M \rightarrow M$  is *definable* when  $d(f(x), y)$  is.

# Definable Functions

## Definition

A function  $f : M \rightarrow M$  is *definable* when  $d(f(x), y)$  is.

## Theorem (A., Bejarano)

*If  $f : M \rightarrow M$  is definable in an o-minimal metric structure  $M$ , then for all  $a < b$ ,  $M$  can be partitioned into finitely many intervals on which either  $f(x) > a$  or  $f(x) < b$ .*



# Definable Functions

## Definition

A function  $f : M \rightarrow M$  is *definable* when  $d(f(x), y)$  is.

## Theorem (A., Bejarano)

*If  $f : M \rightarrow M$  is definable in an o-minimal metric structure  $M$ , then for all  $a < b$ ,  $M$  can be partitioned into finitely many intervals on which either  $f(x) > a$  or  $f(x) < b$ .*

## Proof.

The definable predicate  $r(f(x), a)$  is regulated. Approximate this with an appropriate step function, and partition into the intervals on which it is constant. □

# Definable Sets

## Definition

A set  $D \subseteq M$  is *definable* when it is closed and  $\inf_{y \in D} d(x, y)$  is a definable predicate.

## Theorem (Definable completeness: A., Bejarano)

Let  $M$  be an *o-minimal metric structure* and  $D \subset M$  a definable set. If  $D$  is bounded above (resp. below), then  $D$  has a least upper bound (resp. greatest lower bound).

## Theorem (A., Bejarano)

Let  $M$  be an *o-minimal expansion of MDLO* and  $D \subset M$  a definable set. The complement of  $D$  is a union of countably many intervals, with only finitely many of diameter  $\geq \varepsilon$  for any  $\varepsilon > 0$ .

# Cell Decomposition

By using the bounded alternation numbers of (weakly) regulated functions, we can build distal cell decompositions:

**Theorem (A., Bejarano)**

*Any (weakly) o-minimal metric structure is distal.*

# Cell Decomposition

By using the bounded alternation numbers of (weakly) regulated functions, we can build distal cell decompositions:

## Theorem (A., Bejarano)

*Any (weakly)  $\mathcal{o}$ -minimal metric structure is distal.*

Our main goal is to find more specific  $\mathcal{o}$ -minimal cell decompositions for definable predicates and sets.

Thank you, ASL Model Theory Session!