
TUTORIAL 3

1 Recursive division

Let a and b be two polynomials in $K[x]$ such that $\deg a = 4n$ and $\deg b = 2n$ and take n to be a power of 2. We decompose a and b such that $a(x) = a_h(x)x^{2n} + a_l(x)$ and $b(x) = b_h(x)x^n + b_l(x)$, where $\deg a_h, \deg a_l \leq 2n$ and $\deg b_h, \deg b_l \leq n$.

Consider $D(n)$ as the complexity, in number of arithmetic operations over K , required to perform the Euclidean division of a degree $2n$ polynomial by a degree n polynomial. Similarly, we denote by $M(n)$ the complexity of multiplying two degree n polynomials over R .

We perform the Euclidean division of a_h by b_h (i.e. $a_h = b_h q_h + r_h$, $\deg r_h < \deg b_h$).

1. Show that $\deg(a - bq_h x^n) < 3n$ and that $a - bq_h x^n$ is computable using $D(n) + M(n) + O(n)$ operations.
2. Show that we can finish dividing a by b using another $D(n) + M(n) + O(n)$ operations.
3. What is the value of $D(n)$ if $M(n) = n^\alpha$, $\alpha > 1$?
4. Same question as before, for $M(n) = n(\log n)^\alpha$, $\alpha > 1$.

2 Applications of the Extended Euclidean Algorithm (EEA)

2.1 Computing the inverse

1. Let n be an integer, and $0 \leq a < n$ be such that $\gcd(a, n) = 1$. Give an algorithm that computes $a^{-1} \bmod n$ in time $O(M(\log n) \log \log n)$.
2. Let $P \in K[X]$ be a polynomial of degree d with coefficients in a field K and $Q \in K[X]$ be a polynomial of degree less than d , such that $\gcd(P, Q) = 1$. Prove that Q is invertible modulo P and give an algorithm to compute its inverse using $O(M(d) \log d)$ operations in K .

2.2 Diophantine equation

The aim of this exercise is to describe the set of all solutions (u, v) of the equation

$$au + bv = t \tag{1}$$

1. Show that if $(u, v) = (s_1, s_2)$ is a solution of (1), the general solution is of the form $(u, v) = (s_1 + s'_1, s_2 + s'_2)$ for (s'_1, s'_2) satisfying $as'_1 + bs'_2 = 0$.
2. Find all solutions of $au + bv = 0$ for a, b coprime.
3. Find a solution of (1) for a, b coprime.
4. Observe that t must be divisible by $\gcd(a, b)$.
5. Using the previous questions, give the general solution of (1).

3 Rational function reconstruction

Let K be a field, $m \in K[X]$ of degree $n > 0$, and $f \in K[X]$ such that $\deg f < n$. For a fixed $k \in \{1, \dots, n\}$, we want to find a pair of polynomials $(r, t) \in K[X]^2$, satisfying

$$r = t \cdot f \pmod{m}, \quad \deg r < k, \quad \deg t \leq n - k \quad \text{and} \quad t \neq 0 \quad (2)$$

1. Consider $A(X) = \sum_{l=0}^{N-1} a_l X^l \in K[X]$ a polynomial. Show that if $A(X) = P(X)/Q(X) \pmod{X^N}$, where $P, Q \in K[X]$, $Q(0) = 1$ and $\deg P < \deg Q$, then the coefficients of A , starting from $a_{\deg Q}$ can be computed as a linear recurrent sequence of previous $\deg Q$ coefficients of A . What can you say in the converse setting when the coefficients of A satisfy a linear recurrence relation?
2. Inside (2), consider the case when $m = x^n$. Describe a linear algebra-based method for finding some t and r . (Hint: do **not** use the previous question).
3. Show that, if (r_1, t_1) and (r_2, t_2) are two pairs of polynomials that satisfy (2), then we have $r_1 t_2 = r_2 t_1$.
We will use the Extended Euclidean Algorithm to solve problem (2).
4. Let $r_j, u_j, v_j \in F[X]$ be the quantities computed during the j -th pass of the Extended Euclidean Algorithm for the pair (m, f) , where j is minimal such that $\deg r_j < k$. Show that (r_j, v_j) satisfy (2). What can you say about the complexity of this method?
5. **Application.** Given $2n$ consecutive terms of a recursive sequence of order n , give the recurrence. (Hint: this is where you use question 1). Illustrate your method on the Fibonacci sequence.