

WINEMAKER'S

recipe handbook



Filled with over 100 easy-to-use tested recipes and profuse with identification and selection of plants and fruit for winemaking. Contains more facts than you'll probably want to know.

TABLE OF CONTENTS

INTRODUCTION

PG 2

RECIPES

APPLE, MEDIUM	NO. 1	PG. 3
APPLE, FULL	NO. 2	PG. 3
APPLE-SPICED	NO. 3	PG. 3
APPLE CIDER	NO. 4	PG. 3
APRICOT	NO. 5	PG. 4
BANANA, DRIED & FRESH	NO. 7	PG. 4
BANANA-SPICED	NO. 8	PG. 4
BARLEY	NO. 9	PG. 5
BEET, RED	NO. 11	PG. 5
BEET, SUGAR	NO. 12	PG. 5
BLACKBERRY	NO. 13	PG. 6
BLUEBERRY	NO. 15	PG. 6
BLUEBERRY "PORT"	NO. 16	PG. 6
CARROT	NO. 17	PG. 7
CARROT "HOPS"	NO. 18	PG. 7
CARROT "BLAST"	NO. 19	PG. 7
CHERRY, SWEET	NO. 22	PG. 8
CHERRY, SOUR	NO. 23	PG. 8
CHOCHECHERRY	NO. 24	PG. 8
CHOCHECHERRY, DESSERT	NO. 25	PG. 8
COFFEE	NO. 26	PG. 9
CORN WINE	NO. 28	PG. 9
CORN "WHISKY"	NO. 29	PG. 9
CRABAPPLE	NO. 30	PG. 10
CRABAPPLE & APPLE	NO. 31	PG. 10
CRANBERRY, FRESH	NO. 32	PG. 10
CRANBERRY, JUICE	NO. 33	PG. 10
CRANAPPLE, JUICE	NO. 33	PG. 10
CURRENT, BLACK	NO. 34	PG. 11
CURRENT, RED	NO. 35	PG. 11
DANDELION	NO. 36	PG. 11
ELDERBERRY, FRESH	NO. 37	PG. 12
ELDERBERRY, DRIED	NO. 38	PG. 12
FIRETHORN (Pyranantha)	NO. 39	PG. 12
FRUIT BRANDIES	NO. 40	PG. 13
GLOGI	NO. 41	PG. 13
GOOSEBERRY	NO. 42	PG. 14
GOOSEBERRY "SPARKLE"	NO. 43	PG. 14
GRAPEFRUIT, FRESH	NO. 44	PG. 14
GRAPEFRUIT, JUICE	NO. 45	PG. 14
GRAPEWINES, Introduction	PG. 15	
VITIS VINIFERA	PG. 15	
(European & California Grapes)		
RED WINES	NO. 46	PG. 16
WHITE WINES	NO. 47	PG. 16
SECOND WINES	NO. 48	PG. 17
SECOND WINES, WHITE	NO. 49	PG. 17
THOMPSON SEEDLESS	NO. 50	PG. 17
NATIVE AMERICAN GRAPES	PG. 18	
WILD GRAPES	NO. 51	PG. 18
CONCORD, Fresh	NO. 52	PG. 19
CONCORD, Juice	NO. 53	PG. 19
CONCORD, Fresh Dessert	NO. 54	PG. 19
CONCORD, Juice Dessert	NO. 55	PG. 19
CONCORD, Frozen	NO. 56	PG. 19
WINES from		
GRAPE CONCENTRATES	NO. 57	PG. 20

INSTANT WINES, GRAPE	NO. 58	PG. 20
INSTANT WINES, FRUIT	NO. 59	PG. 20

HONEY WINES/Meads, Etc.

MEAD, DRY	NO. 60	PG. 21
MEAD, SWEET	NO. 61	PG. 21
MELOMEL	NO. 62	PG. 21
MELOMEL	NO. 63	PG. 21
PYMOMEL	NO. 64	PG. 21
METHEGLYN	NO. 65	PG. 21
JELLY & JAM	NO. 66	PG. 22
MALT "QUICKY"	NO. 67	PG. 22
MALT "LIQUEUR"	NO. 68	PG. 22
MELON	NO. 69	PG. 23
MOUNTAIN ASH	NO. 71	PG. 23
ONION	NO. 72	PG. 24
ORANGE, FRESH	NO. 73	PG. 24
ORANGE, FROZEN	NO. 74	PG. 24
OREGON "GRAPE"	NO. 75	PG. 25
PARSNIP	NO. 20	PG. 7
PEACH	NO. 6	PG. 4
PEAR	NO. 76	PG. 25
PERRY	NO. 77	PG. 25
PERSIMMON	NO. 78	PG. 26
PINEAPPLE, FRESH	NO. 79	PG. 26
PINEAPPLE, JUICE	NO. 80	PG. 26
PLUM	NO. 81	PG. 27
PLUM "PORT"	NO. 82	PG. 27
POMEGRANATE	NO. 83	PG. 27
POMEGRANATE & MALT	NO. 84	PG. 27
POTATO	NO. 85	PG. 28
POTATO "MASHER"	NO. 86	PG. 28
PRICKLY PEAR	NO. 87	PG. 28
PUMPKIN	NO. 88	PG. 29
RAISIN	NO. 89	PG. 29
RAISIN "SHERRY"	NO. 90	PG. 29
RASPBERRY	NO. 14	PG. 6
RHUBARB	NO. 91	PG. 30
RHUBARB	NO. 92	PG. 30
RICE "SAKI"	NO. 93	PG. 30
ROSE HIP, FRESH	NO. 94	PG. 31
ROSE HIP, DRIED	NO. 95	PG. 31
RUM POT	NO. 96	PG. 31
SAN GRIA	NO. 97	PG. 31
SERVICEBERRY	NO. 98	PG. 32
STRAWBERRY	NO. 99	PG. 32
TEA	NO. 27	PG. 9
TOMATO, RED	NO. 100	PG. 33
TOMATO, GREEN	NO. 101	PG. 33
TURNIP	NO. 21	PG. 7
WATERMELON	NO. 70	PG. 23
WHEAT	NO. 10	PG. 5

OTHER COMMON WINES and MAKING UP YOUR OWN RECIPES	PG. 33-34
OVER 40 references listed	

COPYRIGHT © 1976
by RAMOND MASSACcesi
All rights reserved

INTRODUCTION

what's in the handbook

Thumbing through these pages you'll find recipes that use the bounty from the open country; from the corner store; and hopefully from your own garden-- and still let the neighbors' apples tree intact. There are country wine recipes like chokecherry, rhubarb, dandelion and corn "whiskey", to name a few. For those of you that believe only grapes make wine, there is a section on grape wines-- fresh harvested, frozen, bottled grape juices, premium concentrates and making "second" wines.

Featured with the recipes are plant selections, many available from nurseries, to help turn your yard into a beautiful and practical source for jelly and winemaking. And, on your country outings use the illustrations and descriptions to help pick out and select some of the plants and fruit.

Aside from the 101 recipes given in the handbook, the last pages contain a unique recipe reference guide that adds over 35 more wine recipes to try, and even how to make up your own recipe.

winemaking equipment

Once the creative spirit of winemaking takes hold, the rewards are many. Compared to the mass produced store wines, the flavorful personally pleasing home wines open up a new world of limitless and interesting varieties. For those of you about to take your first steps into winemaking, it is strongly recommended to use the right equipment. Don't skimp-- its not worth saving a penny at the beginning and turn out less than the best results in the end. The best way to begin is with a winemaker's kit from a wine-making supply shop (avoid department store kits-- they're generally incomplete and more expensive). Most importantly, a wine-making shop can answer your questions and advise. Look for a hydrometer, primary, glass secondary, air lock and syphon. The recipes assume that you have this basic equipment. For fresh fruit winemaking, an acid tester is recommended as is a reusable nylon fermentation bag.

about the recipes

In the old days, wine was made only by the very patient under, at best, hit and miss conditions. To achieve a palatable drink, most wines were considerably sweetened or generously spiced. **THROW AWAY** those hand-me-down recipes-- We've got some updated ones of our own. The selection of recipes in this handbook is based on practicality and availability of ingredients under the reasoning that you can't make what you can't get. Cold juice extraction is used vs. boiling (leftover from jellymaking). Although the color sometimes is lighter, this method yields a fresh, fruitier flavor-- a better table wine.

Follow the recipes as given. **THEY ARE FOR ONE GALLON OF WINE.** To make more wine simply multiply the ingredients by the number of gallons desired (except for yeast-- normally one package will do for up to 5 gal.). If a little extra fermented liquid is left over after transfer, fill up a small extra glass bottle and attach lock. Use later to "top-up" the main bottle or to bottle for consumption. Recipes are in U.S.A. measures. For imperial measure, multiply ingredients by 1.25 (except yeast).

Many wines are drinkable after a few months but most will continue to improve for years. The sourness of new wines decreases-- the flavor mellows. If you wish to consume the wines young or prefer it slightly sweetened, this is done at bottling by adding 1/2 tsp. Stabilizer, and then stir in 1 cup dissolved sugar per gallon.

If the recipes are followed and the wine is allowed time it will clear naturally. Should you wish to speed up the clearing process, Sparkoloid or Bentonite can be used as fining (clearing) agent.

Although the recipes describe the recommended winemaking procedure, this is primarily meant to be recipe and general identification handbook. Consult winemaking shops and books specifically on winemaking for detailed information on the winemaking process. Some books that are highly recommended are: Making Wine, Beer and Merry; The Art of Making Wine; Progressive Winemaking; and Guidelines to Practical Winemaking-- all available from winemaking outlets. Here's to the making and the consuming. Enjoy.

1. APPLE

(Medium Body)

7-8 lb Apples or
2 qts. Juice
4½ pts Water
2 lb Sugar
2 ¼ tsp Acid Blend
½ tsp Pectic Enzyme
¼ tsp Tannin
1 tsp Nutrient
1 Campden, crush
1 pkg Champ. yeast
Later: 1 tsp anti-oxidant

Starting S.G. 1.085-90

2. APPLE

(Full Body)

14-16 lb Apples or
1 gal. Juice
1 lb Sugar
1½ tsp Acid Blend
½ tsp Pectic Enzyme
¼ tsp Tannin
1 tsp Nutrient
1 Campden, crush
1 pkg Champ. yeast
Later: 1 tsp anti-oxidant

Starting S.G. 1.085-90

3. APPLE/SPICE

("Old Fashioned")

7-8 lb Apples or
2 qts. Juice
1½ cup Chopped Light
Raisins or 4 oz.
Grape conc.
4 pts. Water
3/4 oz Cloves
½ oz Ginger Root,
mash
2½ lb Sugar
1 tsp Acid Blend
½ tsp Pectic Enzyme
½ tsp Energizer
1 Campden, crush
1 pkg Sherry yeast

HARD CIDER or **APPLE JACK** refers to a distilled apple cider, a drink developed by ingenuity and necessity. Early Americans took apple cider (about 8%) and froze it in winter to a slushy consistency. The slush (water) on top was skimmed off. What was left was alcohol or "hardened" cider. Nowadays it's double pot stillled and aged in oak 1-5 yrs. Call it Hard Cider, Apple Jack, Cider Whiskey or Apple Brandy-- it's all the same.

VARIETIES TO USE AND PLANT:

In apple wine or apple cider the more flavorful and acid varieties such as Winesap, McIntosh and Jonathans are the best. Delicious apples because of their low acidity should be combined with the more acid varieties. The ideal is a combination of the above varieties. All are available under their common name from nurseries, many in dwarf form.

METHOD:

Only apples that are sound (firm, ripe) and free of rot and worm damage should be used. Wash and sort out apples. Since a better wine is produced by fermenting on the juice (without pulp) it is recommended that the juice be pressed from the apples. You'll find that cutting the apples in smaller pieces helps extract more juice.

If you are starting with juice purchased from the store (make sure no preservatives were added) the above work is already done. In EITHER case, 1. Put the juice into the primary fermentor; immediately add the crushed Campden tablets and Pectic Enzyme to prevent browning and aid clearing. Stir in all other ingredients except Yeast. Cover Primary. 2. After 24 hrs., add yeast. Cover Primary. 3. Stir daily, check Specific gravity. When ferment reaches S.G. of 1.040 (3-5 days) syphon wine off sediment into glass jug secondary. Attach airlock. 4. When ferment is complete (S.G. has reached 1.000-- after 3 weeks) syphon off sediment into clean secondary. Reattach lock. 5. Syphon off sediment again in 2 months and again if necessary until wine is clear before bottling. Add anti-oxidant at bottling.

APPLE CIDER:

For Apple Cider, follow Apple Wine method. When ferment is complete, stir in gently ½ cup dissolved sugar per gallon of Cider. Bottle in champagne or returnable pop bottles. Apply crown caps. Age for 3 months. cool and enjoy.

NOTE: For a small quantity of wine, if a press is not available to do the job, a nylon straining bag can be used. To use: A. Place cut-up apples in bag. Press and strain out juice into primary fermentor. Immediately add Campden tablets (crushed) into juice. B. Keep all pulp in straining bag, tie top and place in primary. C. Stir in all other ingredients except yeast. Cover primary. D. After 24 hrs. strain juice from pulp. Discard pulp and proceed with step 2.

5. APRICOT

2½	Apricots
7 pts	Water
2 lb	Sugar
1¼	Acid Blend
1 tsp	Pectic Enzyme
¼ tsp	Tannin
½ tsp	Energizer
1	Campden, crush
1 pkg	Wine yeast

Starting S.G. 1.085-90

6. PEACH

2½	Peaches
7 pts	Water
2 lb	Sugar
1½	Acid Blend
1 tsp	Pectic Enzyme
¼ tsp	Tannin
½ tsp	Energizer
1	Campden, crush
1 pkg	Wine yeast

Starting S.G. 1.085-90

METHOD:

1. Wash fruit, remove stones and any brown patches, and weigh out correct amount. 2. Mash and strain out juice into primary fermentor. Keep all pulp in nylon straining bag, tie top and place in primary. 3. Stir in all other ingredients except yeast. Cover primary. 4. After 24 hrs., add yeast. Cover primary. 5. Stir daily, check S.G. and press pulp in bag lightly to help extract juice. 6. When ferment reaches S.G. 1.040 (3-5 days) strain juice lightly from bag. Syphon wine off sediment into glass jug secondary. Attach airlock. 7. When ferment is complete (S.G. has reached 1.000--about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 8. Syphon again in 2 months and again if necessary until clear before bottling.



VARIETIES

Apricots and to some extent Peach trees are not recommended in areas prone to late spring frosts. But they both make a handsome tree for shade and of course for fruit. Good all-around Apricot varieties are: Chinese (Mormon), Golden Amber, Royal, Wenatchee and Supurb. Peach varieties: July Elberta, Redhaven and Heath Cling.

7. BANANA

(Fresh or Dried)

3 lb	Bananas, fresh or 8 oz Dried
1½ cup	Chopped Light Raisins or 4 oz Grape Conc.
7 pts	Water
2¼ lb	Sugar
3 tsp	Acid Blend
¼ tsp	Tannin
1 tsp	Nutrient
1 pkg	Wine yeast

Starting S.G. 1.095

8. BANANA/SPICE

3 lb	Bananas, fresh or 8 oz Dried
8 oz	White Grape Con.
1 oz	Ginger Root (bruised)
1 oz	Cloves
7 pts	Water
2½ lb	Sugar
2½ tsp	Acid Blend
½ tsp	Tannin
½ tsp	Energizer
1 pkg	Sherry yeast

METHOD:

1. Slice bananas and skins, put into nylon straining bag and tie. In 1½ qts. water bring to boil, then, simmer for ½ hour. 2. Remove pulp, pour hot liquour over sugar in primary fermentor, and mix thoroughly. 3. Stir in remainder of water (cold) and all other ingredients. Add yeast when cool enough (85° or lower). Cover primary. 4. Stir daily, check S.G. 5. When ferment reaches S.G. 1.040 (3-5 days) strain out and syphon wine off sediment into clean glass jug secondary. Reattach airlock. 6. When ferment is complete (S.G. has reached 1.000--about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. Syphon again in 2 months and again if necessary until clear before bottling.

9. BARLEY


1 lb	Barley or 8 oz Dry Malt	¼ lb	Wheat Lt Grape Conc
1 pt	Lt. Grape Conc	½ pt	Water
6 pt	Water	2¼ lb	Sugar
3½ cup	Sugar	3 tsp	Acid Blend
¼ tsp	Tannin	¼ tsp	Nutrient
2 tsp	Acid Blend	¼ tsp	Tannin
¼ tsp	Nutrient	1	Campden, crush
1	Campden, crush	1 pkg	Wine yeast
1 pkg	Wine yeast		

Starting S.G. 1.085-90

METHOD:

1. Wash grain, soak 24 hrs. in 1 qt. water. Crush again. Pour grain and water through straining bag over primary fermentor, tie bag and leave in primary. If dry malt is substituted for barley, stir and dissolve in 1 qt. hot water. 2. Mix in all ingredients together in primary EXCEPT yeast. Cover primary. 3. After 24 hrs., add yeast. Cover primary. 4. Stir daily and check S.G. 5. When ferment reaches S.G. 1.040 (3-5 days) strain juice lightly from bag. Syphon wine off sediment into glass jug secondary. Attach airlock. 6. When ferment is complete (S.G. has reached 1.000--about 3 weeks) syphon off sediment into clear secondary. Reattach lock. 7. Syphon again in 2 months and again if necessary until clear before bottling.

You may care to sweeten this wine. Before bottling add ½ tsp. Stabilizer, then ¼ lb. dissolved sugar per gallon.



10. WHEAT

¼ lb	Wheat Lt Grape Conc
½ pt	Water
6½ pt	Sugar
3 tsp	Acid Blend
¼ tsp	Nutrient
¼ tsp	Tannin
1	Campden, crush
1 pkg	Wine yeast

11. BEET

2½ lb	Beets
1 gal	Water
2¼ lb	Sugar
2 tsp	Acid Blend
1 tsp	Nutrient
¼ tsp	Tannin
1	Campden, crush
1 pkg	Wine yeast

Starting S.G. 1.095 - 1.100

METHOD:

1. Wash beets, skin the beets and cut up into small pieces. 2. Place in nylon straining bag, tie top and gently boil in 2 qts. of the water until they are tender. 3. Pour hot liquor over sugar in primary fermentor, and mix thoroughly. 4. Put bag with pulp in fermentor, stir in remainder of water (cold) and all other ingredients EXCEPT yeast. Cover primary. 5. After 24 hrs., add yeast. Cover primary. 6. Stir daily, check S.G. and press pulp lightly to help extraction. 7. When ferment reaches S.G. 1.040 (3-5days) strain juice lightly from bag. Syphon wine off sediment into glass jug secondary. Attach airlock. 8. When ferment is complete (S.G. has dropped to 1.000-- about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 9. To aid clearing syphon again in 2 months and again if necessary before bottling.

This wine may be more to your liking slightly sweetened. At bottling add ½ tsp. Stabilizer, then ¼ lb. dissolved sugar per gallon.



12. BEET, SUGAR

2½ lb	Sugar Beets
¼ pt	Grape Conc. or ½ lb Raisins
1 gal	Water
1 ¼ lb	Sugar
¼ tsp	Tannin
2 tsp	Acid Blend
1	Campden, crush
1 pkg	Wine yeast

13. BLACKBERRY 14. RASPBERRY

4 lb	Blackberries (fresh or frozen)	3 lb	Raspberries (fresh or frozen)
7 pts	Water	7 pts	Water
2 1/4 lb	Sugar	2 1/4	Sugar
1/2 tsp	Acid Blend	1/2 tsp	Acid Blend
1/2 tsp	Pectic Enzyme	1/2 tsp	Pectic Enzyme
1 tsp	Nutrient	1 tsp	Nutrient
1	Campden, crush	1	Campden, crush
1 pkg	Wine yeast	1 pkg	Wine yeast

Starting S.G. 1.090-95

METHOD:

Pick berries when they are fully ripened but not moldy. Remove any stems, leaves, and foreign matter (bugs). 1. Wash and drain the berries using nylon straining bag (or press), mash and strain out juice into primary fermentor. Keeping all pulp in straining bag, tie top and place in primary. 2. Stir in all other ingredients EXCEPT yeast. Cover primary. 3. After 24 hrs., add yeast. Cover primary. 4. Stir daily, check hydrometer reading (S.G.) and press pulp lightly to aid extraction. 5. When ferment reaches S.G. 1.030 (about 5 days) strain juice from bag. Syphon wine off sediment into glass secondary. Attach airlock. 6. When ferment is complete (S.G. has dropped to 1.000—about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. To aid clearing, syphon again in 2 months and again if necessary before bottling.

Makes a nice dry wine; but for earlier consumption or for slightly sweetened wine, at bottling: Add 1/2 tsp. Stabilizer, then, stir in 1/4 lb. dissolved sugar per gallon.

VARIETIES

From same family (*Rubus* L.), 1/2"-3/4" red or black fruit. Blackberry take core with, Raspberry don't.

In same family are Dewberry, Loganberry and Boysenberry. Varieties to try: Marion, Ebony King and wild Himalaya Blackberries. Raspberries: Latham, Black Hawk, Sumner and Fairview.



15. BLUEBERRY 16. BLUEBERRY PORT

(fresh or frozen)

2 lb	Blueberries	6 lb	Blueberries
1/2 pt	Red grape conc (for later)	1/2 pt	Lt Dry Malt Water
7 pts	Water	4 pts	Water
1 1/4 lb	Sugar	1 1/4 lb	Sugar
1 1/2 tsp	Acid Blend	1/2 tsp	Acid Blend
1/2 tsp	Pectic Enzyme	1/2 tsp	Pectic Enzyme
1/2 tsp	Energizer	1/2 tsp	Energizer
1	Campden, crush	1	Campden, crush
1 pkg	Wine yeast	1 pkg	Sherry or Port yeast

METHOD:

Use ripe but mold-free berries. Remove stems, leaves and foreign matter. 1. Wash and drain berries. Using nylon straining bag (or press) mash and strain out juice into primary fermentor. Keeping all pulp in straining bag, tie top and place in primary. 2. Stir in all other ingredients EXCEPT yeast. NOTE: In recipe 15, the Red Grape Concentrate is added later to finished wine prior to bottling. Cover primary. 3. After 24 hrs., add yeast. Cover primary. 4. Stir daily, check S.G. and press pulp lightly to aid extraction. 5. When ferment reaches S.G. 1.030 (about 5-8 days), strain juice from bag. Syphon wine off sediment into glass secondary. Attach airlock. 6. When ferment is complete (S.G. has dropped to 1.000—about 4 weeks) syphon off sediment into clean secondary. Reattach lock. 7. To aid clearing syphon again in 2 months and again if necessary before bottling.

In recipe 15 at bottling add 1/2 tsp Stabilizer, then, stir in Red Grape concentrate. This can be done with recipe 16 also for fuller taste or substitute with 1/4 lb. dissolved sugar per gallon.



VARIETIES

Blueberries (*Vaccinium* L.) do best in moist, acid, easy draining conditions. Besides the fruit they yield the plants make a handsome addition to the yard. 1/4"-3/4" round, blue-black berries are in normally loose clusters. Bilberries (see pg. 33-34) are from same family. True Huckleberries (*Gaylussacia* L.) are not in this family--their leaves and fruit characteristics are different. Blueberries to plant: "Earliblue", "Coville", "Bluecrop" and "Berkley".

17. CARROT

3½ lb	Carrots
½ pt	Grape Conc. or 1 lb raisins
1 gal	Water
2 lb	Sugar
3 tsp	Acid Blend
¼ tsp	Tannin
½ tsp	Energizer
1	Campden, crush
1 pkg	Wine yeast

18. CARROT "HOP"

3½	Carrots
½ oz	Dried Hops
1 gal	Water
2½ lb	Brown Sugar
2 tsp	Acid Blend
½ tsp	Energizer
1	Campden, crush
1 pkg	Sherry yeast

19. CARROT BLAST

4¼ lb	Carrots
½ pt	Grape Conc.
1 cup	Dry Malt
1 gal	Water
2¼ lb	Sugar
2¼ tsp	Acid Blend
½ tsp	Energizer
1	Campden, crush
1 pkg	Sherry yeast

20. PARSNIP

3 lb	Parsnips
½ pt	Grape Conc.
1 gal	Water
1¼ lb	Sugar
2½ tsp	Acid Blend
¼ tsp	Tannin
½ tsp	Energizer
1	Campden, crush
1 pkg	Wine yeast

METHOD:

1. Scrub clean, remove tops and rootlings. Chop into pieces, put into nylon straining bag and tie. Boil in the 1 gallon of water until tender, strain out (don't press) juice. 2. Remove pulp, pour hot liquor over sugar in primary fermentor, and mix thoroughly. 3. Stir in remainder of water (cold) and all other ingredients EXCEPT yeast. If hops or raisins are used, put in tied straining bag. Cover primary. 4. After 24 hrs., add yeast. Cover primary. 5. Stir daily and check S.G. 6. When ferment reaches S.G. 1.040 (3-5 days), discard pulp, syphon wine off sediment into glass jug secondary. Attach airlock. 7. When ferment is complete (S.G. has dropped to 1.000--about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 8. To aid clearing syphon again in 2 months and again if necessary before bottling.

21. TURNIP

4 lb	Turnips
½ oz	Dried Hops
1 gal	Water
2½ lb	Sugar
1 tsp	Acid Blend
¼ tsp	Tannin
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine yeast



23. CHERRY, Sour 12. CHERRY, Sweet

(Sweet or Tart)

3 lb	Sour Cherry or	6 lb	Mixed Cherries
4 lb	Dk Sw. Cherry	½ pt	Grape Conc.
7 pts	Water	7 pts	Water
2¼ lb	Sugar	2¼ lb	Sugar
1 tsp	Acid Blend	¼ tsp	Pectic Enzyme
½ tsp	Pectic Enzyme	1 tsp	Nutrient
1 tsp	Nutrient	1	Campden, crush
1	Campden, crush	1 pkg	Sherry or Port yeast
1 pkg	Wine yeast		

Starting S.G. 1.090

Starting S.G. 1.100 - 1.110

METHOD:

1. Wash, remove stems and leaves, and preferably remove pits (use a pitter--either mechanical or a couple of extra hands from around the house). In any case be careful not to break the bitter pits. 2. Using nylon straining bag mash and squeeze out juice into primary fermentor. Keeping all pulp in straining bag, tie top and place in primary. 3. Stir in all other ingredients EXCEPT yeast. Cover primary. 4. After 24 hrs., add yeast. Cover primary. 5. Stir daily, check S.G. and press pulp lightly to aid extraction. 6. When ferment reaches S.G. 1.030 (about 5 days) strain juice from bag. Syphon wine off sediment into glass jug secondary. Attach airlock. 7. When ferment is complete (S.G. has dropped to 1.000--about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 8. To aid clearing syphon again in 2 months and again if necessary before bottling.

VARIETIES

Most cherry trees do not adapt well to a mild winter climate. The handsome trees, the crown of flowers in spring, and the fruit-- whether sour, sweet or crosses-- make them highly desirable. Sour varieties: Montmorency, North Star, Early Richmond and Meteor. Sweet varieties: Bing, Royal Ann, Van and Lambert.



24. CHOKECHERRY 25. CHOKECHERRY

(Dinner Wine)

(Dessert Wine)

2½ lb	Choke Cherry	8 lb	Choke Cherry
½ pt	Red grape conc. or 1 lb raisins	7 pts	Water
7 pts	Water	2¼ lb	Sugar
2¼ lb	Sugar	1 tsp	Pectic Enzyme
½ tsp	Acid Blend	½ tsp	Energizer
½ tsp	Pectic Enzyme	1	Campden, crush
1 tsp	Nutrient	1 pkg	Sherry yeast
1	Campden, crush		
1 pkg	Wine yeast		

Starting S.G. 1.090-95

METHOD:

Pick fruit when they are fully ripe black-purple if you can fight off the birds and other people. 1. Wash and remove stems, leaves, and badly bruised fruit. 2. The pits are too large to remove, so mash cherries without breaking the bitter pits. Using nylon straining bag, mash and squeeze out juice into primary fermentor. Keeping all pulp in straining bag, tie top and place in primary. 3. Stir in all other ingredients EXCEPT yeast. Cover primary. 4. After 24 hrs., add yeast. Cover primary. 5. Stir daily, check S.G. and press pulp lightly to aid in extraction. 6. When ferment reaches S.G. 1.030 (about 5 days) strain juice from bag. Syphon wine off sediment into glass jug secondary. Attach airlock. 7. When ferment is complete (S.G. has dropped to 1.000-- about 3 weeks; NOTE: Recipe No. 25 may finish higher) syphon off sediment into clean secondary. Reattach lock. 8. To aid clearing syphon again in 2 months and again if necessary before bottling.

Here is a dinner wine (recipe 24) that can be left dry or slightly sweetened. Before bottling add ½ tsp. Stabilizer, then, ¼ lb. dissolved sugar per gallon.

VARIETIES

Chokecherries (*Aronia* L.) are 3'-8" shrubs with small white flowers and ¼" round, red to black fruit ripening Aug.-Nov. Red fall color. Black (*A. melanocarpa*), Red (*A. arbutifolia*), and Purple (*A. prunifolia*). Chokecherries are found wild and in nurseries. Shubert Chokecherry (*Prunus virginiana*) is from cherry family (*Prunus* L.) so use sour cherry recipe.



26. COFFEE

5 Tbs	Instant Coffee
7 pts	Water
3 lb	Honey
3 tsp	Acid Blend
¼ tsp	Energizer
1	Campden, crush
1 pkg	Sherry yeast

Starting S.G. 1.095-1.105

27. TEA

5 Tbs	Instant Tea
½ pt	Whit grape conc.
7 pts	Water
2½ lb	Sugar
1 tsp	Acid Blend
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine yeast

28. CORN

1½ lb	Corn
7 pts	Water
2 lb	Honey
3 tsp	Acid Blend
½ tsp	Energizer
1	Campden, crush
1 pkg	Wine yeast

Starting S.G. 1.090-95

29. CORN WHISKEY

2 lb	Corn
½ pt	Whit grape conc.
3	Oranges
12	Peppercorns
1 gal	Water
3½ lb	Sugar
3 tsp	Acid Blend
1 tsp	Energizer
1	Campden, crush
1 pkg	Sherry yeast

Starting S.G. 1.110 - 120

METHOD:

1. Measure out instant flavor and sugar (or honey) into primary fermentor. Pour water (hot) over and stir thoroughly. 2. Stir in all other ingredients EXCEPT yeast. Cover primary. 4. Stir daily and check S.G. 5. When ferment reaches S.G. 1.040 (3-5 days) syphon wine off sediment into glass jug secondary. Attach airlock. 6. When ferment is complete (S.G. has reached 1.000-- about 3 weeks) syphon off sediment into clean secondary. Reattach lock. Syphon again in 2 months and again if necessary until clear before bottling.

METHOD:

Both pleasant wines, the "whisky" wine is a 'sit-down wine' -- you have to sit down to drink this. 1. Wash off ears, cut off and weigh out kernels. Using nylon straining bag, crush and squeeze out juice of kernels into primary fermentor. If used, grate rind of oranges and add peppercorns into bag. Squeeze out juice from orange (discard pulp). With corn pulp in bag, tie top and place in primary. 2. Stir in all other ingredients EXCEPT yeast. Cover primary. 3. After 24hrs., add yeast. Cover primary. 4. Stir daily, check S.G. and press pulp lightly to aid extraction. 5. When ferment reaches 1.040 (about 3-5 days) strain juice lightly from bag. Syphon wine off sediment into glass jug secondary. Attach airlock. 6. When ferment is complete (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. To aid clearing syphon again in 2 months and again if necessary before bottling.



30. CRABAPPLE 31. CRABAPPLE Plus

4½ lb	Crabapples	3½ lb	Crabapples
½ pt	Wht grape conc. or 1 lb Lt raisins	1 qt	Apple juice
7 pts	Water	6 pts	Water
1¼ lb	Sugar	1½	Sugar
½ tsp	Pectic Enzyme	½ tsp	Pectic Enzyme
1 tsp	Nutrient	1 tsp	Nutrient
1	Campden, crush	1	Campden, crush
1 pkg	Champ. yeast	1 pkg	Champ. yeast

Starting S.G. 1.090-95

METHOD:

Select only sound and ripe fruit. Wash lightly and chop up apples, but avoid cutting the bitter seeds. If a press is not available, place cut-up fruit in nylon straining bag. Crush, press and strain out juice into primary fermentor. Immediately add Campden (crushed) into juice. 2. Keep all pulp in bag, tie top and place in primary. 3. Stir in all other ingredients EXCEPT yeast. Cover primary. 4. After 24 hrs., strain juice from pulp. Discard pulp. Add yeast. Cover primary. 5. Stir daily and check S.G. 6. When ferment reaches S.G. of 1.040 (3-5 days) syphon wine off sediment into glass jug secondary. Attach airlock. 7. When ferment is complete (S.G. has dropped to 1.000-- about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 8. Syphon again in 2 months and again if necessary until clear before bottling.

VARIETIES

Crabapples (*Malus* L.) are hardy trees, easy to care for, excellent flowering landscape addition. For jelly and winemaking the smaller apple-like fruit are very welcome. Good varieties to plant: "Dolgo", white blossoms, good flavor; "Hopa", rose-red blossoms; "Royalty", purple foliage, maroon flowers.



32. CRANBERRY 33. CRANBERRY or Cranapple Juice

(Fresh)

3 lb	Cranberries	2 qts	Juice
½ pt	Red grape conc. or 1 lb Dk raisins	2 qts	Water
7 pts	Water	1¼ lb	Sugar
2½ lb	Sugar	2½ tsp	Acid Blend
½ tsp	Pectic Enzyme	½ tsp	Pectic Enzyme
½ tsp	Energizer	½ tsp	Energizer
1	Campden, crush	1	Campden, crush
1 pkg	Wine yeast	1 pkg	Wine yeast

Starting S.G. 1.090-95

METHOD:

Use only well-ripened but sound fruit. 1. Wash and place fruit in nylon straining bag. Press, crush and strain out juice into primary fermentor. Keeping all pulp in bag (add chopped raisins, if used) tie top and place in primary. If using juice all this work is already done. Proceed to step 2. 2. Stir in all other ingredients EXCEPT yeast. Cover primary. 3. After 24 hrs., add yeast. Cover primary. 4. Stir daily, check S.G. and press pulp lightly to aid extraction. 5. When ferment reaches 1.030 (about 5 days) strain juice lightly from bag. Syphon wine off sediment into glass jug secondary. Attach airlock. 6. When ferment is complete (S.G. has dropped to 1.000--about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. Syphon again in 2 months and again if necessary until clear before bottling.



VARIETIES

Cranberry (*Vaccinium* L.) related to blueberry group. The commercial variety American or Large Cranberry (*V. macrocarpum*) yield ¾"-1" round, red fruit. They require a stream with some marsh suitable for a bog - for flooding but quick emptying. Unless you can find these wild, seek them in the store. Lingonberry is a wild variety.

34. CURRANT, BLACK

35. CURRANT, RED

36. DANDELION

(Black - Dry Wine)

(Red - Sweeter)

2½ lb	Black Currants	3½ lb	Red Currants
7 pts	Water	7 pts	Water
2¼ lb	Sugar	3 lb	Sugar
½ tsp	Pectic Enzyme	½ tsp	Pectic Enzyme
1 tsp	Nutrient	1 tsp	Nutrient
1	Camden, crush	1	Camden, crush
1 pkg	Wine yeast	1 pkg	Sherry or Port yeast

7 cups	Dandelion petals
½ pt	Wht grape conc. or 1 lb Lt raisins
1 gal	Hot Water
2 lb	Sugar
3 tsp	Acid Blend
¼ tsp	Tannin
½ tsp	Energizer
1	Camden, crush
1 pkg	Wine yeast

METHOD:

Strip from stems, sort out blemished or moldy fruit, pieces of stem and leaves. Pick fully ripe. 1. Wash and drain currants. Using nylon straining bag (or with press) mash and strain out juice into primary fermentor. Keeping all pulp in bag, tie top and place in primary. 2. Stir in all other ingredients EXCEPT yeast. Cover primary. 3. After 24 hrs., add yeast. Cover primary. 4. Stir daily, check S.G. and press pulp lightly to aid extraction. 5. When ferment drops to S.G. 1.030 (about 5 days) strain juice from bag. Syphon wine off sediment into glass jug secondary. Attach airlock. 6. When ferment is complete (S.G. has reached 1.000-- about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. To aid clearing syphon again in 2 months and again if necessary before bottling.

Red currant can be substituted for black currant and visa-versa in two above recipes. Although recipe 35 produces sweeter wine, wines can be further sweetened to taste by: at bottling add ½ tsp. Stabilizer, then, ¼ lb. dissolved sugar per gallon.

VARIETIES

Currants and Gooseberries (*Ribes* L.) are closely related. Normally currants have smooth canes and gooseberries with spiny canes. Many are decorative, 3'-5' shrubs with attractive foliage. Currants to try: Red Lake, Perfection, Cherry and Fay's Prolific. Gooseberries: Pixwell or Oregon Champion. They do host pine blister rust, so consult nursery.



METHOD:

Cut off and use yellow petals only when flowers are fully opened. Avoid any sprayed plants and discard all green parts as they impart bitterness to the wine. 1. Wash and drain petals. Put petals in bag (add in chopped raisins, if used) tie top and place in primary. 2. Pour over hot water, stir in sugar, and add in all other ingredients EXCEPT yeast. Cover primary. 3. After 24 hrs., add yeast. Cover primary. 4. Stir daily, check S.G., and press pulp lightly to acid extraction. 5. When ferment reaches 1.040 (3-5 days) strain pulp lightly from bag. Syphon wine off sediment into glass jug secondary. Attach airlock. 6. When ferment is complete (S.G. has dropped to 1.000-- about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. Syphon again in 2 months and again if necessary until clear before bottling.

Starting S.G. 1.090-95

37. ELDERBERRY

(Fresh)

3 lb	Elderberries
7 pts	Water
2 1/4 lb	Sugar
1 tsp	Acid Blend
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine yeast

Starting S.G. 1.095-1.100

METHOD:

These highly colored blue-black berries take a little effort in picking but the resulting "Chianti" type wine is very popular. They also make an excellent "Port" type wine by increasing fruit to 4 1/2 lbs., add another 1/4 lb. sugar and use Sherry or Port yeast. 1. Strip from stems, when fully ripe. Wash, sort out blemished and moldy berries, and drain. Using nylon straining bag (or with press) mash and strain out juice into primary fermentor. Keeping all pulp in bag (with dried berries put in with chopped raisins), tie top, and place in primary. 2. Stir in all other ingredients EXCEPT yeast. Cover primary. 3. After 24 hrs., add yeast. Cover primary. 4. Stir daily, check S.G., and press pulp lightly to aid juice extraction. 5. When ferment reaches 1.030 (about 5 days) strain juice lightly from bag. Syphon wine off sediment into glass secondary. Attach airlock. 6. When ferment is complete (S.G. has dropped to 1.000-- about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. To aid clearing syphon again in 2 months and again if necessary before bottling.

Using 1/4 oz. of oak chips per gallon of wine will benefit this wine with an "aged-in-oak" flavor.



VARIETIES

American Elder (*Sambucus Canadensis*) is the most popular. These fast growing, wild looking shrubs yield cluster of black 1/4" round berries perfect for pies, jellies and wines. In the wild and some nurseries also: Red-berried Elder (*S. racemose*) and Blue Elder (*S. glauca*).

38. ELDERBERRY

(Dried)

5 oz	Dried Elderberry
1/2 pt	Red grape conc. or 1 lb Dk raisins
1 gal	Water
2 1/4 lb	Sugar
1 1/2 tsp	Acid Blend
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine yeast

39. FIRETHORN

(*Pyracantha*)

2 qts	Berries
1/3 pt	Grape Conc. or 1/4 lb Dk raisins
6 pts	Water
2 1/2 lb	Sugar
1/2 tsp	Acid Blend
1/2 tsp	Pectic Enzyme
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine yeast

METHOD:

Pick these fire-red berries when fully ripe. Wash, sort out blemished berries, and drain. 1. Using nylon straining bag (or with press) mash and strain out juice into primary fermentor. Keeping all pulp in bag (add in chopped raisins, if used), tie top, and place in primary. 2. Stir in all other ingredients EXCEPT yeast. Cover primary. 3. After 24 hrs., add yeast. Cover primary. 4. Stir daily, check S.G., and press pulp lightly to aid juice extraction. 5. When ferment reaches 1.030 (about 5 days) strain juice lightly from bag. Syphon wine off sediment into glass secondary. Attach airlock. 6. When ferment is complete (S.G. has dropped to 1.000-- about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. To aid clearing syphon again in 2 months and again if necessary before bottling.



VARIETIES

Popularly called "Firethorn" the botanical name is *Pyracantha* L. Grown widely for bright fruits, evergreen foliage, variety of uses and easy growing. Fruits vary in color (normally red to orange), size, and season. "Wyatti", "Lelandi" are hardiest.

40. FRUIT BRANDIES

Fresh fruit, vodka, sugar and time equals a number of deliciously warming brandies. As a drink or poured over desserts the results are rewarding. This is done without fermentation.

Peach Brandy

2½ lbs	Fresh, Ripe Peaches (Weigh without stones)
1½ cups	Sugar
½ cup	Honey
1.5 btl	Vodka (80 Proof)

Apricot Brandy

2 lbs	Fresh, Ripe Apricots (Weigh without stones)
1½ cups	Sugar
½ cup	Honey
1.5 btl	Vodka (80 Proof)

Cherry Brandy

4 cups	Fresh, pitted Cherries (Sweet or Tart)
1½ cups	Sugar
1.5 btl	Vodka (80 Proof)

METHOD:

1. Select fully ripened fruit. Wash and remove stones or pits as these impart a bitter aftertaste.
2. Drain, cut up and mash fruit to facilitate juicing into a food grade plastic container. Add and thoroughly mix in sugar (and honey). Cover container with plastic sheet (securely tie). Let this sit for 24 hrs.
3. Next pour and mix in the 1/5 of Vodka. Recover container. Let this sit for about 4-6 months and allow fruit and Vodka to blend.
4. Then, strain liquor free of pulp into bottles. It's ready!

And don't throw away the fruit pulp. After sitting all that time in alcohol it's bound to be just great on ice creams and a number of other desserts.

NOTE: Brandy can be substituted for Vodka in above recipes.

41. GLOGG

A hot cup of glogg is the Swedish way of offering a warm welcome to a friend on a wintry evening.

Swedish Glogg: Serves 10-15 depending on how cold the weather.

1 cup	Blanched Almonds
12	Cardamom Seeds, peeled
12	Cloves, whole
6	2½" Cinnamon Sticks
6	Candied Oranges or Peel of 1 Orange
1 btl	Burgundy or Claret (26 oz)
1 cup	Dark Raisins
1 btl	Brandy or Aquavit (26 oz)
1 cup	Cube Sugar, crush

Put cardamom seeds, cinnamon, cloves and orange parts in nylon bag and tie top. In large saucepan put bag, ½ bottle of the wine, and the raisins. Cover and simmer for 15 minutes. Add remaining wine and the almond; heat through. In another saucepan warm about 1/3 of the brandy (or aquavit); pour in sugar and light with match to flame. When sugar has melted, pour in remainder of brandy. If necessary extinguish flame with cover lid. Add sugar mixture to wine mixture. Pour in cup with raisins and almonds in each serving.

Instant Glogg:

1 btl	Red wine (26oz)	¼ cup	Dark raisins
¼ pt	Gin	10	Cloves
2½ oz	Blanched almonds	1	Cinnamon stick

Mix and heat (don't boil) all ingredients nuts and raisins, soak for 15-20 min. and serve. Or, when cool bottle and serve later, heated

42. GOOSEBERRY

2½ lb	Gooseberries
7 pts	Water
2¼ lb	Sugar
½ tsp	Pectic Enzyme
1 tsp	Nutrient
1	Camden, crush
1 pkg	Wine yeast

43. "SPARKLE"

2½ lb	Gooseberries
7 pts	Water
1¼ lb	Sugar
½ tsp	Pectic Enzyme
1 tsp	Nutrient
1	Camden, crush
1 pkg	Wine yeast

Starting S.G. 1.080-85

METHOD:

Use firm ripe berries for the still wine and slightly greener berries for the "Sparkle" wine. Wash, sort out blemished or moldy berries, and drain water. 1. The juice is hard to press. Using nylon straining bag mash and strain out juice into primary fermentor. Keeping all pulp in bag, tie top, and place in primary. 2. Stir in all other ingredients EXCEPT yeast. Cover primary. 3. After 24 hrs., add yeast. Cover primary. 4. Stir daily, check S.G., strain pulp lightly to aid juice extraction. 5. When ferment drops to 1.030 (about 5 days) strain juice lightly from bag. Syphon wine off sediment into glass secondary. Attach airlock. 6. When ferment is complete (s. G. has dropped to 1.000-- about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. Syphon again in 2 months and again if necessary until clear before bottling.

For the Gooseberry "Sparkle", when wine is clear and stable: A. Syphon wine into primary fermentor. B. Stir in 2 oz. dissolved sugar per gallon of wine and 1 package Champagne Yeast (for 1-5 gal.) C. As you syphon into Champagne or pop bottles keep stirring. D. Store upright at 65-70° for 2-3 months. E. Check by chilling bottle, then opening.



VARIETIES

Gooseberries come from same family (Ribes L.) as Currants. See varieties under "Currants" pg. 11

44. GRAPEFRUIT

(Fresh)

6	Grapefruit
6 pts	Water
2¼ lb	Sugar
¼ tsp	Tannin
1	Camden, crush
1 pkg	Champ. Yeast

45. GRAPEFRUIT

(Juice)

1½ pt	Grapefruit Juice (unsweetened)
½ pt	Wht grape conc.
6 pt	Water
1 ¼ lb	Sugar
¼ tsp	Tannin
½ tsp	Pectic Enzyme
½ tsp	Energizer
1	Camden, crush
1 pkg	Champ. Yeast

METHOD:

Grate rind from 2 grapefruit (exclude any bitter white pith). Peel fruit and separate segments. 1. Using nylon straining bag (fine mesh) squeeze juice into primary fermentor. Keeping pulp in and grated rind in bag, tie top and place in primary. If using only juice, pour in primary and proceed. 2. Add sugar, pour water (warm) over and stir in all other ingredients EXCEPT yeast. Cover primary. 4. Stir daily, check S.G. and press pulp lightly to aid extraction. 5. When ferment reaches S.G. 1.040 (3-5 days) lightly strain juice from bag. Syphon wine off sediment into glass secondary. Attach airlock. 6. When ferment is complete (S.G. has dropped to 1.000-- about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. To aid clearing syphon again in 2 months and again if necessary before bottling.

Sweeten these wines at bottling as follows: Add ½ tsp. Stabilizer, then, add ¼-½ lb. dissolved sugar per gallon

GRAPE WINES: An Introduction

We all know dandelions, parsnips, elderberries and many other "country" materials produce wine. But, for many, "wine" comes only from the fermentation of grapes. History traces wine and the grape vine as one. If wine were only made from grapes the grape wines would happily fill the void. Commercially they are the most popular.

The skins, pulp and to a limited extent the seeds from different grape varieties produce this wide selection. But, very importantly, so does the way chosen to make the wine. White grape wines are made from the pressed juice of the grape only. It doesn't matter if it's a red or white skinned grape; as long as the pulp is white the wine can be white. Red wines on the other hand are fermented with the skins, pulp and juice. The skins provide the essential difference between white and red wines. They give color, tannins-- that mouth drying taste, and other flavoring constituents. Red wine is made from red-skinned grapes. This difference sometimes makes white wines and red wines seem like worlds apart. Rose' wine fits in there somewhere. Properly made, this wine results from pressing the fermenting grape juice from the pulp before the color becomes a deeper red.

Many people have grapes growing in their backyards. Some are found growing wild in the countryside just waiting to be picked. There are some available in the super market (although these are table eating grapes not wine grapes). Generally in September and October the major harvest of grapes takes place in the northern hemisphere. In some cities winemaking supply shops and produce dealers bring in these fresh-harvested grapes for use by home wine makers. Also, premium wine grapes are made into concentrates and available on a year-around basis. These concentrates make available to us high quality grape varieties from around the world which normally would be unobtainable.

From the recipe viewpoint this book has divided the grape varieties into 2 general groups: 1. European species (*Vitis Vinifera*) grown here. 2. Native American species.

Red grape wines are more comfortable toward the dry side, whereas whites vary from tart to very rich.

The balancing of acidity against sugar from the maturing of the grape to the making of the wine is a factor in determining if a wine is flat and dull or too tart to drink. Here again depending on variety, cool or warm climate and when picked, two major considerations can be tested. Use the acid tester to determine acid content. Because these 2 prime taste factors can be measure easily, here is a guide to the recommended readings at the start of fermentation:

	Specific Gravity	Acid Level (% Tartaric)
White Table Wines	1.080-1.095	.70%
Red Table Wines	1.085-1.100	.65%
Sweet Table Wines	1.090-1.105	.65%
Dessert Wines	1.095-1.110	.50%

Individual varieties may call for exceptions to the general guide above. The starting mixture (must) is adjusted with the addition of sugar and/or fruit acids or to lower high reading by diluting with water. Take readings from samples of pressed juice clear of skins or debris.

VITIS VINIFERA

Of the 32 species which contains over 8000 named grape vines around the world, *Vitis Vinifera* is by all standards the most important. It produces a variety of grapes that accounts for most of the wines produced in Europe and California. The ideal material for winemaking is one that yields a juice with enough sugar, enough acid to add tartness, and other taste aroma factors to give the wine balance with character.

The *Vitis Vinifera* species of grape has many varieties that do just that. Most of these varieties can be made into wine from the pulp and juice only. The normal ingredients to assure a proper ferment and in a few cases very minor corrections of sugar and acid levels are all that's needed.

46. RED WINES

RED TABLE WINE (Vitis Vinifera)

16-18 lb Grapes (depending on variety)

Sugar, add if low to S.G. 1.090-1.100

Acid Blend, add if low to .65% tartaric

1 Campden, crush

1 pkg Wine Yeast

METHOD:

Use grapes which are fully ripened and free of mold. Remove from stems. Discard leaves and any foreign matter. 1. For small quantities (5-10 gal) crush by hand into primary fermentor. For larger quantities use grape crusher. DO NOT fill primary to top. Allow at least 1/5 of primary for grape must to expand. 2. Test sample of clear juice and adjust acid and sugar content, if necessary. 3. Stir in campden. Cover primary. After 12 hrs. add yeast. Recover primary. 4. Pulp raises to form a "cap". Stir daily, keep cap punched down and wet and check S.G. 5. When ferment reaches S.G. 1.030 (about 5-6 days) syphon free-run juice into secondary. Strain pulp (or use press) and add this juice to secondary. 6. When ferment is complete (S.G. has dropped to 1.000--about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. To aid clearing syphon again in 2 months and again if necessary before bottling.

Red grape wines can benefit from the judicious use of aging in oak barrels or even oak chips.

47. WHITE WINES

WHITE TABLE WINE (Vitis Vinifera)

16-18 lb Grapes (depending on variety)

Sugar, add if low to S.G. 1.085-1.095

Acid Blend, add if low to .70% tartaric


1 tsp Nutrient

1 Campden, crush

1 pkg Wine Yeast

METHOD:

Use grapes which are fully ripened and free of mold. Remove from stems. Discard leaves and any foreign matter. 1. Crush grapes into extra primary. Quickly stir in Campden. 2. Immediately press out juice with press (with small quantities nylon mesh bag can be used) into another primary fermentor. 3. Adjust sugar and acid levels of juice and nutrient. Cover primary. 4. After 12 hours add yeast. Cover primary. Ferment at 70°F or lower. 5. When ferment reaches 1.040 (3-5 days) syphon off sediment into clean glass secondary. Attach airlock. 6. When ferment is complete (S.G. has dropped to 1.000--about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. Syphon again in 2 months and again if necessary until clear before bottling.



48. SECOND WINES, Red

With the European and California grape varieties it is possible to make what is called a very acceptable "second wine" using the grape pulp. It is a way of doubling the amount of wine made and cutting costs per gallon. Aside from putting water, sugar, fruit acid and nutrient back into the pulp grape concentrate is added to provide body and vinous character.

RED 2nd WINE

After wine has fermented down to S.G. 1.030 use pulp (unpressed). Measure amount of wine juice obtained in first wine. For every gallon measured add to pulp the following:

1 gal	Water
8-10 oz	Red Grape Concentrate
1 1/2 lb	Sugar
2 tsp	Acid Blend
1/8 tsp	Tannin
1/2 tsp	Nutrient

Use yeast already present in pulp. Hydrometer will indicate lower than expected reading because of alcohol already in pulp. Follow recipe. When ferment is down to S.G. 1.010 syphon juice to secondary. Press Juice from pulp and add. Proceed to clear.

49. SECOND WINES, White

After grapes have been pressed for first wine use pulp for "second wine". Since they are fermented at same time plan equipment use accordingly. For every gallon of juice extracted add to pulp the following:

1 gal	Water
8-10 oz	Grape Concentrate
1 1/2 lb	Sugar
2 tsp	Acid Blend
1/8 tsp	Tannin
1/2 tsp	Nutrient
1 pkg	Wine Yeast

METHOD:

1. Ferment in primary down to S.G. 1.020, then syphon secondary. Press out additional juice from pulp if desired. 2. Allow to clear as per first run white wine.

50. Thompson Seedless

16 lb	Grapes
1/2-3/4 cup	Sugar
2 tsp	Acid Blend
1/4 tsp	Tannin
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine Yeast

METHOD:


From the *Vitis Vinifera* species this is really an eating grape. It is generally lower in sugar and acid than the normal table wine grape. 1. Wash, remove from stems and discard any moldy or badly bruised grapes. Using nylon straining bag or press, crush grapes and extract maximum juice from pulp. Let juice run into primary fermentor and immediately add campden. Discard pulp. 2. Stir in all other ingredients EXCEPT yeast. Cover primary. 3. After 12 hours add yeast. Cover primary. 4. Check S.G. and when ferment reaches S.G. 1.040 (3-5 days) syphon wine off sediment into glass secondary. Attach airlock. 5. When ferment is complete (S.G. has dropped to 1.000--about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 6. To aid clearing syphon again in 2 months and again if necessary before bottling.

NATIVE AMERICAN GRAPES

Before the grape varieties of Europe were brought over and planted here, there were already growing an abundance of grape vines. Our native American varieties are mainly characterized by: a distinguishable pungent odor to the point of masking any other; higher in fruit acid but lower in sugar; and many varieties tolerate cold well below 0°F. A native variety, *Vitis Labrusca*, contains some of the most popularly planted vines-- Concord, Catawba, Niagara, etc. In most cases these grapes must be balanced by first diluting the juice to lower the acid level, sugar added back to assure proper alcohol production and pectic enzyme added to eliminate pectin (good for jelly, bad for wine). Whether you make wine from our native grapes which require major adjustments or from the natural juice of European varieties use the guide given on page 15.

RECIPE SELECTION:

Depending on the acid level and sugar content other native varieties such as: Catawba, Niagara, Delaware, or hybrids such as Baco, Aurore, Chelois; and grapes mainly found in the Southeast form the Muscadine family, the same care and the same recipes can be used as a guide. If the grape yields a juice balanced in acid and sugar, use recipes 46-50. If the acid levels are much too high, use recipes 51-56.



51. WILD GRAPES

Fox Grape, "Mustang" Grape
(Oregon Grapes see Recipe 75)

6 lb	Grapes
6 pts	Water
2 lb	Sugar
½ tsp	Pectic Enzyme
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine Yeast

Keep your acid tester and hydrometer handy. As with all wild fruit the sugar and acid content varies greatly from year to year and even from one location to another. The recipe above is a general recipe to use which you may have to adjust.

METHOD:

1. Wash, remove from stems and discard any moldy or badly bruised grapes. Using nylon straining bag or press, crush grapes and extract maximum juice from pulp. 2. Add enough water to juice to adjust acid level to .65% (tartaric). 3. Then add sugar to adjust Specific Gravity to 1.095-1.100. Add remaining ingredients including pulp tied in straining bag, EXCEPT for yeast. Cover primary. 4. After 24 hours, add yeast. Cover primary. 5. Stir daily, check S.G. and press pulp lightly to aid extraction. 6. When ferment reaches S.G. 1.030 (5 days) lightly press juice from bag. Then syphon wine off sediment into glass secondary. Attach the airlock. 7. When ferment is complete (S.G. has dropped to 1.000-- about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 8. To aid clearing syphon again in 2 months and again if necessary before bottling.

This makes a robust wine-- give it time to age. You may prefer this sweetened back a bit. Do this before bottling by adding ½ tsp stabilizer, then ¼ to ½ lb dissolved sugar per gallon.

52. CONCORD

(Fresh-Tablewine)

6lb	Fresh Grapes
5 pts	Water
3¼ cup	Sugar
½ tsp	Pectic Enzyme
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine yeast

Starting S.G. 1.090-95
Acid Level .65%

54. CONCORD

(Fresh-Dessert)

16lb	Fresh Grapes
1¼ cup	Sugar
1 tsp	Pectic Enzyme
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine yeast

56. CONCORD

(Frozen)

32 oz	Frozen Grape Concentrate
6 pts	Water
1 cup	Sugar
1 tsp	Acid Blend
½ tsp	Pectic Enzyme
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine yeast

53. CONCORD

(Juice-Tablewine)

3 pts	Pure Juice (Welch's or equal)
5 pts	Water
3¼ cup	Sugar
2 tsp	Acid Blend
½ tsp	Pectic Enzyme
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine yeast

55. CONCORD

(Juice-Dessert)

1 gal	Pure Juice (Welch's or equal)
1¼ cup	Sugar
1 tsp	Pectic Enzyme
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine yeast

METHOD:

Concord is the most popularly planted native grape. But, there are several other grape juices available in the store that can be used. 1. Wash, remove from stems, and discard moldy or badly bruised grapes. Using nylon straining bag or press crush grapes and extract juice into primary fermentor. 2. Keeping pulp in bag, tie up, and place primary. If you are using store purchased juice this work is already done. In either case: 3. Stir in all ingredients EXCEPT yeast. Cover primary. 4. After 24 hours add yeast. Cover primary. 5. Stir daily, check S.G. and press pulp lightly to aid extraction. 6. When ferment reaches S.G. 1.030 (5-6 days) lightly press juice from bag. Then syphon wine off sediment into glass secondary. Attach airlock. 7. When ferment is complete (S.G. has dropped to 1.000-- about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 8. To aid clearing syphon again in 2 months and again if necessary before bottling.

The wines made from recipes 52, 53, and 56 can be left dry but most prefer it slightly sweetened. At bottling add ½ tsp Stabilizer, then stir in ½ cup dissolved sugar per gallon.

Wines from recipes 54 and 55 should be sweetened back to S.G. 1.010-1.020. At bottling add ½ tsp Stabilizer then stir in ¾-1 cup dissolved sugar per gallon.

57. Wines from Grape Concentrates

For those who prefer winemaking without the work— premium grape juice concentrates are the answer. These concentrates are available for winemaking throughout the year. From France, Italy, Spain and California come premium wine grapes that normally only the wineries have access to. They are now freely available from the many home winemaker's stores. Most are in container sizes to make 1 or 5 gallons of wine. This recipe for 1 gallon of wine can be used as a general guide:

1 Can	Concentrate
	Water to 1 gallon
	Sugar to S.G. 1.085-1.095
	Acid Blend to .65% tartaric
1 tsp	Nutrient
1/8-1/4 tsp	Tannin
1	Campden, crush
1 pkg	Wine Yeast

METHOD:

All the work of fruit preparation has been done. 1. Mix all ingredients EXCEPT yeast in primary fermentor. Cover primary. 2. After 24 hours add yeast. Cover primary. 3. Stir daily and check Specific Gravity. 4. When ferment reaches S.G. 1.040 (3-5 days) syphon off sediment into glass secondary. Attach airlock. 5. When ferment is complete (S.G. has dropped to 1.000-- about 3 weeks) syphon to clean secondary. 6. To aid clearing syphon again in 2 months and again if necessary before bottling.

USE OF GRAPE CONCENTRATES TO IMPROVE OTHER WINES

The body and vinosity (wine taste quality) can be improved with the addition of small quantities of high quality grape concentrates. For every gallon of wine add 6-8 oz. of concentrate; reduce sugar added by 1/2 lb and acid blend by 1/2 tsp.

58. Instant Wines, GRAPE

1 1/2	Red grape conc.	3 lb	Blkberry, Rspb or Strawb.
7 1/2 pts	Water		
1/4-1 lb	Sugar	7 pts	Water
2 1/2 tsp	Acid Blend	1 1/2 lb	Sugar
1/4 tsp	Tannin	1/2 tsp	Acid Blend
1/4 tsp	Energizer	1/2 tsp	Pectic Enzyme
1	Campden, crush	1/4 tsp	Energizer
1 pkg	Wine yeast	1	Campden, crush
		1 pkg	Wine yeast

Later: 1 tsp Sparkolloid; 1 tsp Stabilizer
Starting S.G. 1.075 Acid Level .50-.55%

METHOD:

Instant wines, ready for drinking within a few weeks of starting ferment make pleasant drinks in warmer weather. Such wines have a low alcohol content of 7-8% and they should be consumed within 6-8 months. They have a light fresh appeal with a hint of sweetness. Hold back and refrigerate 1/2 cup Grape Concentrate or juice from 1 lb fruit. With recipe 59, using nylon straining bag mash and squeeze juice from fruit into primary fermentor. Leave pulp in bag, tie top and put in primary. With recipe 58 no processing is necessary. 2. Mix in all ingredients EXCEPT yeast. Cover primary and ferment at 70°-80° F. 3. Stir daily, check Specific Gravity, squeeze any pulp lightly. 4. At 1.030 (5 days) syphon off sediment (strain off juice from pulp) into glass secondary. Add Sparkolloid. Attach airlock. 5. When ferment reaches 1.000-.995 (12-14 days) the wine may be transferred to clean secondary for clearing (4-6 weeks). Add Stabilizer, then stir in concentrate or fruit juice and 1/3 cup dissolved sugar. Or with the use of a filter unit (after mixing in ingredients) syphon directly into bottles.

60. MEAD, Dry

2½ lb	Honey
7 pts	Water
3 tsp	Malic Acid
1½ tsp	Tartaric Acid
¼ tsp	Tannin
¼ tsp	Energizer
1	Campden, crush
1 pkg	Wine yeast, preferably German types or Champ.

61. MEAD, Sweet

3 ¼ lb	Honey
7 pts	Water
3 tsp	Malic Acid
1½ tsp	Tartaric Acid
¼ tsp	Tannin
¼ tsp	Energizer
2	Campden, crush
1 pkg	Sauternes yeast

62. MELOMEL

(Honey & Fruit)

2lb	Blackcurrants or
2½ lb	Raspberry
2½ lb	Honey
6 pts	Water
2 tsp	Acid Blend
½ tsp	Pectic Enzyme
¼ tsp	Energizer
1	Campden, crush
1 pkg	Wine yeast

63. MELOMEL

1¼ pt	Orange Juice
2pt	Pineapple juice (unsweetened)
½ pt	White grape conc.
1¼ lb	Honey
6 pts	Water
1½ tsp	Acid Blend
1/16tsp	Tannin
¼ tsp	Energizer
1	Campden, crush
1 pkg	Wine yeast

64. PYMENT

(Honey & Grape)

1½ lb	Honey
1 pt	White grape conc. or 2lb
6½ pt	Water
3 tsp	Acid Blend
¼ tsp	Tannin
¼ tsp	Energizer
1	Campden, crush
1 pkg	Wine yeast, pre-

65. METHEGLYN

(Honey & Herbs)

3 lb	Honey
4	Cloves
½ oz	Root Ginger
1 oz	Elderflower (optional)
7 pts	Water
3 tsp	Acid Blend
¼ tsp	Energizer
1	Campden, crush
1 pkg	Wine yeast

METHOD:

1. Wash and drain any fruit; put fruit and herbs into nylon straining bag. Crush and strain out juice into primary fermentor. Keeping pulp in bag, tie top and place in primary. 2. In primary, dissolve honey in ½ gallon of warm water with acid and tannin. Add remaining ingredients EXCEPT yeast. Cover primary. 3. After 24 hours add yeast. Cover primary. 4. Stir daily, check S.G. and press pulp lightly to aid extraction. 5. When ferment reaches S.G. 1.030 (5-6 days) lightly press juice from bag. Then syphon wine off sediment into glass secondary. Attach airlock. 6. When ferment is complete (S.G. has dropped to 1.000—about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. To aid clearing syphon again in 2 months and again if necessary before bottling.

66. JELLY and JAM

3 lb	Jelly or Jam (any flavor)
6 pts	Water
1 lb	Sugar
2 tsp	Acid Blend
1/4 tsp	Tannin
1 tsp	Nutrient
1 tsp	Pectic Enzyme
1 pkg	Wine yeast

Starting S.G. 1.095-1.100

METHOD:

The selection is limitless-- Strawberry, Grape, Peach, Raspberry, Orange, Plum, etc. 1. Scoop jam or jelly into primary fermentor. Stir in water (use cool not warm) and jelly or jam thoroughly. 2. Stir in all other ingredients, then, add yeast. Cover primary. 3. Stir daily, check S.G. 4. When ferment reaches S.G. 1.040 (3-5 days) syphon off wine sediment into glass secondary. Attach airlock. 5. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 6. To aid clearing syphon again in 2 months and again if necessary before bottling.

If you wish a slightly sweet wine at bottling: add 1/2 tsp stabilizer, then stir in 1/4 lb dissolved sugar per gallon.

67. MALT 'QUICKY' 68. MALT 'LIQUEUR'

1 cup	Lt. Malt Extract (dry or liquid)	1 1/2 lb	Lt. Malt Extr. (dry or liquid)
1 pt	Wht grape conc. or 5 cups raisins	1/2 pt	Wht grape conc
7 pts	Water	6 pts	Water
1/4 cup	Sugar	2 1/2 lb	Sugar
1/4 tsp	Acid Blend	3 1/2 tsp	Acid Blend
1 tsp	Nutrient	1/4-1/2 tsp	Tannin
1	Campden, crush	1 tsp	Nutrient
1 pkg	Wine yeast	1	Campden, crush
		1 pkg	Wine yeast

METHOD:

1. Pour malt in primary fermentor. If using raisins, chop up, put in straining bag and put in primary. 2. Stir in all other ingredients, EXCEPT yeast. Cover primary. 3. After 24 hours add yeast. Cover primary. 4. Stir daily, check S.G. 5. When ferment reaches S.G. 1.040 (3-5 days) syphon off wine sediment into glass secondary. Attach airlock. 6. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. To aid clearing syphon again in 2 months and again if necessary before bottling.

The Malt "Liqueur" (recipe 68) may be further sweetened at bottling: add 1/2 tsp Stabilizer, then stir in 1/4-1/2 lb dissolved sugar per gallon.

69. MELON

3 lb	Melon centers	2 qts	Watermelon juice
7 pts	Water	2 qts	Water
2 lb	Sugar	1½ lb	Sugar
2 tsp	Acid Blend	2½ tsp	Acid Blend
½ tsp	Tannin	¼ tsp	Tannin
1 tsp	Nutrient	1 tsp	Nutrient
1 pkg	Wine yeast	1	Campden, crush
		1 pkg	Wine yeast

Starting S.G. 1.085

METHOD:

1. Cut melons in quarters, remove all rind parts entirely and discard seeds. Cut "meat" of melons into cubes. 2. Using nylon straining bag, mash and squeeze out juice into primary fermentor. Keeping pulp in straining bag, tie top and place in primary. 3. Stir in all other ingredients, EXCEPT yeast. Cover primary. 4. After 24 hours add yeast. Cover primary. 5. Stir daily, check S.G. 6. When ferment reaches S.G. 1.040 (3-5 days) lightly press juice from bag. Syphon off wine sediment into glass secondary. Attach airlock. 7. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 8. To aid clearing syphon again in 2 months and again if necessary before bottling.

70. WATERMELON

2 qts	Watermelon juice
2 qts	Water
1½ lb	Sugar
2½ tsp	Acid Blend
¼ tsp	Tannin
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine yeast

Starting S.G. 1.085

71. MT. ASH

(Rowanberry)

2 lb	Berries
7 pts	Water
2 lb	Sugar
1 tsp	Acid Blend
½ tsp	Pectic Enzyme
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine yeast

METHOD:

Use ripe, sweet berries and if possible pick after 1st frost. 1. Remove stem, rinse off and drain berries. 2. Put in nylon straining bag, crush and squeeze juice into primary fermentor. Keeping pulp in bag, tie top and place in primary. 3. Stir in all other ingredients, EXCEPT yeast. Cover primary. 4. After 24 hours, add yeast. Cover primary. 5. Stir daily, check S.G. and press pulp lightly to aid extraction. 6. When ferment reaches S.G. 1.040 (3-5 days) squeeze juice lightly from bag. Syphon off wine sediment into glass secondary. Attach airlock. 7. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 8. To aid clearing syphon again in 2 months and again if necessary before bottling.

To sweeten wine at bottling, add ½ tsp Stabilizer then stir in ¼ lb dissolved sugar per gallon.



VARIETIES

Mountain Ash (*Sorbus L.*) normally a 20'-30' tree. White flowers followed by attractive clusters of ¼" bright orange fruit. American Mt. Ash (*S. Americana*) and European Mt. Ash (*S. Aucuparia*) are the two popular varieties. Sometimes referred to as Rowanberry.

72. ONION

½ lb	Onions
½	Potatoes)
½ pt	Wht grape conc. or 1 lb Raisins
7 pts	Water
2¼ lb	Sugar
2 tsp	Acid Blend
1 tsp	Nutrient
1	Campden.crush
1 pkg	Wine yeast

METHOD:

Don't tell your friends what it is until they've tried some. It's a surprising wine that even works well for marinating meats or whatever. 1. Wash and peel onions. Chop up onions and potatoes. 2. Put in nylon straining bag, tie top and put in pot with water. Boil until tender (about 15 min.) Strain off and press pulp into pot. Discard pulp. Pour liquour in pot into primary fermentor. 3. Stir in all other ingredients, EXCEPT yeast. If using raisins, chop up and put in bag, tie, put in primary. Cover primary. 4. After 24 hours, add yeast. Cover primary. 5. Stir daily, check S.G. 6. When ferment reaches S.G. 1.040 (3-5 days) strain juice off raisins and syphon off wine sediment into glass secondary. Attach airlock. 7. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 8. To aid clearing syphon again in 2 months and again if necessary before bottling.

73. ORANGE

8-9	Oranges
½ pt	Wht grape conc. or 1 lb Lt raisins
6 pts	Water
2½ lb	Sugar
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine yeast

74. ORANGE

(Frozen)

12 oz	Frozen Juice
½ pt	Wht grape conc. or 1 cup Lt dry Malt
7 pts	Water
2½ lb	Sugar
½ tsp	Energizer
1	Campden,crush
1 pkg	Wine yeast

METHOD:

Juice oranges like Temples work the best. Scrape the rind of 2 oranges. Discard the bitter white pith and peels. 1. Using nylon straining bag, separate orange segments, squeeze juice into primary fermentor. Keeping pulp and rind in bag, put chopped raisins in bag, tie top and place in primary. If using frozen juice pour in primary 2. Add sugar, pour over water (warm) and stir in all other ingredients EXCEPT yeast. Cover primary. 3. After 24 hours, add yeast. Cover primary. 4. Stir daily, check S.G. and with fresh orange recipe press pulp lightly to aid extraction. 5. When ferment reaches S.G. 1.040 (3-5 days) squeeze juice lightly from bag. Syphon off wine sediment into glass secondary. Attach airlock. 6. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. To aid clearing syphon again in 2 months and again if necessary before bottling.

These wines may be sweetened at bottling by: Add ½ tsp. Stabilizer then add ¼-½ lb dissolved sugar per gallon.

75. OREGON 'GRAPE'

(Mahonia)

3 lb	Oregon Grapes
½ pt	Red grape conc. or 1 lb Dk raisins
7 pts	Water
1¼ lb	Sugar
½ tsp	Pectic Enzyme
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine yeast

METHOD:

Although they look like grapes these fruit come from a different part of the plant world. But, for winemaking purposes these berries can be treated as wild grapes. The sugar and acid content will vary greatly so use your hydrometer and acid tester to adjust to Specific Gravity-- 1.095-1.100; Acid level .65%. If you can wait, pick late season. 1. Wash, remove from stems, and discard moldy or badly bruised berries. Using nylon straining bag or press crush berries and strain juice into primary fermentor. 2. Keeping pulp in bag, tie top and place in primary. 3. Stir in all other ingredients, EXCEPT yeast. Cover primary. 4. After 24 hours, add yeast. Cover primary. 5. Stir daily, check S.G. and press pulp lightly to aid extraction. 6. When ferment reaches S.G. 1.030 (3-5 days) lightly press juice from bag. Syphon off wine sediment into glass secondary. Attach airlock. 7. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 8. To aid clearing syphon again in 2 months and again if necessary before bottling.

At bottling add ½ tsp Stabilizer, then ¼ lb dissolved sugar per gallon.



VARIETIES

Oregon Grape (*Mahonia aquifolium*), ½" blue-black berries in clusters look grape-like, with holly-like leaves. Attractive all year. Along with Desert Mahonia (*M. fremontii*), California Holly Grape (*M. pinnata*), Creeping Mahonia (*M. repens*) found in the wild and some from nursery.

76. PEAR

4 lb	Pears
6 pts	Water
4 cups	Sugar
2½ tsp	Acid Blend
½ tsp	Pectic Enzyme
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine yeast

77. PERRY

1 gal	Pear Juice (12-14lb) to 1.060 S.G. Pectic Enzyme Energizer Campden, crush Champ. yeast
Sugar	
½ tsp	
¾ tsp	
1	
1 pkg	

METHOD:

Use ripe to firm-ripe pears. 1. Wash, drain, remove stems, cut in half and core. Cut into smaller pieces. 2. Using nylon straining bag, (or press) mash and strain out juice into primary fermentor. As juice is extracted immediately add campden to prevent spoilage and browning. Keeping all pulp in bag, tie top and place in primary. 3. Stir in all other ingredients EXCEPT yeast. Cover primary. 4. After 24 hours, add yeast. Cover primary. 5. Stir daily, check S.G. and press pulp lightly to aid extraction. 6. When ferment reaches S.G. 1.040 (3-5 days) strain juice lightly from bag. Syphon off wine sediment into glass secondary. Attach airlock. 7. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 8. To aid clearing syphon again in 2 months and again if necessary before bottling.

For Perry, when ferment is complete gently stir in ¼ cup dissolved sugar per gallon. Bottle in champagne or returnable pop bottles. Apply crown caps. Age for 3 months. Cool and enjoy.

VARIETIES

Pear trees (*Pyrus communis*) make a handsome addition to the garden. Properly pruned and cared for they take heavy soil better than most. Good ones to plant are: Bartlett, Bosc, Packham Triumph, Seckel and Clapp Favorite. Most pears prefer cool winters.



78. PERSIMMON

3 lb	Persimmons
7 pts	Water
2½ lb	Sugar
2½ tsp	Acid Blend
½ tsp	Pectic Enzyme
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine yeast

METHOD:

Pick fully ripe preferably after 1st frost. 1. Wash and drain Persimmons. Cut in half and remove seeds. Cut into chunks and using nylon straining bag, crush, mash and squeeze out juice into primary fermentor. 2. Keeping pulp in bag, tie top and place in primary. Stir in all other ingredients, EXCEPT yeast. Cover primary. 3. After 24 hours, add yeast. Cover primary. 4. Stir daily, check S.G. and press pulp lightly to aid extraction. 6. When ferment reaches S.G. 1.040 (3-5 days) strain juice lightly from bag. Syphon off wine sediment into glass secondary. Attach airlock. 6. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. To aid clearing syphon again in 2 months and again if necessary before bottling.

If a slightly sweetened wine is more to taste, then, at bottling add ½ tsp Stabilizer, then stir in ¼-½ lb dissolved sugar per gallon.

VARIETIES:

Persimmons (*Diospyros* L.) besides the fruit have outstanding ornamental qualities. One of the best fruit trees for ornamental use-- good in the garden or as a small shade tree. The American Persimmon (*D. Virginiana*) is a 20'-30' tree with round, yellow to orange 1½"-2" fruit (puckery until soft-ripe, then very sweet). Oriental persimmon (*D. kaki*) attractive autumn color even in warm winter areas. Several varieties include: Chocolate, Fuyu and Hachiya.

79. PINEAPPLE

(Fresh)

3 lb	Pineapple
6 pts	Water
1¼ lb	Sugar
¼ tsp	Tannin
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine yeast

Starting S.G. 1.090-95

80. PINEAPPLE

(Juice)

1 gal	Pineapple Juice (unsweetened)
5 pts	Water
2 lb	Sugar
½ tsp	Acid Blend
¼ tsp	Tannin
1	Campden, crush
1 pkg	Wine yeast

METHOD:

Select fully ripe pineapples. Cut off stalk and outside rind. 1. Cut in half and remove hard core. Chop in small pieces. Put in nylon straining bag, mash and squeeze out juice into primary fermentor. Keeping all pulp in bag, tie top and place in primary. With pineapple juice (recipe 80) this work is already done. 2. Stir in all other ingredients EXCEPT yeast. Cover primary. 3. After 24 hours, add yeast. Cover primary. 4. Stir daily, check S.G. and press pulp lightly to aid extraction. 5. When ferment reaches S.G. 1.040 (3-5 days) strain juice lightly from bag. Syphon off wine sediment into glass secondary. Attach airlock. 6. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. To aid clearing syphon again in 2 months and again if necessary before bottling.

These wines are very pleasant as a dry wine. If a slightly sweetened wine is more to taste, at bottling add ½ tsp Stabilizer, then stir in ¼ lb dissolved sugar per gallon.

81. PLUM

4 lb	Plums, pitted	4 lb	Plums, pitted
6 pts	Water	½ pt	Red grape conc.
2 lb	Sugar	6 pts	Water
½ tsp	Acid Blend	2¼ lb	Sugar
½ tsp	Pectic Enzyme	½ tsp	Pectic Enzyme
1 tsp	Nutrient	1 tsp	Nutrient
1	Campden, crush	1	Campden, crush
1 pkg	Wine yeast	1 pkg	Sherry or Port yeast

METHOD:

You can use these recipes for any plum-type fruit—home grown or store bought; Italian, Damson, Yellow, Greengage, or any sweet plum. With wild plums, which are generally high in acid, use acid tester or cut down to 3 lbs per gallon. 1. Wash, drain, remove stones, chop into smaller pieces. 2. Put in nylon straining bag, crush and squeeze juice into primary fermentor. Keeping all pulp in bag, tie top and place in primary. 3. Stir in all other ingredients EXCEPT yeast. Cover primary. 4. After 24 hours, add yeast. Cover primary. 5. Stir daily, check S.G. and press pulp lightly to aid extraction. 6. When ferment reaches S.G. 1.040 (3-5 days) squeeze juice lightly from bag. Syphon off wine sediment into glass secondary attach airlock. 7. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 8. To aid clearing syphon again in 2 months and again if necessary before bottling.

To sweeten wine, at bottling add ½ tsp Stabilizer, then stir in ¼-½ lb dissolved sugar per gallon.



VARIETIES

Japanese (*Prunus salicina*) and European (*p. domestica*) Plums are attractive as well as productive 15'-20' trees. European plums are more tart and not as juicy as Japanese plums. Here are some self-fertile varieties to try: Damson, Greengage, Italiana Plum, Santa Rosa and Stanley.

82. PLUM 'PORT'

83. POMEGRANATE

6	Pomegranates	8	Pomegranates
½ pt	Wht grape conc.	1 cup	Dry Mlt. Extret.
6 pts	Water	6 pts	Water
1¼ lb	Sugar	2 lb	Sugar
2 tsp	Acid Blend	2 tsp	Acid Blend
¼ tsp	Tannin	¼ tsp	Tannin
1 tsp	Pectic Enzyme	1 tsp	Pectic Enzyme
1 tsp	Nutrient	1 tsp	Nutrient
1	Campden, crush	1	Campden, crush
1 pkg	Wine yeast	1 pkg	Wine Yeast

84. POME. & MALT

METHOD:

1. Wash, peel, removing all yellow skin. Split open pomegranates and scrape out seeds. 2. Chop into small pieces. Put in nylon straining bag, crush fruit and squeeze juice into primary fermentor. Keeping pulp in bag, tie top, and place in primary. 3. Stir in all other ingredients EXCEPT yeast. Cover primary. 4. After 24 hours, add yeast. Cover primary. 5. Stir daily, check S.G. and press pulp lightly to aid extraction. 6. When ferment reaches S.G. 1.050 5-6 days) squeeze juice lightly from bag. Syphon off wine sediment into glass secondary attach airlock. 7. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 8. To aid clearing syphon again in 2 months and again if necessary before bottling.

If a slightly sweetened wine is preferred, at bottling add ½ tsp Stabilizer, then stir in ¼ lb dissolved sugar per gallon.

VARIETIES

Pomegranates (*Punica granatum*) tolerate great heat but fruiting varieties need deep watering. Available in tree or shrub form, 1½'-12' high. The two popular varieties to try are: "Wonderful", 10' high, large orange-red flowers followed by reddish fruit and "Sweet", a pomegranate similar to Wonderful but with sweeter, less acid fruit.



85. POTATO

2 lb Potatoes
½ pt Wht grape conc.
7 pts Water
2¼ lb Sugar
3 tsp Acid Blend
½ tsp Tannin
1 tsp Nutrient
1 pkg Wine yeast

86. "MASHER"

4 lb Potatoes
½ pt Wht grape conc
or 1 lb raisins
7 pts Water
2½ lb Sugar or Honey
3 tsp Acid Blend
½ tsp Energizer
1 pkg Wine Yeast

METHOD:

An old favorite that kept many a pioneer going. 1. Wash and scrub potatoes clean. Slice up potatoes and boil in water just until tender. Strain out juice, remove potatoes and pour the boiled mixture into primary fermentor. If you have nylon straining bag use this to contain potatoes while boiling. 2. Stir in all other ingredients EXCEPT yeast. Cover primary. 3. When mixture cools (below 85°F.) add yeast. Cover primary. 4. Stir daily, check S.G. 5. When ferment reaches S.G. 1.040 (3-5 days) strain juice from raisins, if used, syphon wine into glass secondary. Attach airlock. 6. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. To aid clearing syphon again in 2 months and again if necessary before bottling.

Sweeten wine at bottling by adding ½ tsp Stabilizer and then stir in ¼ lb dissolved sugar per gallon.



87. PRICKLY PEAR

3 qts Prickly Pear
1¼ cup Raisins or ½ pt Grape Conc.
2 pts Water
2 lb Sugar
2 tsp Acid Blend
1 tsp Pectic Enzyme
¼ tsp Energizer
1 Campden, crush
1 pkg Sherry or Port Yeast

METHOD:

Don't head for the grocery store. These "pears" are sometimes purple, sometimes orange, sticker covered pear-shape fruits growing on the top edge of flat pad-shaped cactus. Cut off fruit. 1. Break or singe off stickers. Wash, chop and mash up fruit. Chop up raisins, if used. Put all in nylon straining bag, tie top, and leave bag in primary fermentor. Using nylon straining bag or press crush berries and strain juice into primary fermentor. 2. Stir in all other ingredients, EXCEPT yeast. Cover primary. 3. After 24 hours, add yeast. Cover primary. 4. Stir daily, check S.G. and press pulp lightly to aid extraction. 5. When ferment reaches S.G. 1.040 (3-5 days) lightly press juice from bag. Syphon off wine sediment into glass secondary. Attach airlock. 6. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. To aid clearing syphon again in 2 months and again if necessary before bottling.

Wine can be sweetened at bottling by adding ½ tsp Stabilizer and then stir in ¼ lb dissolved sugar per gallon

VARIETIES

Prickly Pear (*Opuntia* L.) is a large genus; in fact one variety or another can be found in most parts of the U.S. Easter Prickly Pear (*O. compressa*) with red or purple fruit. Hunger Cactus (*O. polyacantha*) yields spiny fruit. The illustration shows the repeated flat pads. Flowers appear followed by fruit, sometimes spiny.



88. PUMPKIN

4 lb	Pumpkin "meat"
7 pts	Water
2¼ lb	Sugar
2½ tsp	Acid Blend
¼ tsp	Tannin
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine Yeast

METHOD:

1. Wash, cut open pumpkin, remove seeds and stringy material. Rough peel skin. Grind or mash up pumpkin into nylon straining bag. Squeeze juice out into primary fermentor, tie top, and leave bag in primary fermentor. Using nylon straining bag or press crush berries and strain juice into primary fermentor. 2. Stir in all other ingredients, EXCEPT yeast. Cover primary. 3. After 24 hours, add yeast. Cover primary. 4. Stir daily, check S.G. and press pulp lightly to aid extraction. 5. When ferment reaches S.G. 1.040 (3-5 days) lightly press juice from bag. Syphon off wine sediment into glass secondary. Attach airlock. 6. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. To aid clearing syphon again in 2 months and again if necessary before bottling.

You may prefer to age this wine dry or sweeten it back slightly. Sweeten at bottling: Add ½ tsp Stabilizer, then add in ¼ lb dissolved sugar per gallon.



89. RAISIN

2 lb	Raisins
7 pts	Water
1¼ lb	Sugar
3½ tsp	Acid Blend
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine yeast

90. RAISIN SHERRY

3 lb	Raisins
1 cup	Lt Dry Malt
6 pts	Water
2 lb	Sugar
½ tsp	Acid Blend
½ tsp	Energizer
1	Campden, crush
1 pkg	Sherry Yeast

METHOD:

1. Chop or use mincer on seedless light or dark raisins. Place in nylon straining bag, tie top, and put in primary fermentor. 2. Stir in all other ingredients EXCEPT yeast. Cover primary. 3. Stir daily, check S.G. 4. When ferment reaches S.G. 1.040 (3-5 days) strain juice from raisins, if used, syphon wine into glass secondary. Attach airlock. 5. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 6. To aid clearing syphon again in 2 months and again if necessary before bottling.

Recipe 89 makes a good dry wine. If you prefer it slightly sweetened, then add ¼ lb dissolved sugar per gallon.

Recipe 90, The "Sherry" can have additional doses of ¼ lb dissolved sugar. Add every time it continues to work down to S.G. 1.000.

91. RHUBARB

3 lb	Rhubarb	5 lb	Rhubarb
¼ pt	Wht grape conc.	6 pts	Water
7 pts	Water	2 lb	Sugar
2¼ lb	Sugar	½ oz	Precip. Chalk
¼ tsp	Tannin	¼ tsp	Tannin
1 tsp	Nutrient	1 tsp	Nutrient
1	Campden, crush	1	Campden, crush
1 pkg	Wine yeast	1 pkg	Wine Yeast

Starting S.G. 1.095-1.100 Starting S.G. 1.095-1.100
At bottling: Add 1/8 tsp. anti-oxidant

METHOD:

Select and use stock only. Wash, drain and cut into small pieces. 1. Place rhubarb in nylon straining bag, mash and squeeze out juice into primary fermentor. Keeping all pulp in straining bag, tie top, and put in primary fermentor. Mix in sugar and campden. Cover primary. 2. Let mixture sit for 2-3 days stirring once daily. Then strain juice from bag. Discard pulp. 3 Stir in all other ingredients (use cold water), then add yeast. Cover primary. 4. Stir daily and check Specific Gravity. 5. When ferment reaches S.G. 1.040 (3-5 days) syphon wine into glass secondary. Attach airlock. 6. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. To aid clearing syphon again in 2 months and again if necessary before bottling.

You may prefer them dry or slightly sweetened back. To sweeten: Add ½ tsp Stabilizer, then stir in ¼ lb dissolved sugar per gallon.

For and old fashioned German Rhubarb wine add to either recipe ¼ oz. Ginger Root (bruised) and ¼ oz Cloves per gallon. Tie up in straining bag, then remove at S.G. 1.040.

92. RHUBARB


93. RICE "SAKI"

2½ lb	Rice
½ pt	Grape conc or 1 lb Lt Raisins
7 pts	Water (hot)
2¼ lb	Sugar or Honey
3 tsp	Acid Blend
¼ tsp	Energizer
1	Campden, crush
1 pkg	Sherry Yeast

METHOD:

Use husked or raw rice if available, rather than polished rice. 1. Wash and coarse crush rice. Place rice and chopped raisins in nylon straining bag, tie top and place in primary fermentor. 2. Pour hot water over and stir in all other ingredients, EXCEPT yeast. Cover primary. 3. After 48 hours, add yeast and energizer. Cover primary. 4. Stir daily, check S.G. and press pulp lightly to aid extraction. 5. When ferment reaches S.G. 1.050 (2-3 days) add another ¼ lb dissolved sugar per gallon. 6. At S.G. 1.030 (6-7) days strain juice from bag. Syphon off wine sediment into glass secondary. Attach airlock. 7. At S.G. 1.020 add another ¼ lb dissolved sugar per gallon. 8. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 9. To aid clearing syphon again in 2 months and again if necessary before bottling.

Optional: You may continue to build up alcohol by adding additional doses of sugar until ferment ceases. To sweeten add before bottling ½ tsp Stabilizer, then add ¼ lb dissolved sugar per gallon.



94. ROSE HIP

(Fresh)

2½ lb Fresh Rosehips
7½ pts Water (hot)
2 lb Sugar
1 tsp Acid Blend
½ tsp Energizer
1 Campden, crush
1 pkg Wine yeast

95. ROSE HIP

(Dried)

5 oz Rosehips (dried shells) or 10 oz (dried, whole)
1½ cup Lt Raisins or 3 oz Dried Fig or 4 oz grape conc.
1 gal Water
2 lb Sugar
1 Campden, crush
1 pkg Wine yeast

METHOD:

With fresh hips, cut off stalks and any foreign matter. In the wild the red hips can be found on the dried stalks even through winter. 1. Wash and drain. Crush or run through mincer (dried or fresh). Put in nylon straining bag, (if using raisins or fig, chop and put in bag), tie top and place in primary. 2. Pour over hot water and stir in all other ingredients EXCEPT yeast. Cover primary. 3. After 24 hours, add yeast. Cover primary. 4. Stir daily, check S.G. and press pulp lightly to aid extraction. 5. When ferment reaches S.G. 1.040 (3-5 days) strain juice lightly from bag. Syphon off wine sediment into glass secondary. Attach airlock. 6. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 7. To aid clearing syphon again in 2 months and again if necessary before bottling.

To sweeten: Before bottling add ½ tsp Stabilizer, then add ¼ lb dissolved sugar per gallon. With recipe No. 94 for a Sherry type wine start with Sherry Yeast and at 1.000 continue to add ¼ lb dissolved sugar per gallon until ferment ceases.



96. RUM POT

Sometimes called "drunken fruit" this delicious mix grows and grows. When it gets going you can give it to friends 1 cup at a time. Always leave 1 cup for yourself

1 cup Canned Fruit Cocktail
1 cup Sugar
pinch of Wine Yeast
Liquor, optional

1. Put fruit, then sugar in glass jar or small crock (1 qt). Sprinkle on pinch of yeast. Partly cover. 2. After 24 hours, stir. 3. Let sit partly covered for 2 weeks. 4. Then again add 1 cup fruit and 1 cup sugar. Partly cover stir, let sit for 2 weeks. 5. Repeat step 4 as required. Keep at room temperature. Refrigerate to stop action. Add to ice cream, pudding or use as a glaze. Also excellent with a touch of Brandy, White Wine, Creme de Menthe or Galliana.

97. SAN GRIA

Call it Sant Gria, Sangaree, or San Gria here are some recipes for some different "true" Sangrias:

1 Bottle Red or White Wine
1 Whole Sliced Orange
1 Whole Sliced Lemon
¾ Cup Sugar, or to taste
6 oz Soda Water
¾ oz Brandy, Grand Marnier, or Cointreau
6-8 Strawberries, halved
or 4-6 slices Elberta Peaches

Slice oranges and lemons thinly into server. Pour in wine, stir in sugar thoroughly, add liquor, and sprinkle on fruit. Place in refrigerator to cool. Just before serving add chilled soda and ice cubes.

98. SERVICE BERRY

3 lb	Berries
7 pts	Water
2 1/4 lb	Sugar
3 tsp	Acid Blend
1/2 tsp	Pectic Enzyme
1/2 tsp	Energizer
1	Camden, crush
1 pkg	Wine Yeast

Starting S.G. 1.095-1.100

METHOD:

1. The best time to pick these berries is after the first frost. Remove stems, leaves, and foreign matter. Wash and drain. 2. Using nylon straining bag mash and strain out juice into primary fermentor. Keeping all pulp in straining bag, tie top, and place in primary. 3. Stir in all other ingredients, EXCEPT yeast. Cover primary. 4. After 24 hours, add yeast and Cover primary. 5. Stir daily, check S.G. and press pulp lightly to aid extraction. 6. When ferment reaches S.G. 1.030 (5 days) syphon off wine sediment into glass secondary. Attach airlock. 7. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 8. To aid clearing syphon again in 2 months and again if necessary before bottling.

To sweeten: Before bottling add 1/2 tsp Stabilizer, then add 1/4 lb dissolved sugar per gallon.

VARIETIES

Serviceberries (*Amelanchier* L.) commonly called Juneberry also Shadbush and Sarvisberry. An attractive shrub anywhere from 5'-25', white flowers, 1/4" bluish-black small crab-apple like fruit. Saskatoon Serviceberry (*A. alnifolia*) is a common variety. Oblong leaf Juneberry (*A. canadensis*) can be found as tree also. Some are available from nursery.



99. STRAWBERRY

3 1/2 lb	Strawberries (fresh or frozen)
7 pts	Water
2 lb	Sugar
1 tsp	Acid Blend
1/4 tsp	Tannin
1/2 tsp	Pectic Enzyme
1 tsp	Nutrient
1	Camden, crush
1 pkg	Wine Yeast

Starting S.G. 1.090-95

METHOD:

Pick berries fully ripe. Remove any stems or leaves or foreign matter. 1. Wash and drain berries. 2. Using nylon straining bag (or press) mash and strain out juice into primary fermentor. Keeping all pulp in straining bag, tie top, and place in primary. 3. Stir in all other ingredients, EXCEPT yeast. Cover primary. 4. After 24 hours, add yeast and Cover primary. 5. Stir daily, check S.G. and press pulp lightly to aid extraction. 6. When ferment reaches S.G. 1.030 (about 5 days) syphon off wine sediment into glass secondary. Attach airlock. 7. When ferment reaches 1.000-- (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 8. To aid clearing syphon again in 2 months and again if necessary before bottling.

This makes a good dry wine, but you may slightly sweeten it also. Before bottling add 1/2 tsp Stabilizer, then add 1/4 lb dissolved sugar per gallon.



VARIETIES

Strawberries (*Fragaria* L.) are a popular fruit that needs little introduction. They can be found in the wild and if lucky you can spot the small 1/4" wild fruit. Here are some ever bearing commercial varieties: Ogalala, Streamliner, Ozark Beauty and Sequoia. Pinch first blossoms for plant strength. Some can be used as ground cover.

100. TOMATO, Red 101. TOMATO, Green

3½ lb	Red Tomatoes	3½ lb	Green Tomato
1 cup	Raisins	6	Cloves
6 pts	Water	½	Ginger Root
1½ lb	Sugar		(bruised)
2½ tsp	Acid Blend	6 pts	Water
¼ tsp	Tannin	2½ lb	Sugar
1 tsp	Nutrient	2 tsp	Acid Blend
1	Campden, crush	¼ tsp	Tannin
1 pkg	Wine yeast	1 tsp	Nutrient
		1	Campden, crush
		1 pkg	Wine yeast

METHOD:

1. Wash tomatoes. Remove any bruised portions. Cut into pieces. 2. Using nylon straining bag, mash and squeeze out juice into primary fermentor. Keeping all pulp in straining bag, (put ginger root and clove in bag) tie top and place in primary. 3. Stir in all ingredients EXCEPT yeast. Cover primary. 4. After 24 hours, add yeast. Cover primary. 5. Stir daily, check S.G. and press pulp lightly to aid extraction. 6. When ferment reaches 1.040 (3-5 days) lightly press juice from bag. Syphon off wine sediment into glass secondary. Attach airlock. 7. When ferment reaches 1.000— (about 3 weeks) syphon off sediment into clean secondary. Reattach lock. 8. To aid clearing syphon again in 2 months and again if necessary before bottling.

The wines can be left towards the dry side or you may prefer them sweetened back slightly. Before bottling add ¼ tsp Stabilizer, then add ¼ dissolved sugar per gallon.

Other Common Wines and Making Up Your Own Recipes

Most of the popular domesticated fruit and their found in the wild versions are covered in the hand book. Many of the wild plants listed here can also be found in local commercial nurseries suitable for your area. On these last pages we have listed other wine making material which use the same method and all other ingredients as recipes already given Simply substitute the ingredients.

There are also hundreds of other fruits and berries to be found in the wild. Some of the plants have been cultivated for garden use. Many are better left for the animals of the wild, but if you posses an adventurous spirit, jump right in. If they are edible and have a good taste they can be made into acceptable wines. Here is a basic recipe to follow for making up your own recipe and for those that are in doubt:

2-3 lb	Fruit or Berries
7 pts	Water
2-2½ lb	Sugar, adjust to 1.090-1.100
½-3 tsp	Acid Blend, adjust to .50-65% (tartaric)
½ tsp	Pectic Enzyme
1 tsp	Nutrient
1	Campden, crush
1 pkg	Wine Yeast

METHOD:

Sometimes it is necessary to dilute the pure juice below the normal acid level, then fruit acid blend (using acid tester) added back. This is done in order to balance the otherwise overpowering flavor of some fruit. The hydrometer is then used in adding enough sugar to achieve correct alcohol production. The goal is wine that tastes good with food-- a table wine. For these wines a hydrometer and acid tester are essential. Further information on how to make the wine is given on page 18, recipe 51, "Wild Grapes".

reference list

This is a combined list of other common wine making materials and fruits that may be native to your area.

BARBERRY (*Berberis L.*), ½" oval berry, bright red in drooping clusters, ripens Sept.-Oct.
 BEAUTYBERRY (*Callicarpa americana*), ¼" round, purple berries in clusters around leaf base, ripens Aug.-Oct. Also called French Mulberry.
 BILBERRY (*Vaccinium L.*), ¼" dark blue berry; also ½" pale purple Deerberry (*V. stamineum*); ¼ black Sparkelberry (*V. arboreum*); and the various cranberries and blueberries belong to the same family. Use Blueberry recipe pg 6. Most ripen July-Sept

BEARBERRY (*Arctostaphylos uva-ursi*), ¼" round, red, 5-10 seeds, ripen Aug.-Sept. Evergreen shrub also called Kinnikinnik. Excellent nursery variety;

"Radiant"

BLACKHAWK (*Viburnum prunifolium*), ¾" oval, blue-black, sweet taste, ripens Sept.-Oct. Also in family is Squashberry (*V. edule*), ¾" oval, red-orange, excellent fruit; Nannyberry (*V. lentago*) or Sheepberry, ½" oval. Use recipe above.

BOYSENBERRY (*Rubus L.*), See and use Blackberry recipe pg. 6.

BUFFALOBERRY (*Shepherdia L.*), ¼ oval berry, red or yellow, ripens June-Aug. Hardy, medium shrub with silver leaves.

CORALBERRY (*Symphoricarpos L.*), 3/16" round, purple to coral, ripens Sept.-Nov. Also called Indian Current. In this group is Wolfberry (*S. occidentalis*) ¾" round, dull white to blackish berry; and Snowberry (*S. albus*), ¼"-½" round, white (not evergreen) Snowberry.

CORN or MAPLE SYRUP-- use the same quantities (substitute Honey) as Mead recipes pg. 21.

CROWBERRY (*Empetrum L.*), ¼" or less round, black or purple, 6-9 seeds, ripens July-Nov. Use Raspberry recipe pg. 6.

DATES-- 1½ lbs. Use Raisin recipe pg. 29.

DEWBERRY (*Rubus L.*) See and use blackberry recipe pg. 6.

FIGS-- 2 lbs. Dried, or 4 lbs. Fresh. Use Raisin recipe pg. 29.

FLOWER PETAL or FLAVORED WINES-- Use Dandelion recipe pg 11 for following wines: 1 ¼ qts. Carnation petals; 2 qts, Clover petals; 1 qt fresh or 2 oz dried Elderflowers; 2 pts. Honeysuckle blossoms; 1 oz dried Limeflowers; 2 qts. Rosepetals; 2 oz dried Sarsaparilla; or 7 pts. Woodruff petals.

GUAVA-- 2½ lbs fruit plus 1 tsp Tannin. Although fruits are not similar, use method and all other ingredients of Persimmon recipe pg 26.

HAWTHORN BERRY (*Crataegus L.*), from tree of same name, ½"-¾" round oval berry, red with dotted surface, ripens Sept.-Oct. Columbian, Washington, Winter King and Pinnatifida varieties good fruiting trees. Use Mt. Ash recipe pg. 23.

HONEYSUCKLE (*Lonicera L.*), ¼" oval, specifically Mountain Fly Honeysuckle (*L. villosa*), blue to black fruit ripens July-Sept.

HUCKLEBERRY (*Gaylussacia L.*), ¾" round, blue-black, blueberry like but with 10 large seeds. Ripens June-Aug. In same group are Buckberry (*G. ursina*), and Dangleberry (*G. frondosa*). Use blueberry recipe pg 6.

JUNEBERRY (*Amelanchier L.*), same as Serviceberry, see pg 32. Also called Sarvisberry or Shadbush.

JUNIPER BERRY (*Juniperus communis*), ¼" oval, berry-like, bluish-black color. An evergreen. Numerous upright or spreading varieties available.

LEMON-- 7/8" lemons. Use Orange recipe pg 24

LINGOBERRY-- A small wild Cranberry. See and use Cranberry recipe pg 10.

LOGANBERRY-- In Raspberry family. See and use Raspberry recipe pg 6.

MULLBERRY (*Morus L.*), ½"-¾" Blackberry like fruit, juicy, dark red to black Black (*M. Nigra*) or persian Mullberry the best to plant (30' trees).

PARTRIDGE BERRY (*Mitchella repens*), ¼" round, single berry looks like two partly-joined berries, bright red, ripen Aug-Sept.

PAW PAW (*Asimina Adans*), 2½"-5" elongated oval, pulpy flesh with several large seeds, greenish yellow turns brown, ripens Aug-Oct.

PRUNE-- 3 lbs. , remove pits and mash. Use raisin recipe pg. 29.

QUINCE-- 3 lbs fruit, use only ½" tsp acid blend with Persimmon recipe pg. 26.

SALAL BERRY (*Gaultheria Shallon*), ¼"-¾" round, purple berry, ripens Sept.-Nov. Related to Teaberry (*G. procumbens*) also called checkerberry; has wintergreen odor. Use Raspberry recipe pg. 6.

SALMON BERRY (*Rubus spectabilis*), salmon pink edible fruit. One of the wild versions in the same family as Raspberry. Use the Raspberry recipe pg. 6.

TANGERINE-- 12-15 fruit. Use Orange recipe pg. 24.

WINTERBERRY (*Ilex verticillata*), ¼" bright red berry, ripens Sept.-Oct. Also called black Alder.

WINEBERRY (*Aristotelia racemosa*), ¼" red to black berries. Both male and female plant need to fruit.

SOME PLANTS TO AVOID:

Poison Ivy, Poison Oak and Poison Sumac should be avoided. All parts of these plants contain a dangerous skin irritant. Fruit of all three are about 3/16" round, in loose clusters, and a waxy white color.



Poison SUMAC

Poison IVY

Poison OAK



PAW PAW

HUCKLEBERRY



ALSO SEE ILLUSTRATIONS ON BACK COVER



BEARBERRY
(Kinnikinnick)



CROWBERRY CORALBERRY



BUFFALOBERRY



BEAUTYBERRY



PARTRIDGE
BERRY



BILBERRY



BLACK HAW



JUNIPER BERRY



TEADBERRY

