

Spatial Gaussian Process Regression for Bayesian Optical Flow Estimation


Andy Wang & Luka Orquiza

2 December 2025

Table of Contents

1. Background and Motivation
2. Probabilistic Method
3. Formulation of GP Regression
4. Implementation and Results
5. Discussion and Next Steps

Motion in Image Sequences



Given two sequential images $I : \mathcal{D} \times [0, T] \rightarrow \mathcal{R}$, i.e.

$$I(x, y, t) \text{ and } I(x, y, t + \delta t)$$

(for fixed t and δt), the optical flow at time t is a vector field

$$\mathbf{f}(x, y, t) = (u(x, y, t), v(x, y, t))$$

that transforms one image into the next:

$$I(x + u(x, y, t), y + v(x, y, t), t + \delta t) \simeq I(x, y, t)$$

Lucas-Kanade Method

Brightness constancy assumption:

$$\frac{d}{dt} I(x(t), y(t), t) = \frac{\partial I}{\partial x} u + \frac{\partial I}{\partial y} v + \frac{\partial I}{\partial t} = 0, \quad u := \frac{dx}{dt}, v := \frac{dy}{dt}$$

Underdetermined, so assume same $(u(x, y), v(x, y)) \forall (x, y) \in W$:

$$\underbrace{\begin{bmatrix} I_x(x_1, y_1) & I_y(x_1, y_1) \\ \vdots & \vdots \\ I_x(x_n, y_n) & I_y(x_n, y_n) \end{bmatrix}}_{\nabla I^T} \underbrace{\begin{bmatrix} u \\ v \end{bmatrix}}_{\mathbf{f}} = \underbrace{\begin{bmatrix} -I_t(x_1, y_1) \\ \vdots \\ -I_t(x_n, y_n) \end{bmatrix}}_{-\mathbf{I}_t}$$



Overdetermined: least squares solution is $\mathbf{v} = (\mathbf{A}^T \mathbf{A})^{-1} \mathbf{A}^T \mathbf{b}$:

$$\begin{bmatrix} u \\ v \end{bmatrix} = \begin{bmatrix} \sum I_x(x_i, y_i)^2 & \sum I_x(x_i, y_i) I_y(x_i, y_i) \\ \sum I_x(x_i, y_i) I_y(x_i, y_i) & \sum I_y(x_i, y_i)^2 \end{bmatrix}^{-1} \begin{bmatrix} -\sum I_x(x_i, y_i) I_t(x_i, y_i) \\ -\sum I_y(x_i, y_i) I_t(x_i, y_i) \end{bmatrix}$$

Optional weighting of pixels via diagonal matrix \mathbf{W} :

$$\mathbf{v} = (\mathbf{A}^T \mathbf{W} \mathbf{A})^{-1} \mathbf{A}^T \mathbf{W} \mathbf{b}$$

Lucas-Kanade Results



Probabilistic Motivation

Classical methods produce deterministic point estimates.
Uncertainty is inherent in optical flow estimation, due to:

- ▶ Image noise
- ▶ Brightness changes
- ▶ Low contrast regions
- ▶ Object occlusion
- ▶ Aperture problem
- ▶ Incompatible motions in localized regions

Quantification of confidence is desirable for safety-critical applications, e.g.:

- ▶ Autonomous navigation
- ▶ Computer-integrated surgery
- ▶ Real-time surveillance

Wang-Orquiza Method

Suppose $\mathbf{y} = \mathbf{A}\mathbf{x} + \boldsymbol{\eta}$, \mathbf{A} fixed, \mathbf{x} fixed unknown, \mathbf{y} observed, $\boldsymbol{\eta} \sim \mathcal{N}(\mathbf{0}, \mathbf{H})$:

$$\hat{\mathbf{x}}_{\text{MLE}}(\mathbf{y}) = \arg \max_{\mathbf{x}} f_{\mathbf{y}}(\mathbf{y}|\mathbf{x}) = \arg \max_{\mathbf{x}} f_{\boldsymbol{\eta}}(\mathbf{y} - \mathbf{A}\mathbf{x}) = \boldsymbol{\Sigma}_{\hat{\mathbf{x}}} \mathbf{A}^T \mathbf{H}^{-1} \mathbf{y}$$

$$\boldsymbol{\Sigma}_{\hat{\mathbf{x}}} = \text{Cov}[\hat{\mathbf{x}}_{\text{MLE}}(\mathbf{y}) - \mathbf{x}] = (\mathbf{A}^T \mathbf{H}^{-1} \mathbf{A})^{-1}$$

Same assumptions as Lucas-Kanade, plus additive Gaussian noise:



$$\mathbf{I}_t = -\nabla \mathbf{I}^T \mathbf{f} + \boldsymbol{\eta} \quad \boldsymbol{\eta} \sim \mathcal{N}(\mathbf{0}, \mathbf{H})$$

Results above give:

$$\hat{\mathbf{f}} = -\boldsymbol{\Sigma}_{\hat{\mathbf{f}}} \nabla \mathbf{I} \mathbf{H}^{-1} \mathbf{I}_t \quad \boldsymbol{\Sigma}_{\hat{\mathbf{f}}} = (\nabla \mathbf{I} \mathbf{H}^{-1} \nabla \mathbf{I}^T)^{-1}$$

Poor texture around $\mathbf{x} \implies \text{small } \nabla \mathbf{I} \implies \text{large } \boldsymbol{\Sigma}_{\hat{\mathbf{f}}}.$

Wang-Orquiza Results



Spatial Gaussian Processes

2-dimensional Gaussian process (distribution over functions of 2D inputs):

$$\mathbf{f}(\cdot) = \begin{bmatrix} u(\cdot) \\ v(\cdot) \end{bmatrix} \sim \mathcal{GP}(\mathbf{m}(\cdot), \mathbf{k}(\cdot, \cdot'))$$

Example mean & covariance kernel (affine & RBF):

$$\mathbf{m}(\cdot) = \mathbf{A}(\cdot) + \mathbf{b}, \quad \mathbf{k}(\cdot, \cdot') = \exp\left(-\frac{\|\cdot - \cdot'\|^2}{2\lambda^2}\right) \mathbf{\Sigma}$$

Joint distribution of function values at finite subset of points:

$$\begin{bmatrix} \mathbf{f}(\mathbf{x}_1) \\ \vdots \\ \mathbf{f}(\mathbf{x}_n) \end{bmatrix} \sim \mathcal{N}\left(\begin{bmatrix} \mathbf{m}(\mathbf{x}_1) \\ \vdots \\ \mathbf{m}(\mathbf{x}_n) \end{bmatrix}, \begin{bmatrix} \mathbf{k}(\mathbf{x}_1, \mathbf{x}_1) & \cdots & \mathbf{k}(\mathbf{x}_1, \mathbf{x}_n) \\ \vdots & \ddots & \vdots \\ \mathbf{k}(\mathbf{x}_n, \mathbf{x}_1) & \cdots & \mathbf{k}(\mathbf{x}_n, \mathbf{x}_n) \end{bmatrix}\right)$$

Equivalently:

$$\mathbf{F} := \begin{bmatrix} \mathbf{U}(\mathbf{X}) \\ \mathbf{V}(\mathbf{X}) \end{bmatrix} \sim \mathcal{N}\left(\boldsymbol{\mu} := \begin{bmatrix} \boldsymbol{\mu}_u \\ \boldsymbol{\mu}_v \end{bmatrix}, \mathbf{K} := \begin{bmatrix} \mathbf{K}_{uu} & \mathbf{K}_{uv} \\ \mathbf{K}_{vu} & \mathbf{K}_{vv} \end{bmatrix}\right)$$

Gaussian Process Regression

Through Wang-Orquiza method, we obtain noisy observations $\tilde{\mathbf{F}}$:

$$\tilde{\mathbf{f}}(\mathbf{x}) = \mathbf{f}(\mathbf{x}) + \boldsymbol{\eta}(\mathbf{x}), \quad \mathbf{H}(\mathbf{x}) \sim \mathcal{N}(\mathbf{0}, \boldsymbol{\Sigma}_{\tilde{\mathbf{f}}})$$

The noisy distribution is:

$$\tilde{\mathbf{F}} | \mathbf{F} \sim \mathcal{N}(\mathbf{F}, \boldsymbol{\Sigma}_{\tilde{\mathbf{f}}}), \quad \mathbf{F} \sim \mathcal{N}(\boldsymbol{\mu}, \mathbf{K}) \implies \tilde{\mathbf{F}} \sim \mathcal{N}(\boldsymbol{\mu}, \mathbf{K} + \boldsymbol{\Sigma}_{\tilde{\mathbf{f}}})$$

We want to estimate true optical flows: we form the joint distribution.

$$\begin{bmatrix} \mathbf{F} \\ \tilde{\mathbf{F}} \end{bmatrix} \sim \mathcal{N}\left(\begin{bmatrix} \boldsymbol{\mu} \\ \boldsymbol{\mu} \end{bmatrix}, \begin{bmatrix} \mathbf{K} & \mathbf{K} \\ \mathbf{K} & \mathbf{K} + \boldsymbol{\Sigma}_{\tilde{\mathbf{f}}} \end{bmatrix}\right)$$

We derive the conditional distribution (posterior):

$$\mathbf{F} | \tilde{\mathbf{F}} \sim \mathcal{N}\left(\boldsymbol{\mu} + \mathbf{K}(\mathbf{K} + \boldsymbol{\Sigma}_{\tilde{\mathbf{f}}})^{-1}(\tilde{\mathbf{F}} - \boldsymbol{\mu}), \mathbf{K} - \mathbf{K}(\mathbf{K} + \boldsymbol{\Sigma}_{\tilde{\mathbf{f}}})^{-1}\mathbf{K}\right)$$

Mean serves as estimate, covariance as uncertainty.

Parameter Fitting

Mean/covariance parameters (e.g. lengthscale λ^2) are optimized by maximizing the marginal likelihood of observations $\tilde{\mathbf{F}}$. This likelihood is:

$$p(\tilde{\mathbf{F}}) \propto \frac{1}{\sqrt{\det(\mathbf{K} + \boldsymbol{\Sigma}_{\tilde{\mathbf{f}}})}} \exp\left(-\frac{1}{2}(\tilde{\mathbf{F}} - \boldsymbol{\mu})^T (\mathbf{K} + \boldsymbol{\Sigma}_{\tilde{\mathbf{f}}})^{-1} (\tilde{\mathbf{F}} - \boldsymbol{\mu})\right)$$

In practice we minimize negative log-likelihood via gradient descent:

$$-\log p(\tilde{\mathbf{F}}) \propto \frac{1}{2}(\tilde{\mathbf{F}} - \boldsymbol{\mu})^T (\mathbf{K} + \boldsymbol{\Sigma}_{\tilde{\mathbf{f}}})^{-1} (\tilde{\mathbf{F}} - \boldsymbol{\mu}) + \frac{1}{2} \log \det(\mathbf{K} + \boldsymbol{\Sigma}_{\tilde{\mathbf{f}}})$$

We minimize this numerically via gradient descent.

Parameter Fitting (cont.)

Let φ be a parameter of \mathbf{m} , $\mathbf{y} = \tilde{\mathbf{F}} - \boldsymbol{\mu}$, $\mathbf{K}_y = \mathbf{K} + \sigma_\eta^2 \mathbf{I}$:

$$\frac{\partial \mathcal{L}}{\partial \varphi} = \frac{1}{2} \left[\left(\frac{\partial \mathbf{y}^T}{\partial \varphi} \right) \mathbf{K}_y^{-1} \mathbf{y} + \mathbf{y}^T \mathbf{K}_y \left(\frac{\partial \mathbf{y}}{\partial \varphi} \right) \right] = \frac{1}{2} \left(2 \mathbf{y}^T \mathbf{K}_y \left(\frac{\partial \mathbf{y}}{\partial \varphi} \right) \right) = -\mathbf{y}^T \mathbf{K}_y \left(\frac{\partial \mathbf{m}}{\partial \varphi} \right)$$


Let θ be a parameter of \mathbf{k} :

$$\frac{\partial \mathcal{L}}{\partial \theta} = \mathbf{y}^T \left(\frac{\partial \mathbf{K}_y^{-1}}{\partial \theta} \right) \mathbf{y} + \frac{\partial |\mathbf{K}_y|}{\partial \theta} = -\frac{1}{2} \mathbf{y}^T \mathbf{K}^{-1} \left(\frac{\partial \mathbf{K}_y}{\partial \theta} \right) \mathbf{K}^{-1} \mathbf{y} + \frac{1}{2} \text{tr} \left(\mathbf{K}_y^{-1} \left(\frac{\partial \mathbf{K}_y}{\partial \theta} \right) \right)$$

RBF parameter λ^2 :

$$\frac{\partial \mathbf{K}_y}{\partial \lambda} = \mathbf{K} \odot \frac{\|\mathbf{x} - \mathbf{x}'\|}{\lambda^3}$$

GP Regression Results



Discussion

Our maximum-likelihood method produced an initial estimate and uncertainty.

Our Gaussian-process Bayesian method provided updated these (posterior), after enforcing a global smoothness constraint (Gaussian process prior).

Possible next steps:

- ▶ Obtain better observations: FlowNet, RAFT, etc.
- ▶ Experiment with more complex covariance kernels.
- ▶ Perform segmentation, smooth each segment independently.
- ▶ Extend to spatiotemporal regression (3D, multiple frames).