Assignment 2 + Entropy, KL Divergence (SNLP Tutorial 3)

Vilém Zouhar, Awantee Deshpande, Julius Steuer

11th, 12th May 2021

Assignment 2

- Exercise 1: Perplexity Calculation
- Exercise 2: Formulating n-gram models
- Exercise 3: Perplexity Calculation for n-grams
- Bonus: Alternative metric to perplexity

- Information Content
- Entropy
- Joint entropy
- Conditional entropy
- Mutual Information
- Cross-entropy
- KL-Divergence

Concepts and formulations. (Express also as expectation values.)

Information Content

 $I(x) = -\log p(x)$

- Entropy
- Joint entropy
- Conditional entropy
- Mutual Information
- Cross-entropy
- KL-Divergence

- Information Content
- Entropy
- Joint entropy
- Conditional entropy
- Mutual Information
- Cross-entropy
- KL-Divergence

•
$$I(x) = -\log p(x)$$

•
$$H(X) = -\sum_{x \in X} p(x) \cdot \log p(x)$$

- Information Content
- Entropy
- Joint entropy
- Conditional entropy
- Mutual Information
- Cross-entropy
- KL-Divergence

•
$$I(x) = -\log p(x)$$

•
$$H(X) = -\sum_{x \in X} p(x) \cdot \log p(x)$$

•
$$H(X,Y) = -\sum_{x \in X} \sum_{y \in Y} p(x,y) \cdot \log p(x,y)$$

- Information Content
- Entropy
- Joint entropy
- Conditional entropy
- Mutual Information
- Cross-entropy
- KL-Divergence

•
$$I(x) = -\log p(x)$$

•
$$H(X) = -\sum_{x \in X} p(x) \cdot \log p(x)$$

•
$$H(X,Y) = -\sum_{x \in X} \sum_{y \in Y} p(x,y) \cdot \log p(x,y)$$

•
$$H(X,Y) = -\sum_{x \in X} \sum_{y \in Y} p(x,y) \cdot \log p(y \mid x)$$

- Information Content
- Entropy
- Joint entropy
- Conditional entropy
- Mutual Information
- Cross-entropy
- KL-Divergence

•
$$I(x) = -\log p(x)$$

•
$$H(X) = -\sum_{x \in X} p(x) \cdot \log p(x)$$

•
$$H(X,Y) = -\sum_{x \in X} \sum_{y \in Y} p(x,y) \cdot \log p(x,y)$$

•
$$H(X,Y) = -\sum_{x \in X} \sum_{y \in Y} p(x,y) \cdot \log p(y \mid x)$$

•
$$I(X;Y) = -\sum_{x,y} p(x,y) \cdot \log \frac{p(x,y)}{p(x) \cdot p(y)}$$

- Information Content
- Entropy
- Joint entropy
- Conditional entropy
- Mutual Information
- Cross-entropy
- KL-Divergence

•
$$I(x) = -\log p(x)$$

•
$$H(X) = -\sum_{x \in X} p(x) \cdot \log p(x)$$

•
$$H(X, Y) = -\sum_{x \in X} \sum_{y \in Y} p(x, y) \cdot \log p(x, y)$$

•
$$H(X,Y) = -\sum_{x \in X} \sum_{y \in Y} p(x,y) \cdot \log p(y \mid x)$$

•
$$I(X;Y) = -\sum_{x,y} p(x,y) \cdot \log \frac{p(x,y)}{p(x) \cdot p(y)}$$

•
$$H(p,q) = -\sum_{x} p(x) \cdot \log q(y)$$

- Information Content
- Entropy
- Joint entropy
- Conditional entropy
- Mutual Information
- Cross-entropy
- KL-Divergence

•
$$I(x) = -\log p(x)$$

•
$$H(X) = -\sum_{x \in X} p(x) \cdot \log p(x)$$

•
$$H(X,Y) = -\sum_{x \in X} \sum_{y \in Y} p(x,y) \cdot \log p(x,y)$$

•
$$H(X,Y) = -\sum_{x \in X} \sum_{y \in Y} p(x,y) \cdot \log p(y \mid x)$$

•
$$I(X;Y) = -\sum_{x,y} p(x,y) \cdot \log \frac{p(x,y)}{p(x) \cdot p(y)}$$

•
$$H(p,q) = -\sum_{x} p(x) \cdot \log q(y)$$

•
$$D(p||q) = -\sum_{x \in X} p(x) \cdot \log \frac{p(x)}{q(x)}$$

Chain Rule:

$$H(X, Y) = H(X) + H(Y|X)$$

$$H(X_1...X_n) = H(X_1) + H(X_2 \mid X_1) + ... + H(X_n \mid X_1, ..., X_{n-1})$$

Chain Rule:

$$H(X, Y) = H(X) + H(Y|X)$$

$$H(X_1...X_n) = H(X_1) + H(X_2 \mid X_1) + ... + H(X_n \mid X_1, ..., X_{n-1})$$

Mutual Information and Entropy

$$I(X; Y) = H(X) - H(X \mid Y) = H(X) + H(Y) - H(X, Y)$$

Chain Rule:

$$H(X, Y) = H(X) + H(Y|X)$$


$$H(X_1...X_n) = H(X_1) + H(X_2 \mid X_1) + ... + H(X_n \mid X_1, ..., X_{n-1})$$

Mutual Information and Entropy


$$I(X; Y) = H(X) - H(X \mid Y) = H(X) + H(Y) - H(X, Y)$$

Apply to 3 variables

$$I(X; Y \mid Z) = I((X; Y)|Z) = H(X \mid Z) - H(X \mid Y, Z)$$



 $Source: \ https://syncedreview.com/2020/11/30/synced-tradition-and-machine-learning-series-part-1-entropy/$



Source: https://en.wikipedia.org/wiki/Information_diagram

Examples - Entropy calculation

$X \setminus Y$	0	1
0	1/2	1/6
1	1/3	0

Find

- \bullet H(X), H(Y)
- \bullet H(X,Y)
- H(X|Y), H(Y|X)
- I(X; Y)
- H(Y) H(Y|X)

Examples - Entropy of functions

What is the (in)equality relationship between H(X) and H(Y) when

- y = f(x) # general case
- $y = 2^x$
- y = sin(x)

KL-divergence

Question: Can we use the chain rule on KL-Divergence?

KL-divergence

Question: Can we use the chain rule on KL-Divergence?

$$D(p(x,y) || q(x,y)) = D(p(x) || q(x)) + D(p(y | x) || q(y | x))$$

Applications of KL Divergence:

- Bayesian inference
- Compression techniques
- Variational autoencoders

Resources

- http://csustan.csustan.edu/~tom/sfi-csss/info-theory/info-lec.pdf
- https://www.cs.cmu.edu/~odonnell/toolkit13/lecture20.pdf