

MA 519: Homework 9

Max Jeter, Carlos Salinas

October 27, 2016

PROBLEM 9.1 (HANDOUT 13, # 7)

Let X have a *double exponential* density $f(x) = \frac{1}{2\sigma}e^{-\frac{|x|}{\sigma}}$, $-\infty < x < \infty$, $\sigma > 0$.

- (a) Show that all moments exist for this distribution.
- (b) However, show that the MGF exists only for restricted values. Identify them and find a formula.

SOLUTION. For part (a), we show that $E(X^n) < \infty$ for all $n \in \mathbb{N}$. By direct calculation, we have

$$\begin{aligned} E(X^n) &= \int_{-\infty}^{\infty} x^n f(x) dx \\ &= \int_{-\infty}^{\infty} \frac{x^n}{2\sigma} e^{-\frac{|x|}{\sigma}} dx \\ &= \underbrace{\int_{-\infty}^0 \frac{x^n}{2\sigma} e^{\frac{x}{\sigma}} dx}_L + \int_0^{\infty} \frac{x^n}{2\sigma} e^{-\frac{x}{\sigma}} dx \end{aligned}$$

making the substitution $x \mapsto -y$ to L and relabeling y to x again, we see that

$$\begin{aligned} &= \int_{-\infty}^0 \frac{x^n + (-1)x^n}{2\sigma} e^{-\frac{x}{\sigma}} dx \\ &= \begin{cases} 0 & \text{if } n \text{ is odd,} \\ I := \int_0^{\infty} \frac{x^n}{\sigma} e^{-\frac{x}{\sigma}} dx & \text{if } n = 2k \text{ is even.} \end{cases} \end{aligned}$$

To evaluate I we use integration by parts recursively to arrive at

$$\begin{aligned} I &= \int_{-\infty}^0 \frac{x^n}{\sigma} e^{-\frac{x}{\sigma}} \\ &= (-0 + 0) + \int_0^{\infty} n\sigma x^{n-1} e^{-\frac{x}{\sigma}} dx \\ &= (-0 + 0) + (-0 + 0) + \int_0^{\infty} n(n-1)\sigma^2 x^{n-2} e^{-\frac{x}{\sigma}} dx \\ &\vdots \\ &= (-0 - 0) + \cdots + (-0 + 0) + (-0 + n!\sigma^n) \\ &= n!\sigma^n. \end{aligned} \tag{9.1}$$

Therefore, $E(X^n) < \infty$ for all $n \in \mathbb{N}$, i.e., all moments of this distribution exist.

For part (b), the MGF associated to f is

$$m(t) = \sum_{i=0}^{\infty} \frac{t^i E(X^i)}{i!} = \sum_{i=0}^{\infty} t^i \sigma^i.$$

This is a geometric series and so converges for all $-\frac{1}{\sigma} < t < \frac{1}{\sigma}$. ■

PROBLEM 9.2 (HANDOUT 13, # 16)

Give an example of each of the following phenomena:

- (a) A continuous random variable taking values in $[0, 1]$ with equal mean and median.
- (b) A continuous random variable taking values in $[0, 1]$ with mean equal to twice the median.
- (c) A continuous random variable for which the mean does not exist.
- (d) A continuous random variable for which the mean exists, but the variance does not exist.
- (e) A continuous random variable with a PDF that is not differentiable at zero.
- (f) a positive continuous random variable for which the mode is zero, but the mean does not exist.
- (g) A continuous random variable for which all moments exist.
- (h) A continuous random variable with median equal to zero, and 25th and 75th percentiles equal to 1.
- (i) A continuous random variable X with mean equal to median equal to mode equal to zero, and $E(\sin X) = 0$.

SOLUTION. ■

PROBLEM 9.3 (HANDOUT 13, # 17)

An exponential random variable with mean 4 is known to be larger than 6. What is the probability that it is larger than 8?

SOLUTION.

■

PROBLEM 9.4 (HANDOUT 13, # 18)

(Sum of Gammas). Suppose X, Y are independent random variables, and $X \sim G(\alpha, \lambda)$, $Y \sim G(\beta, \lambda)$. Find the distribution of $X + Y$ by using moment-generating functions.

SOLUTION. ■

PROBLEM 9.5 (HANDOUT 13, # 19)

(*Product of Chi Squares*). Suppose X_1, X_2, \dots, X_n are independent chi square variables, with $X_i \sim \chi_{m_i}^2$. Find the mean and variance of $\prod_{i=1}^n X_i$.

SOLUTION. ■

PROBLEM 9.6 (HANDOUT 13, # 20)

Let $Z \sim N(0, 1)$. Find

$$P(0.5 < |Z - \tfrac{1}{2}| < 1.5); \quad P\left(\frac{e^Z}{1+e^Z} > \tfrac{3}{4}\right); \quad P(\Phi(Z) < 0.5).$$

SOLUTION. ■

PROBLEM 9.7 (HANDOUT 13, # 21)

Let $Z \sim N(0, 1)$. Find the density of $\frac{1}{Z}$. Is the density bounded?

SOLUTION. ■

PROBLEM 9.8 (HANDOUT 13, # 22)

The 25th and the 75th percentile of a normally distributed random variable are -1 and 1 . What is the probability that the random variable is between -2 and 2 ?

SOLUTION.

■