

MA 523: Homework 5

Carlos Salinas

October 4, 2016

PROBLEM 5.1

Prove that Laplace's equation $\Delta u = 0$ is rotation invariant; that is, if O is an orthogonal $n \times n$ matrix and we define $v(x) := u(Ox)$, $x \in \mathbb{R}^n$, then $\Delta v = 0$.

SOLUTION.

■

PROBLEM 5.2

Let $n = 2$ and U be the halfplane $\{x_2 > 0\}$. Prove that

$$\sup_U u = \sup_{\partial U} u$$

for $u \in C^2(U) \cap C(\bar{U})$ which are harmonic in U under the additional assumption that u is bounded from above in \bar{U} . (The additional assumption is needed to exclude examples like $u = x_2$.)

[Hint: Take for $\varepsilon > 0$ the harmonic function

$$u(x_1, x_2) + \varepsilon \ln \sqrt{x_1^2 + (x_2 + 1)^2}.$$

Apply the maximum principle to a region $\{x_1^2 + (x_2 + 1)^2 < a^2, x_2 > 0\}$ with large a . Let $\varepsilon \rightarrow 0$.]

SOLUTION. ■

PROBLEM 5.3

Let $U \subset \mathbb{R}^n$ be an open set. We say $v \in C^2(U)$ is subharmonic if

$$-\Delta v \leq 0 \quad \text{in } U.$$

- (a) Let $\varphi: \mathbb{R}^m \rightarrow \mathbb{R}$ be smooth and convex. Assume u^1, \dots, u^m are harmonic in U and

$$v := \varphi(u_1, \dots, u_m).$$

Prove v is sub harmonic.

[Hint: Convexity for a smooth function $\varphi(z)$ is equivalent to $\sum_{j,k=1}^m \varphi_{z_j, z_k}(z) \xi_j \xi_k \geq 0$ for any $\xi \in \mathbb{R}^m$.]

- (b) Prove $v := |Du|^2$ is subharmonic, whenever u is harmonic. (Assume that harmonic functions are C^∞ .)

SOLUTION. ■