

MA 519: Homework 9

Max Jeter, Carlos Salinas

October 27, 2016

PROBLEM 9.1 (HANDOUT 13, # 7)

Let X have a *double exponential* density $f(x) = \frac{1}{2\sigma}e^{-\frac{|x|}{\sigma}}$, $-\infty < x < \infty$, $\sigma > 0$.

- (a) Show that all moments exist for this distribution.
- (b) However, show that the MGF exists only for restricted values. Identify them and find a formula.

SOLUTION. For part (a), we show that the moments $m_n := E(X^n)$ for all $n \in \mathbb{N}$. By direct calculation, we have

$$\begin{aligned} m_n &= \int_{-\infty}^{\infty} x^n f(x) dx \\ &= \int_{-\infty}^{\infty} \frac{x^n}{2\sigma} e^{-\frac{|x|}{\sigma}} dx \\ &= \underbrace{\int_{-\infty}^0 \frac{x^n}{2\sigma} e^{\frac{x}{\sigma}} dx}_L + \int_0^{\infty} \frac{x^n}{2\sigma} e^{-\frac{x}{\sigma}} dx, \end{aligned}$$

making the substitution $x \mapsto -y$ to L and relabeling y to x again, the above becomes

$$\begin{aligned} &= \int_0^{\infty} \frac{x^n + (-1)x^n}{2\sigma} e^{-\frac{x}{\sigma}} dx \\ &= \begin{cases} 0 & \text{if } n \text{ is odd,} \\ I := \int_0^{\infty} \frac{x^n}{\sigma} e^{-\frac{x}{\sigma}} dx & \text{if } n = 2k \text{ is even.} \end{cases} \end{aligned}$$

To evaluate I we apply integration by parts repeatedly to arrive at

$$\begin{aligned} I &= \int_{-\infty}^0 \frac{x^n}{\sigma} e^{-\frac{x}{\sigma}} \\ &= (-0 + 0) + \int_0^{\infty} n\sigma x^{n-1} e^{-\frac{x}{\sigma}} dx \\ &= (-0 + 0) + (-0 + 0) + \int_0^{\infty} n(n-1)\sigma^2 x^{n-2} e^{-\frac{x}{\sigma}} dx & \vdots \\ &= (-0 - 0) + \cdots + (-0 + 0) + (-0 + n!\sigma^n) \\ &= n!\sigma^n. \end{aligned}$$

Therefore, m_n exist and are finite for all $n \in \mathbb{N}$.

For part (b), the MGF associated to f is given by the series

$$m(t) = \sum_{n=0}^{\infty} \frac{t^n m_n}{n!} = \sum_{k=1}^{\infty} t^{2k} \sigma^{2k}. \quad (9.1)$$

This series is geometric and, as such, converges for all $-\frac{1}{\sigma} < t < \frac{1}{\sigma}$, in which case (9.1) becomes

$$m(t) = \frac{1}{1 - t^2 \sigma^2}. \quad \blacksquare$$

PROBLEM 9.2 (HANDOUT 13, # 10)

Suppose X has Cauchy distribution as in # 6. Which of the following functions have finite expectation

$$X; \quad -X; \quad |X|; \quad \frac{1}{X}; \quad \sin X; \quad \ln |X|; \quad e^X; \quad e^{-|X|}?$$

SOLUTION. Suppose $X \sim \text{Cauchy}(0, 1)$. Then the PDF of X is given by the expression

$$f(x) = \frac{1}{\pi(x^2 + 1)}.$$

Now we proceed to find the expectations of (i) X , (ii) $-X$, (iii) $\frac{1}{X}$, (iv) $\sin X$, (v) $\ln |X|$, (vi) e^X , (vii) $e^{-|X|}$.

For (i), the expectation does not even exist. We repeat the argument given in class: Consider

$$E(X) = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{x}{x^2 + 1} dx = \lim_{x_1, x_2} \frac{1}{\pi} \int_{-x_1}^{x_2} \frac{x}{x^2 + 1} dx.$$

Then, making the substitution $u = x^2 + 1$, $du = 2x dx$, the integral above evaluates to

$$E(X) = \lim_{x_1, x_2 \rightarrow \infty} \frac{1}{2\pi} \ln \left(\frac{x_2^2 + 1}{x_1^2 + 1} \right).$$

However, the limit of this expression is undefined! Fix positive real number α and let $x_2 = \alpha x_1$. Then

$$E(X) = \lim_{x_1 \rightarrow \infty} \frac{1}{2\pi} \ln \left(\frac{\alpha^2 x_1^2 + 1}{x_1^2 + 1} \right) = \frac{1}{2\pi} \ln \alpha^2.$$

This value is distinct for each α . Therefore, the limit is not unique and so is undefined.

For (ii), the CDF of $-X$ is given by

$$\begin{aligned} F_{-X}(x) &= P(-X \leq x) \\ &= P(X \geq -x) \\ &= 1 - P(X < -x) \\ &= 1 - \int_{-\infty}^{-x} \frac{1}{\pi(y^2 + 1)} dy \\ &= \frac{1}{2} - \frac{1}{\pi} \tan^{-1}(-x). \end{aligned}$$

Thus, the PDF of $-X$ is

$$f_{-X}(x) = \frac{dF_{-X}(x)}{dx} = \frac{1}{\pi(x^2 + 1)}.$$

Thus, $-X \sim \text{Cauchy}(0, 1)$ and as we have previously shown, $E(X) = E(-X)$ is undefined.

For (iii), we have

$$X = \begin{cases} X & \text{if } X > 0, \\ -X & \text{if } X \leq 0. \end{cases}$$

Thus, the CDF of $|X|$ is

$$\begin{aligned}
 F_{|X|}(x) &= P(|X| \leq x) \\
 &= P(-x \leq X \leq x) \\
 &= \frac{1}{\pi} \int_{-x}^x \frac{1}{y^2 + 1} dy \\
 &= \frac{1}{\pi} (\tan^{-1}(x) - \tan^{-1}(-x)) \\
 &= \frac{2}{\pi} \tan^{-1}(x),
 \end{aligned}$$

and hence its PDF is

$$f_{|X|}(x) = \frac{dF_{|X|}(x)}{dx} = \begin{cases} \frac{2}{\pi(x^2+1)} & \text{if } x \geq 0, \\ 0 & \text{otherwise.} \end{cases}$$

Thus,

$$\begin{aligned}
 E(X) &= \int_0^\infty \frac{2x}{x^2 + 1} dx \\
 &= \lim_{x \rightarrow \infty} \ln(x^2 + 1) \\
 &= \infty.
 \end{aligned}$$

For (iv), as discussed in class $\frac{1}{X} \sim \text{Cauchy}(0, 1)$. Let us show this. First, we find the CDF of X :

$$\begin{aligned}
 F_{\frac{1}{X}}(x) &= P\left(\frac{1}{X} \leq x\right) \\
 &= P\left(X \geq \frac{1}{x}\right) \\
 &= 1 - P\left(X < \frac{1}{x}\right) \\
 &= 1 - \frac{1}{\pi} \int_{-\infty}^{\frac{1}{x}} \frac{1}{y^2 + 1} dy \\
 &= 1 - \tan^{-1}\left(\frac{1}{x}\right) - \frac{1}{2} \\
 &= \frac{1}{2} - \tan^{-1}\left(\frac{1}{x}\right).
 \end{aligned}$$

Thus, the PDF of $\frac{1}{X}$ is

$$\begin{aligned}
 f_{\frac{1}{X}}(x) &= \frac{dF_{\frac{1}{X}}(x)}{dx} \\
 &= -\left(-\frac{1}{x^2}\right) \left(\frac{1}{\left(\frac{1}{x}\right)^2 + 1}\right) \\
 &= \frac{1}{x^2 + 1}.
 \end{aligned}$$

Thus, $\frac{1}{X} \sim \text{Cauchy}(0, 1)$ so its expectation is not defined.

For (v), the CDF of $\sin X$ is given by

$$F_{\sin X}(x) = P(\sin X \leq x),$$

for $-1 \leq x \leq 1$, we have

$$\begin{aligned} &= \sum_{-\infty < n < \infty} P((2n+1)\pi - \sin^{-1} x \leq X \leq (2n+2)\pi + \sin^{-1} x) \\ &= \frac{1}{\pi} \sum_{-\infty < n < \infty} \int_{(2n+1)\pi - \sin^{-1} x}^{(2n+2)\pi + \sin^{-1} x} \frac{1}{y^2 + 1} dy \\ &= \frac{1}{\pi} \sum_{-\infty < n < \infty} \tan^{-1}((2n+2)\pi + \sin^{-1} x) + \tan^{-1}((2n+1)\pi - \sin^{-1} x). \end{aligned}$$

Thus, the PDF of $\sin X$ is ■

PROBLEM 9.3 (HANDOUT 13, # 16)

Give an example of each of the following phenomena:

- (a) A continuous random variable taking values in $[0, 1]$ with equal mean and median.
- (b) A continuous random variable taking values in $[0, 1]$ with mean equal to twice the median.
- (c) A continuous random variable for which the mean does not exist.
- (d) A continuous random variable for which the mean exists, but the variance does not exist.
- (e) A continuous random variable with a PDF that is not differentiable at zero.
- (f) a positive continuous random variable for which the mode is zero, but the mean does not exist.
- (g) A continuous random variable for which all moments exist.
- (h) A continuous random variable with median equal to zero, and 25th and 75th percentiles equal to 1.
- (i) A continuous random variable X with mean equal to median equal to mode equal to zero, and $E(\sin X) = 0$.

SOLUTION. First, note that $[0, 1]$ is a probability space under the standard Lebesgue measure on \mathbb{R} . Therefore, it makes sense to consider $X: [0, 1] \rightarrow \mathbb{R}$ random variables.

For part (a), consider the random variable $X: [0, 1] \rightarrow \mathbb{R}$ defined by $x \mapsto x$ with $X \sim \text{Uniform}[0, 1]$. Then the mean is

$$\mu = \int_{-\infty}^{\infty} xf(x) dx = \int_0^1 x dx = \frac{1}{2}$$

and the median is

$$m = \inf\{x : F(x) = x \geq 0.5\} = \frac{1}{2}.$$

For part (b), consider again the random variable $X(x) = x$ for $x \in [0, 1]$, but this time let

$$f(x) = \begin{cases} & \\ & \end{cases},$$

be the PDF of X . Then the mean is ■

PROBLEM 9.4 (HANDOUT 13, # 17)

An exponential random variable with mean 4 is known to be larger than 6. What is the probability that it is larger than 8?

SOLUTION.

■

PROBLEM 9.5 (HANDOUT 13, # 18)

(Sum of Gammas). Suppose X, Y are independent random variables, and $X \sim \Gamma(\alpha, \lambda)$, $Y \sim \Gamma(\beta, \lambda)$. Find the distribution of $X + Y$ by using moment-generating functions.

SOLUTION. ■

PROBLEM 9.6 (HANDOUT 13, # 19)

(*Product of Chi Squares*). Suppose X_1, X_2, \dots, X_n are independent chi square variables, with $X_i \sim \chi_{m_i}^2$. Find the mean and variance of $\prod_{i=1}^n X_i$.

SOLUTION. ■

PROBLEM 9.7 (HANDOUT 13, # 20)

Let $Z \sim \text{Normal}(0, 1)$. Find

$$P\left(0.5 < \left|Z - \frac{1}{2}\right| < 1.5\right); \quad P\left(\frac{e^Z}{1 + e^Z} > \frac{3}{4}\right); \quad P(\Phi(Z) < 0.5).$$

SOLUTION.

■

PROBLEM 9.8 (HANDOUT 13, # 21)

Let $Z \sim \text{Normal}(0, 1)$. Find the density of $\frac{1}{Z}$. Is the density bounded?

SOLUTION. ■

PROBLEM 9.9 (HANDOUT 13, # 22)

The 25th and the 75th percentile of a normally distributed random variable are -1 and 1 . What is the probability that the random variable is between -2 and 2 ?

SOLUTION.

