

MA 523: Homework 6

Carlos Salinas

October 12, 2016

PROBLEM 6.1

For $n = 2$ find Green's function for the quadrant $\{x_1 > 0, x_2 > 0\}$ by repeated reflection.

Solution.

■

PROBLEM 6.2

(Precise form of Harnack's inequality) Use Poisson's formula for the ball to prove

$$\frac{r^{n-2}(r-|x|)}{(r+|x|)^{n-1}}u(0) \leq u(x) \leq \frac{r^{n-2}(r+|x|)}{(r-|x|)^{n-1}}u(0)$$

whenever u is positive and harmonic in $B(0, r) = \{x \in \mathbb{R}^n : |x| < r\}$.

Solution.

■

PROBLEM 6.3

Let $P_k(x)$ and $P_m(x)$ be homogeneous harmonic polynomials in \mathbb{R}^n of degrees k and m respectively; i.e.,

$$\begin{aligned} P_k(\lambda x) &= \lambda^k P_k(x), & P_m(\lambda x) &= \lambda^m P_m(x) & \text{for every } x \in \mathbb{R}^n, \lambda > 0, \\ \Delta P_k &= 0, & \Delta P_m &= 0 & \text{in } \mathbb{R}^n. \end{aligned}$$

(a) Show that

$$\frac{\partial P_k}{\partial \nu} = k P_k(x), \quad \frac{\partial P_m}{\partial \nu} = m P_m(x) \quad \text{on } \partial B(0, 1)$$

where $B(0, 1) = \{x \in \mathbb{R}^n : |x| < 1\}$ and ν is the outward normal on $\partial B(0, 1)$.

(b) Use (a) and Green's formula to prove that

$$\int_{\partial B(0, 1)} P_k(x) P_m(x) d\sigma = 0, \quad \text{if } k \neq m.$$

Solution.

■