

## MA 523: Homework 2

Carlos Salinas

September 9, 2016



**Problem 2.1**

Verify assertion (36) in [E, §3.2.3], that when  $\Gamma$  is not flat near  $x^0$  the noncharacteristic condition is

$$D_p F(p^0, z^0, x^0) \cdot \nu(x^0) \neq 0.$$

(Here  $\nu(x^0)$  denotes the normal to the hypersurface  $\Gamma$  at  $x^0$ ).

**Solution.** ► First, note that the condition

$$D_p F(p^0, z^0, x^0) \cdot \nu(x^0) \neq 0 \tag{2.1}$$

reduces to the standard noncharacteristic boundary condition if  $\Gamma$  is flat near  $x^0$  because in such case we have  $\nu(x^0) = (0, \dots, 0, 1)$  so

$$\begin{aligned} 0 &\neq D_p F(p^0, z^0, x^0) \cdot (0, \dots, 0, 1) \\ &= F_{p_n}(p^0, z^0, x^0). \end{aligned}$$

To show (2.1), we will first straighten the boundary near  $x^0$  and then apply the noncharacteristic boundary conditions to the straightened region. Write  $y = \Phi(x)$  where

$$\begin{cases} \Phi^1(x) = x_1, \\ \vdots \\ \Phi^n(x) = x_n - \varphi(x_1, \dots, x_{n-1}), \end{cases}$$

with  $\varphi(x_1^0, \dots, x_{n-1}^0) = x_n^0$  and let  $\Psi = \Phi^{-1}$  and  $v(y) = u(\Phi(x))$ . Then the image of  $\Gamma$  under  $\Phi$  is flat near  $y^0 = \Phi(x^0) = (x_1^0, \dots, x_{n-1}^0, 0)$  so we can apply the standard noncharacteristic boundary conditions on the PDE,

$$0 \neq G_{p_n}(p, z, y), \tag{2.2}$$

where  $G$  is the PDE  $F$  after applying the transformation  $\Phi$ , i.e., the PDE

$$G(Dv, v, y) = F(Dv(y)D\Phi(\Psi(y)), v(y), \Psi(y)).$$

We are done after we relate (2.2) to the original PDE  $F$ . Note that by the equation above we have

$$p(y)D\Phi(\Psi(y)) = \begin{cases} p_1(y) - p_n(y)\varphi_{x_1}(y_1, \dots, y_{n-1}), \\ \vdots \\ p_{n-1} - p_n(y)\varphi_{x_{n-1}}(y_1, \dots, y_{n-1}), \\ p_n(y). \end{cases}$$

Thus,

$$G_{p_n}(p, z, \tilde{y}) = F_{p_n}(pD\Phi(\Psi(\tilde{y})), z, \Psi(\tilde{y}))$$

which, by the chain rule, is equal to

$$\begin{aligned} &= -F_{p_1} \varphi_{x_1}(\tilde{y}) - \cdots - F_{p_{n-1}} \varphi_{x_{n-1}} + F_{p_n} \\ &= DF(p^0, z^0, x^0) \cdot (-D\varphi(x_1^0, \dots, x_{n-1}^0), 1) \\ &= DF(p^0, z^0, x^0) \cdot v(x^0) \end{aligned}$$

◀

**Problem 2.2**

Show that the solution of the quasilinear PDE

$$u_t + a(u)u_x = 0$$

with initial conditions  $u(x, 0) = g(x)$  is given implicitly by

$$u = g(x - a(u)t).$$

Show that the solution develops a shock (becomes singular) for some  $t > 0$ , unless  $a(g(x))$  is a nondecreasing function of  $x$ .

**Solution.** ►

◀

**Problem 2.3**

Show that the function  $u(x, t)$  defined by  $t \geq 0$  by

$$u(x, t) = \begin{cases} -\frac{2}{3} \left( t + \sqrt{3x + t^2} \right) & \text{for } 4x + t^2 > 0 \\ 0 & \text{for } 4x + t^2 < 0 \end{cases}$$

is an (unbounded) entropy solution of the conservation law  $u_t + (u^2/2)_x = 0$  (*inviscid Burger's equation*).

**Solution.** ►

◀