

## MA 523: Homework 2

Carlos Salinas

September 9, 2016



**Problem 2.1**

Verify assertion (36) in [E, §3.2.3], that when  $\Gamma$  is not flat near  $x^0$  the noncharacteristic condition is

$$D_p F(p^0, z^0, x^0) \cdot \nu(x^0) \neq 0.$$

(Here  $\nu(x^0)$  denotes the normal to the hypersurface  $\Gamma$  at  $x^0$ ).

**Solution.** ► First, note that the condition

$$D_p F(p^0, z^0, x^0) \cdot \nu(x^0) \neq 0 \tag{2.1}$$

reduces to the standard noncharacteristic boundary condition if  $\Gamma$  is flat near  $x^0$  because in such case we have  $\nu(x^0) = (0, \dots, 0, 1)$  so

$$\begin{aligned} 0 &\neq D_p F(p^0, z^0, x^0) \cdot (0, \dots, 0, 1) \\ &= F_{p_n}(p^0, z^0, x^0). \end{aligned}$$

We shall verify the noncharacteristic condition (2.1) by first flattening the boundary near  $x^0$  and then applying the noncharacteristic boundary conditions to the flattened region. Assuming some degree of regularity near  $x^0$ , e.g., that the boundary of  $U$  be smooth, we may express  $\Gamma$  near  $x^0$  as the graph of a smooth function  $f: \mathbb{R}^{n-1} \rightarrow \mathbb{R}$ , i.e.,  $x = (x_1, \dots, x_{n-1}, f(x_1, \dots, x_{n-1}))$  on  $\Gamma$  and  $x_n \geq f(y)$  after reorienting the coordinate axes. Then we flatten out  $\Gamma$  via the map  $\Phi(x): \mathbb{R}^n \rightarrow \mathbb{R}^n$  given by

$$\begin{cases} y_1 = x_1 = \Phi^1(x), \\ \vdots \\ y_{n-1} = x_{n-1} = \Phi^{n-1}(x), \\ y_n = x_n - f(x_1, \dots, x_{n-1}) = \Phi^n(x) \end{cases}$$

and write  $y = \Phi(x)$ . Let  $\Psi = \Phi^{-1}$  and rewrite our PDE  $F$  in terms of  $y$  as follows,

$$0 = F(Du(\Psi(y)), u(\Psi(y)), \Psi(y)). \tag{2.2}$$

Since  $\Delta = \Phi(\Gamma)$  is flat near  $y^0 = \Phi(x^0) = (y_1^0, \dots, y_{n-1}^0, 0)$ , we may apply the standard noncharacteristic condition on (2.2) and get

$$0 \neq F_{u_{y_n}}(Du(\Psi(y^0)), u(\Psi(y^0)), \Psi(y^0)).$$

Before we move on to finding an expression for this derivative, let us consider the gradient  $Du(\Psi(y))$ . By the chain rule, we have

$$\begin{aligned} u_{y_i}(\Psi(y)) &= \sum_{j=1}^n u_{x_j}(\Psi(y)) \frac{\partial x_j}{\partial y_i} \\ &= u_{x_i}(\Psi(y)) + u_{x_n}(\Psi(y)) f_{y_i}(y_1, \dots, y_{n-1}), \\ u_{y_n}(\Psi(y)) &= \sum_{j=1}^n u_{x_j}(\Psi(y)) \frac{\partial x_j}{\partial y_n} \\ &= u_{x_n}(\Psi(y)), \end{aligned}$$

Then, substituting  $u_{y_n}$  for  $u_{x_n}$ , we have

$$u_{y_i}(\Psi(y)) = u_{x_i}(\Psi(y)) + u_{y_n}(\Psi(y))f_{y_i}(y_1, \dots, y_{n-1}),$$

Now, by the chain rule on (2.2), we have

$$\begin{aligned} 0 &\neq F_{u_{y_n}}(Du(\Psi(y^0)), u(\Psi(y^0)), \Psi(y^0)) \\ &= F_{u_{y_n}}(u_{x_1} + u_{y_n}f_{y_1}, \dots, u_{x_{n-1}} + u_{y_n}f_{y_{n-1}}, u_{y_n}, z^0, x^0) \\ &= F_{u_{x_1}}f_{y_1} + \dots + F_{u_{x_{n-1}}}f_{y_{n-1}} + F_{u_{x_n}} \\ &= D_p F(p^0, z^0, x^0) \cdot (Df(x^0), 1) \\ &= D_p F(p^0, z^0, x^0) \cdot v(x^0), \end{aligned}$$

as we set out to show. ◀

**Problem 2.2**

Show that the solution of the quasilinear PDE

$$u_t + a(u)u_x = 0$$

with initial conditions  $u(x, 0) = g(x)$  is given implicitly by

$$u = g(x - a(u)t).$$

Show that the solution develops a shock (becomes singular) for some  $t > 0$ , unless  $a(g(x))$  is a nondecreasing function of  $x$ .

**Solution.** ► Using the method of characteristics, the characteristic ODEs are

$$\dot{t} = 1, \quad \dot{x} = a(z), \quad \dot{z} = 1.$$

◀

**Problem 2.3**

Show that the function  $u(x, t)$  defined by  $t \geq 0$  by

$$u(x, t) = \begin{cases} -\frac{2}{3} \left( t + \sqrt{3x + t^2} \right) & \text{for } 4x + t^2 > 0 \\ 0 & \text{for } 4x + t^2 < 0 \end{cases}$$

is an (unbounded) entropy solution of the conservation law  $u_t + (u^2/2)_x = 0$  (*inviscid Burger's equation*).

**Solution.** ►

◀