

# MA 519: Homework 6

Max Jeter, Carlos Salinas

October 6, 2016



## PROBLEM 6.1 (HANDOUT 8, # 2)

Identify the parameters  $n$  and  $p$  for each of the following binomial distributions:

- (a) # boys in a family with 5 children;
- (b) # correct answers in a multiple choice test if each question has a 5 alternatives, there are 25 questions, and the student is making guesses at random.

*SOLUTION.* For part (a), the distribution is binomial with  $k$  being the number of children in a given family and  $p$  the probability that a child is born, say, male. In this case, we can reasonably assume that  $p = 0.5$ . Thus, the binomial distribution is given by  $\text{Binom}(5, 0.5)$ .

For part (b), we use similar reasoning and we have  $\text{Binom}(25, 0.2)$  where  $k = 25$  is the number of questions and  $p = 1/5 = 0.2$  the probability of guessing a question correctly. ■

## PROBLEM 6.2 (HANDOUT 8, # 10)

A newsboy purchases papers at 20 cents and sells them for 35 cents. He cannot return unsold papers. If the daily demand for papers is modeled as a  $\text{Binom}(50, 0.5)$  random variable, what is the optimum number of papers the newsboy should purchase?

*SOLUTION.* Let  $X \sim \text{Binom}(50, 0.5)$  denote the daily demand for papers and  $n$  the number of copies bought by the newsboy. Then, the random variable  $S = \min\{X, n\}$  denotes the number of copies actually sold by the newsboy. His daily profit is, therefore, measured by the random variable

$$Y = 0.35S - 0.25n.$$

Now let us compute the average sales of the newsboy. By the linearity of expected value, we have

$$\begin{aligned} E(S) &= \sum_{k=0}^n kP(S = k) \\ &= \sum_{k=0}^{n-1} kP(\min\{X, n\} = k) + nP(\min\{X, n\} = n), \end{aligned}$$

where  $P(\min\{X, n\} = k) = P(X = k)$  the probability that there is a demand for  $k$  copies, and  $P(\min\{X, n\} = n) = P(X \geq n)$  the probability that the demand exceeds the number of copies the newsboy bought, giving us

$$\begin{aligned} &= \sum_{k=0}^{n-1} kP(X = k) + nP(X \geq n) \\ &= \sum_{k=0}^{n-1} kP(X = k) + n(1 - P(X < n)) \\ &= n + \sum_{k=0}^{n-1} kP(X = k) - nP(X < n) \\ &= n + \sum_{k=0}^{n-1} kP(X = k) - n \sum_{k=0}^{n-1} P(X = k) \\ &= n + \sum_{k=0}^{n-1} k \binom{n-1}{k} 0.5^k 0.5^{n-1-k} - n \sum_{k=0}^{n-1} \binom{n-1}{k} 0.5^k 0.5^{n-1-k} \\ &= n + \sum_{k=0}^{n-1} k \binom{n-1}{k} 0.5^{n-1} - n \sum_{k=0}^{n-1} \binom{n-1}{k} 0.5^{n-1} \\ &= n + (n-1)0.5 - n0.5^{n-1}2^{n-1} \\ &= \frac{n-1}{2}. \end{aligned}$$

Thus,

$$E(Y) = 0.35E(S) - 0.25n = 0.175(n-1) - 0.25n =$$

■

## PROBLEM 6.3 (HANDOUT 8, # 12)

How many independent bridge dealings are required in order for the probability of a preassigned player having four aces at least once to be  $1/2$  or better? Solve again for some player instead of a given one.

*SOLUTION.* Without loss of generality, assume North is the preassigned player. Let  $X$  be the number of dealings required for North to have a probability greater than or equal to  $1/2$  ■

## PROBLEM 6.4 (HANDOUT 8, # 13)

A book of 500 pages contains 500 misprints. Estimate the chances that a given page contains at least three misprints.

*SOLUTION.* Let  $X$  be the number of misprints on the given page. The probability that a given misprint is on that page is  $1/500$ . Now,  $P(X \geq 3) = 1 - P(X = 0) - P(X = 1) - P(X = 2)$ . Also,

$$\begin{aligned}P(X = 0) &= \left(\frac{499}{500}\right)^{500} \\P(X = 1) &= 500 \left(\frac{1}{500}\right) \left(\frac{499}{500}\right)^{499} \\P(X = 2) &= \frac{500 \cdot 499}{2} \left(\frac{1}{500}\right)^2 \left(\frac{499}{500}\right)^{498}\end{aligned}$$

So that

$$\begin{aligned}P(X \geq 3) &= 1 - P(X = 0) - P(X = 1) - P(X = 2) \\&= 1 - \left(\frac{499}{500}\right)^{500} - 500 \left(\frac{1}{500}\right) \left(\frac{499}{500}\right)^{499} - \frac{500 \cdot 499}{2} \left(\frac{1}{500}\right)^2 \left(\frac{499}{500}\right)^{498} \\&\approx 0.08\end{aligned}$$

that is, the probability that the given page has at least 3 misprints is about 8 percent. ■

## PROBLEM 6.5 (HANDOUT 8, # 14)

Colorblindness appears in 1 per cent of the people in a certain population. How large must a random sample (with replacements) be if the probability of its containing a colorblind person is to be 0.95 or more?

*SOLUTION.* Let  $n$  be the sample size. The probability of the sample containing no colorblind people is  $0.99^n$ . Solving the equation  $0.99^n = 0.05$ , we see that (for the naturals) taking  $n = 299$  is (minimally) sufficient for  $0.99^n$  to be less than 0.05.

The probability that the sample has some colorblind person is equal to  $1 - 0.99^n$ . This is at least 95 percent if  $0.99^n$  is less than 0.05. That is, having 299 people in the sample is (minimally) sufficient for there to be a 95 percent chance of having some colorblind person. ■



PROBLEM 6.6 (HANDOUT 8, # 15)

Two people toss a true coin  $n$  times each. Find the probability that they will score the same number of heads.

SOLUTION. ■

## PROBLEM 6.7 (HANDOUT 8, # 16)

Binomial approximation to the hypergeometric distribution. A population of TV elements is divided into red and black elements in the proportion  $p : q$  (where  $p + q = 1$ ). A sample of size  $n$  is taken without replacement. The probability that it contains exactly  $k$  red elements is given by the hypergeometric distribution of II, 6. Show that as  $n \rightarrow \infty$  this probability approaches  $\text{Binom}(n, p)$ .

SOLUTION. ■

## PROBLEM 6.8 (HANDOUT 9, # 3)

Suppose  $X, Y, Z$  are mutually independent random variables, and  $E(X) = 0, E(Y) = -1, E(Z) = 1, E(X^2) = 4, E(Y^2) = 3, E(Z^2) = 10$ . Find the variance and the second moment of  $2Z - Y/2 + eX$ , where  $e$  is the number such that  $\ln e = 1$ .

*SOLUTION.* Let  $W = 2Z - Y/2 + eX$ . Then

$$\begin{aligned}
 E(W^2) &= E((2Z - Y/2 + eX)^2) \\
 &= E(4Z^2 - 2ZY + 4eZX + Y^2/4 - eYX + e^2X^2) \\
 &= 4E(Z^2) - 2E(Z)E(Y) + 4eE(Z)E(X) + E(Y^2)/4 - eE(Y)E(X) + e^2E(X^2) \\
 &= 4 \cdot 10 + 2 + 3/4 + e^2 \cdot 4 \\
 &= \frac{171}{4} + 4e^2 \\
 &\approx 72.31
 \end{aligned}$$

and

$$\begin{aligned}
 \text{Var}(W) &= E(W^2) - E(W)^2 \\
 &= \frac{171}{4} + 4e^2 - (2E(Z) + E(Y)/2 + eE(X))^2 \\
 &= \frac{171}{4} + 4e^2 - (2 - 1/2)^2 \\
 &= \frac{171}{4} + 4e^2 - (3/2)^2 \\
 &= \frac{81}{2} + 4e^2 \\
 &\approx 70.06
 \end{aligned}$$

That is, the second moment is “about 72.31” and the variance is “about 70.06”

■

## PROBLEM 6.9 (HANDOUT 9, # 14)

(*Variance of Product*). Suppose  $X, Y$  are independent random variables. Can it ever be true that  $\text{Var}(XY) = \text{Var}(X) \text{Var}(Y)$ ?

If it can, when?

SOLUTION. ■