

MA 519: Homework 11

Max Jeter, Carlos Salinas

November 10, 2016

PROBLEM 11.1 (DASGUPTA 7.2 (A), (B), (C), (D), (E))

- (a) Suppose $E|X_n - c|^\alpha \rightarrow 0$, where $0 < \alpha < 1$. Does X_n necessarily converge in probability to c ?
- (b) Suppose $a_n(X_n - \theta) \xrightarrow{\mathcal{L}} N(0, 1)$. Under what condition on a_n can we conclude that $X_n \xrightarrow{\mathcal{P}} \theta$?
- (c) $o_p(1) + O_p(1) = ?$
- (d) $o_p(1)O_p(1) = ?$
- (e) $o_p(1) + o_p(1)O_p(1) = ?$

SOLUTION. For part (a) we show that indeed $E(|X_n - c|^\alpha) \rightarrow 0$ implies $X_n \xrightarrow{\mathcal{P}} c$. Let $\varepsilon > 0$ be given. By Markov's inequality, we have

$$P(|X_n - c| > \varepsilon) = P(|X_n - c|^\alpha > \varepsilon^\alpha) \leq \frac{E(|X_n - c|^\alpha)}{\varepsilon^\alpha}.$$

Since $E(|X_n - c|^\alpha) \rightarrow 0$ as $n \rightarrow \infty$, it follows that $P(|X_n - c| > \varepsilon) \rightarrow 0$ as $n \rightarrow \infty$; i.e., X_n converges to c in probability.

For part (b), suppose $a_n(X_n - \theta) \xrightarrow{\mathcal{L}} N(0, 1)$; i.e., $P(|a_n(X_n - \theta)| \leq x) \rightarrow \Phi(x)$ as $n \rightarrow \infty$. In words $X_n \xrightarrow{\mathcal{P}} \theta$ means that for every $\varepsilon > 0$ and every $\eta > 0$ there exists a positive integer N depending on ε and η such that $n \geq N$ implies

$$P(|X_n - \theta| \geq \varepsilon) < \eta.$$

First, let us find the PDF of the sequence $a_n(X_n - \theta)$. Let f_n denote the PDF of X_n , then the CDF of $a_n(X_n - \theta)$ is

$$\begin{aligned} P(|a_n(X_n - \theta)| \leq x) &= P(-x \leq a_n(X_n - \theta) \leq x) \\ &= P\left(-\frac{x}{a_n} + \theta \leq X_n \leq \frac{x}{a_n} + \theta\right) \\ &= \int_{-x/a_n + \theta}^{x/a_n + \theta} f(y) dy \\ &= f(x/a_n + \theta) - f(-x/a_n + \theta), \end{aligned}$$

therefore its PDF is

$$\begin{aligned} \frac{dP}{dx}(|a_n(X_n - \theta)| \leq x) &= \frac{d}{dx} \left[f\left(\frac{x}{a_n} + \theta\right) - f\left(-\frac{x}{a_n} + \theta\right) \right] \\ &= \frac{1}{a_n} (f(x/a_n + \theta) + f(-x/a_n + \theta)) \end{aligned}$$

For part (c), suppose $\{a_n\}$ and $\{b_n\}$ are sequences such that $a_n = o_p(1)$ and $b_n = O_p(1)$, then for the sequence $\{c_n := a_n + b_n\}$ the most we can expect is $c_n = O_p(1)$. Indeed, we know that if a sequence is $o_p(1)$ then it is also $O_p(1)$ therefore there exists K_1 and K_2 such that $|a_n| \leq K_1$, $|b_n| \leq K_2$ for all $n \geq 1$. Therefore, $|c_n| \leq K_1 + K_2$ for all $n \geq 1$.

For part (d), suppose $\{a_n\}$ and $\{b_n\}$ are sequences such that $a_n = o_p(1)$ and $b_n = O_p(1)$, then for the sequence $\{c_n := a_n b_n\}$ the most we can expect is $c_n = O_p(1)$. Again, since $\{a_n\}$ is $o_p(1)$ it

is $O_p(1)$ so there exists a constant $K_1 \geq 0$ such that $|a_n| \leq K_1$ for all $n \geq 1$ and similarly for $\{b_n\}$ there exists a constant K_2 such that $|b_n| \leq K_2$ for all $n \geq 1$. Therefore, $|c_n| \leq K_1 K_2$ for all $n \geq 1$ so $c_n = O_p(1)$.

For part (e), suppose $\{a_n\}$, $\{b_n\}$, and $\{c_n\}$ are sequences such that $a_n, b_n = o_p(1)$ and $c_n = O_p(1)$, then for the sequence $\{d_n := a_n + b_n c_n\}$ the most we can expect is $d_n = O_p(1)$ since there exists constants K_1 , K_2 , and K_3 such that $|a_n| \leq K_1$, $|b_n| \leq K_2$, and $|c_n| \leq K_3$ for all $n \geq 1$. This implies that $|d_n| \leq K_1 + K_2 K_3$ for all $n \geq 1$. Thus, $d_n = O_p(1)$. ■

PROBLEM 11.2 (DASGUPTA 7.3 [MONTE CARLO])

Consider the purely mathematical problem of finding a definite integral $\int f(x) dx$ for some (possibly complicated) function $f(x)$. Show that the SLLN provides a method for approximately finding the value of the integral by using appropriate averages $\frac{1}{n} \sum_{k=1}^n f(X_k)$.

Numerical analysts call this Monte Carlo integration.

SOLUTION.

■

PROBLEM 11.3 (DASGUPTA 7.4 (A), (B))

Suppose X_1, \dots , are i.i.d. and that $E(X_1) = \mu \neq 0$, $\text{Var}(X_1) = \sigma^2 < \infty$. Let $S_{m,p} = \sum_{k=1}^m X_k^p$, $m \geq 1$, $p = 1, 2$.

- (a) Identify with proof the almost sure limit of $S_{m,1}/S_{n,1}$ for fixed m , and $n \rightarrow \infty$.
- (b) Identify with proof the almost sure limit of $S_{n-m,1}/S_{n,1}$ for fixed m , and $n \rightarrow \infty$.

SOLUTION. ■

PROBLEM 11.4 (DASGUPTA 7.5 (A))

Let A_n , $n \geq 1$, A be events with respect to a common sample space Ω .

- (a) Prove that $I_{A_n} \xrightarrow{\mathcal{L}} I_A$ if and only if $P(A_n) \rightarrow P(A)$.

SOLUTION. ■

PROBLEM 11.5 (DASGUPTA 7.11 [SAMPLE MAXIMUM])

Let X_k , $k \geq 1$, be an i.i.d. sequence, and $X_{(n)}$ the maximum of X_1, \dots, X_n . Let $\xi(F) = \sup\{x : F(x) < 1\}$, where F is the common CDF of the X_k . Prove that $X_{(n)} \xrightarrow{\text{a.s.}} \xi(F)$.

SOLUTION. ■

PROBLEM 11.6 (DASGUPTA 7.14 (A))

Suppose X_k are i.i.d. standard Cauchy. Show that

- (a) $P(|X_n| > n \text{ infinitely often}) = 1$.

SOLUTION.

■

PROBLEM 11.7 (DASGUPTA 7.16 [COUPON COLLECTION])

Cereal boxes contain independently and with equal probability exactly one of n different celebrity pictures. Someone having the entire set of n pictures can cash them in for money. Let W_n be the minimum number of cereal boxes one would need to purchase to own a complete set of the pictures. Find a sequence a_n such that $W_n/a_n \xrightarrow{\mathcal{P}} 1$.
(*Hint:* Approximate the mean of W_n .)

SOLUTION. Let $X_n \sim \text{Geom}(\frac{n-k}{n})$

■

PROBLEM 11.8 (DASGUPTA 7.17)

Let $X \sim \text{Bin}(n, p)$. Show that $(X_n/n)^2$ and $X_n(X_n - 1)/(n(n - 1))$ both converging in probability to p^2 . Do they converge almost surely?

SOLUTION.

■

PROBLEM 11.9 (DASGUPTA 7.21)

Let X_1, X_2, \dots , be i.i.d. $U[0, 1]$. Let

$$G_n = (X_1 \cdots X_n)^{1/n}.$$

Find c such that $G_n \xrightarrow{\mathcal{P}} c$.

SOLUTION.

■

PROBLEM 11.10 (DASGUPTA 7.30 [CONCEPTUAL])

Suppose $X_n \xrightarrow{\mathcal{L}} X$, and also $Y_n \xrightarrow{\mathcal{L}} X$. Does this mean that $X_n - Y_n$ converge in distribution to (the point mass at) zero?

SOLUTION.

■

PROBLEM 11.11 (DASGUPTA 7.31 (A))

- (a) Suppose $a_n(X_n - \theta) \rightarrow N(0, \tau^2)$; what can be said about the limiting distribution of $|X_n|$, when $\theta \neq 0$, $\theta = 0$?

SOLUTION. ■