

# MA571 Problem Set 2

Carlos Salinas

September 7, 2015



**Problem 2.1 (Munkres §17, p. 100, 2)**

Show that if  $A$  is closed in  $Y$  and  $Y$  is closed in  $X$ , then  $A$  is closed in  $X$ .

*Proof.* Let  $C$  denote the closure of  $A$  in  $X$  then, by Theorem 17.4,  $A = \bar{A} = C \cap Y$  is the closure of  $A$  in  $Y$ . Thus,  $A$  is closed in  $X$  since it is the intersection of two closed subsets of  $X$ . ■

**Problem 2.2 (Munkres §17, p.100, 3)**

Show that if  $A$  is closed in  $X$  and  $B$  is closed in  $Y$ , then  $A \times B$  is closed in  $X \times Y$ .

*Proof.* Before proceeding we will prove the following set theoretic result (which was adapted from Exercises 2(n) and 2(o) from §1, p.14 of Munkres):

**Lemma 5.** *For sets  $A, B, C$  and  $D$  the following equalities hold:*

- (a)  $(A \times B) \cap (C \times D) = (A \cap C) \times (B \cap D)$ .
- (b)  $A \times (B \setminus C) = (A \times B) \setminus (A \times C)$ .
- (c)  $(A \setminus C) \times B = (A \times B) \setminus (C \times B)$ .

that is, the Cartesian product distributes over taking complements.

*Proof of Lemma 4.* (a) The equality follows (rather straightforwardly) from the definition of the Cartesian product and the complement of a set for  $x \times y \in (A \times B) \cap (C \times D)$  if and only if  $x \times y \in A \times B$  and  $x \times y \in C \times D$  if and only if  $x \in A$  and  $x \in C$  and  $y \in B$  and  $y \in D$  if and only if  $x \in A \cap C$  and  $y \in B \cap D$  if and only if  $x \times y \in (A \cap C) \times (B \cap D)$ .

(b) The point  $x \times y \in A \times (B \setminus C)$  if and only if  $x \in A$  and  $y \in B \setminus C$  if and only if  $x \in A$  and  $y \in B$  and  $y \notin C$  if and only if  $x \times y \in A \times B$  and  $x \times y \notin A \times C$  if and only if  $x \times y \in (A \times B) \setminus (A \times C)$ .

(c) The very same argument as part (b) can be used, taking  $B$  to be a subset of  $A$  and replacing (where appropriate)  $A$  by  $A \setminus B$  and  $B \setminus C$  by  $C$ , to prove that

$$(A \setminus B) \times C = (A \times C) \setminus (B \times C). \quad \blacklozenge$$

Now, since  $A$  is closed in  $X$  and  $B$  is closed in  $Y$ , their complements,  $X \setminus A$  and  $Y \setminus B$  are, by definition, open in  $X$  and  $Y$ , respectively. Then, the sets

$$(X \setminus A) \times Y \quad \text{and} \quad X \times (Y \setminus B)$$

are open since they are basic open sets in the product topology on  $X \times Y$ . So, applying Lemma 4(b) and (c), their complements

$$(X \times Y) \setminus (X \setminus A) \times Y = A \times Y \quad \text{and} \quad (X \times Y) \setminus X \times (Y \setminus B) = X \times B$$

are closed in  $X \times Y$ . At last, we have that

$$(A \times Y) \cap (X \times B)$$

is the intersection of closed sets, hence, by Theorem 17.1(b), is closed. By Lemma 4(a),

$$(A \times Y) \cap (X \times B) = (A \cap X) \times (Y \cap B) = A \times B$$

so  $A \times B$  is closed in  $X \times Y$ . ■

**Problem 2.3 (Munkres §17, p.101, 6(b))**

Let  $A$ ,  $B$  and  $A_\alpha$  denote subsets of a space  $X$ . Prove the following:

(b)  $\overline{A \cup B} = \bar{A} \cup \bar{B}$ .

*Proof.* By definition, the closure of a set is the intersection of all closed sets which contain it therefore,  $\overline{A \cup B} \subset \bar{A} \cup \bar{B}$  since  $\bar{A} \cup \bar{B}$  is a closed set, by Theorem 17.1(a), which contains  $A \cup B$ . To see the reverse containment note that  $\bar{A} \subset \overline{A \cup B}$  since  $\overline{A \cup B}$  is a closed set which contains  $A$ . Similarly  $\bar{B} \subset \overline{A \cup B}$  so  $\bar{A} \cup \bar{B} \subset \overline{A \cup B}$ . Therefore,  $\overline{A \cup B} = \bar{A} \cup \bar{B}$  holds.

Naturally this results extends, by induction, to the case of finite unions of sets. ■

**Problem 2.4 (Munkres §17, p. 101, 6(c))**

Let  $A$ ,  $B$  and  $A_\alpha$  denote subsets of a space  $X$ . Prove the following:

(b)  $\overline{\bigcup A_\alpha} \supset \bigcup \overline{A_\alpha}$ .

*Proof.* Let  $C$  denote the set  $\overline{\bigcup A_\alpha}$ . It is clear, by the definition of the closure of a set, that  $\bar{A}_\alpha \subset C$  for every  $\alpha$  since  $C$  is a closed set which contains  $A_\alpha$ , so  $\bigcup_\alpha \bar{A}_\alpha \subset C$ .

The reverse is not true in general; in fact, as Theorem 17.1(3) suggests, an arbitrary union of closed sets is not even necessarily closed. For a concrete example consider the family  $A_r = \{r\}$  for  $r \in \mathbf{Q}$ . The closure of a point  $r$  in  $\mathbf{R}$  is itself since its complement,  $\mathbf{R} \setminus \{r\}$ , is the union of the open intervals  $(-\infty, r)$  and  $(r, \infty)$ ; in particular,  $\{r\}$  is the “smallest” closed set containing  $\{r\}$ . Hence, we see that the union

$$\bigcup_{r \in \mathbf{Q}} \bar{A}_r = \mathbf{Q},$$

but, by Example 6,  $\bar{\mathbf{Q}} = \mathbf{R}$ . ■

**Problem 2.5 (Munkres §17, p. 101, 7)**

Criticize the following “proof” that  $\overline{\bigcup A_\alpha} \subset \bigcup \bar{A}_\alpha$ : if  $\{A_\alpha\}$  is a collection of sets in  $X$  and if  $x \in \overline{\bigcup A_\alpha}$ , then every neighborhood  $U$  of  $x$  intersects  $\bigcup A_\alpha$ . Thus  $U$  must intersect some  $A_\alpha$ , so  $x$  must belong to the closure of some  $A_\alpha$ . Therefore,  $x \in \bigcup \bar{A}_\alpha$ .

*Critique.* The main argument, that “ $x$  must belong to the closure of some  $A_\alpha$ ”, is what is wrong here. The point  $x$  may belong to the closure of multiple  $A_\alpha$ ’s, in fact uncountably many of them, so that one would have to prove that if  $x$  belongs to the closure of some family  $A_\beta$  of set, then  $x$  must belong to the union of their closures. This takes us right back to what we are trying to prove. ■

**Problem 2.6 (Munkres §17, p. 101, 9)**

Let  $A \subset X$  and  $B \subset Y$ . Show that in the space  $X \times Y$ ,

$$\overline{A \times B} = \bar{A} \times \bar{B}.$$

*Proof.*

■



**Problem 2.7 (Munkres §17, p. 101, 10)**

Show that every order topology is Hausdorff.

*Proof.*

■

**Problem 2.8 (Munkres §17, p. 101, 13)**

Show that  $X$  is Hausdorff if and only if the *diagonal*  $\Delta = \{x \times x \mid x \in X\}$  is closed in  $X \times X$ .

*Proof.*

■

**Problem 2.9 (Munkres §18, p. 111, 4)**

Given  $x_0 \in X$  and  $y_0 \in Y$ , show that the maps  $f: X \rightarrow X \times Y$  and  $g: Y \rightarrow X \times Y$  defined by

$$f(x) = x \times y_0 \quad \text{and} \quad g(y) = x_0 \times y$$

are imbeddings.

*Proof.*

■

**Problem 2.10 (Munkres §18, p. 111-112, 8(a,b))**

Let  $Y$  be an ordered set in the order topology. Let  $f, g: X \rightarrow Y$  be continuous.

- (a) Show that the set  $\{x \mid f(x) \leq g(x)\}$  is closed in  $X$ .
- (b) Let  $h: X \rightarrow Y$  be the function

$$h(x) = \min\{f(x), g(x)\}.$$

Show that  $h$  is continuous. [*Hint:* Use the pasting lemma.]

*Proof.*

■

**Problem 2.11**

Given:  $X$  is a topological space with open sets  $U_1, \dots, U_n$  such that  $\bar{U}_i = X$  for all  $i$ . Prove that the closure of  $U_1 \cap \dots \cap U_n$  is  $X$ .

*Proof.*

■