

MA 519: Homework, Midterms and Practice Problems Solutions

Carlos Salinas

Last compiled: October 24, 2016

Contents

1	Homework Solutions	2
1.1	Homework 1	2
1.2	Homework 2	6
1.3	Homework 3	9
1.4	Homework 4	12
1.5	Homework 5	14
1.6	Homework 6	16
1.7	Homework 7	18
1.8	Homework 8	20
2	Finals and Midterms	22
2.1	Midterm 1	22
3	Midterms, Exams, and Qualifying Exams	23
3.1	Qualifying Exams, August '99	23
3.2	Qualifying Exam, January '06	24
3.3	Qualifying Exam, August '14	26

1 Homework Solutions

These are my (corrected) solutions to DasGupta's Math/Stat 519 homework for the fall semester of 2016.

1.1 Homework 1

PROBLEM 1.1.1 (Handout 1, # 5). A closet contains five pairs of shoes. If four shoes are selected at random, what is the probability that there is at least one complete pair among the four?

SOLUTION. Let A denote the event “at least one complete pair is among the four shoes that were selected at random”. We compute the probability $P(\Omega \setminus A)$, i.e., the probability of the event “there is no complete pair among the four shoes that were selected at random”.

Under the equal likeliness assumption, the problem reduces to combinatorics. There are $\binom{10}{4} = 210$ ways to choose four shoes from five pairs. Therefore, $|\Omega| = 210$ and every point $x \in \Omega$ has a corresponding probability of $\frac{1}{210}$. Now, in order to avoid selecting a pair, we can choose one of two (a left or a right) shoe from four pairs of shoes which we can choose in $\binom{5}{4} = 5$ ways.

Thus, the probability that there is at least one pair of shoes among the four shoes selected at random is

$$\begin{aligned} P(A) &= 1 - P(\Omega \setminus A) \\ &= 1 - \frac{5 \cdot 2^4}{210} \\ &= 1 - \frac{8}{21} \\ &\approx 0.619. \end{aligned}$$

■

PROBLEM 1.1.2 (Handout 1, # 7). A gene consists of 10 subunits, each of which is normal or mutant. For a particular cell, there are 3 mutant and 7 normal subunits. Before the cell divides into 2 daughter cells, the gene duplicates. The corresponding gene of cell 1 consists of 10 subunits chosen from the 6 mutant and 14 normal units. Cell 2 gets the rest. What is the probability that one of the cells consists of all normal subunits.

SOLUTION. Let A denote the event “at least one of the two cells consists of all normal subunits”. Assuming we can arrange these subunits in any order, each daughter cell can receive 10 of the 20 subunits in $\binom{20}{10} = 184756$ ways. Hence, $|\Omega| = 184756$. Now, suppose one of the daughter cells receives a full 10 normal subunits. There are $\binom{14}{10} = 1001$ ways to do this. Since we can do this for the other cell as well, there are $2 \cdot 1001 = 2002$ ways of giving daughter one cell a full 10 normal subunits. Thus, the probability that at least one of the two daughter cells consists entirely of normal subunits is

$$P(A) = \frac{2002}{184756} \approx 0.011.$$

■

PROBLEM 1.1.3 (Handout 1, # 9). From a sample of size n , r elements are sampled at random. Find the probability that none of the N prespecified elements are included in the sample, if sampling is

- (a) with replacement;
- (b) without replacement.

Compute it for $r = N = 10$, $n = 100$.

SOLUTION. For part (a), the sample space contains $|\Omega| = n^r$ elements; these correspond to all of the possible draws of r with replacement. If n elements are specified, $n - N$ have not and therefore, there are $(n - N)^r$ ways of drawing r of the non-prespecified elements. Thus, the probability that none of the N prespecified elements are drawn if we sample r elements randomly with replacement is

$$p_1(n, N, r) = \frac{(n - N)^r}{n^r}. \quad (1)$$

The argument for part (b) is similar. There number of elements in the sample space is $|\Omega| = \binom{n}{r}$; these correspond to the possible draws of r elements without replacement. As before, $n - N$ of the elements have not been prespecified and therefore, we have $\binom{n - N}{r}$ ways of drawing r of the non-prespecified elements. Thus, the probability that none of the N prespecified elements are drawn if we sample r elements randomly without replacement is

$$p_2(n, N, r) = \frac{\binom{n - N}{r}}{\binom{n}{r}}. \quad (2)$$

Finally, we compute p_1 and p_2 for $r = N = 10$, $n = 100$ using equations (1) and (2) above:

$$\begin{aligned} p_1(100, 10, 10) &\approx 0.349, \\ p_2(100, 10, 10) &\approx 0.330. \end{aligned} \quad \blacksquare$$

PROBLEM 1.1.4 (Handout 1, # 11). Let E , F , and G be three events. Find expressions for the following events:

- (a) only E occurs;
- (b) both E and G occur, but not F ;
- (c) all three occur;
- (d) at least one of the events occurs;
- (e) at most two of them occur.

SOLUTION. For part (a), the statement “only E occurs” is interpreted as “we only consider those x in E , F , or G such that they belong only to E ”. In set-theoretic language, this is written

$$E \cap (\Omega \setminus F) \cap (\Omega \setminus G).$$

For part (b), \blacksquare

PROBLEM 1.1.5 (Handout 1, # 12). Which is more likely:

- (a) Obtaining at least one six in six rolls of a fair die, or
- (b) Obtaining at least one double six in six rolls of a pair of fair dice.

SOLUTION. For part (a), we compute the probability of the complement of the event, i.e., the event that we do not roll a 6 in 6 rolls of a fair die. This is

$$P(A) = 1 - P(\neg A) = 1 - \left(\frac{5}{6}\right)^6 \approx 0.665.$$

For part (b), we, again, compute the probability of the complement of the event, i.e., the event that no double 6 is rolled. This is

$$P(A) = 1 - P(\neg A) = 1 - \left(\frac{35}{36}\right)^6 \approx 0.156.$$

Therefore, it is more likely that you roll a 6 in 6 rolls of a fair die than that you roll a double 6 in 6 rolls of a pair of fair dice. ■

PROBLEM 1.1.6 (Handout 1, # 13). There are n people are lined up at random for a photograph. What is the probability that a specified set of r people happen to be next to each other?

SOLUTION. The r specified people can stand as a group starting at positions $1, 2, \dots, n - r + 1$. They can be permuted among themselves in $r!$ ways. The remaining $n - r$ people can be permuted among themselves in $(n - r)!$ ways. Thus, we have

$$P = \frac{(n - r + 1)r!(n - r)!}{n!} = \frac{(n - r + 1)!r!}{n!}.$$

■

PROBLEM 1.1.7 (Handout 1, # 16). Consider a particular player, say North, in a Bridge game. Let X be the number of aces in his hand. Find the distribution of X .

SOLUTION. The number of aces in North's hand X is a random variable which takes integer values between 0 and 4. Therefore, we must compute the probabilities $P(X = x)$, $0 \leq x \leq 4$.

From a deck with 52 cards, 13 cards can be selected in $\binom{52}{13}$ ways among which the player will have x number of aces in $\binom{4}{x}\binom{48}{13-x}$ ways. Thus, the distribution is given by

$$P(X = x) = \frac{\binom{4}{x}\binom{48}{13-x}}{\binom{52}{13}}.$$

The explicit values are approximately

$$\begin{aligned} P(X = 0) &\approx 0.304, & P(X = 1) &\approx 0.439, \\ P(X = 2) &\approx 0.213, & P(X = 3) &\approx 0.041, \\ P(X = 4) &\approx 0.003. \end{aligned}$$

■

PROBLEM 1.1.8 (Handout 1, # 20). If 100 balls are distributed completely at random into 100 cells, find the expected value of the number of empty cells.

Replace 100 by n and derive the general expression. Now approximate it as n tends to ∞ .

SOLUTION. Let X be the random variable denoting the number of empty cells. Define indicator variables X_1, \dots, X_n by

$$X_i = \begin{cases} 1 & \text{if the } i^{\text{th}} \text{ cell is empty,} \\ 0 & \text{otherwise.} \end{cases}$$

Then

$$X = \sum_{i=1}^n X_i$$

and so the mean of this random variable is given by

$$\begin{aligned} E(X) &= \sum_{i=1}^n E(X_i) \\ &= \sum_{i=1}^n P(X_i = 1) \\ &= \sum_{i=1}^n \frac{(n-1)^n}{n^n} \\ &= n \left(1 - \frac{1}{n} \right)^n, \end{aligned}$$

which approaches ∞ as $n \rightarrow \infty$ since $\left(1 - \frac{1}{n}\right)^n \rightarrow e^{-1}$.

For $n = 100$, we have $E(X) \approx 36.60$.

■

1.2 Homework 2

PROBLEM 1.2.1 (Handout 2, # 5). Four men throw their watches into the sea, and the sea brings each man one watch back at random. What is the probability that at least one man gets his own watch back?

SOLUTION. Suppose 4 men throw their watches into the sea. Label these men “the i^{th} man” for $1 \leq i \leq 4$. Let A_i denote the probability that the i^{th} man gets his own watch back. Then the probability that at least one man gets his own watch back is the event $A = A_1 \cup A_2 \cup A_3 \cup A_4$. By the inclusion-exclusion principle, we have

$$\begin{aligned} P(A) = & P(A_1) + P(A_2) + P(A_3) + P(A_4) \\ & - P(A_1 \cap A_2) - P(A_1 \cap A_3) - P(A_1 \cap A_4) \\ & - P(A_2 \cap A_3) - P(A_2 \cap A_4) - P(A_3 \cap A_4) \\ & + P(A_1 \cap A_2 \cap A_3) + P(A_1 \cap A_2 \cap A_4) \\ & + P(A_1 \cap A_3 \cap A_4) + P(A_2 \cap A_3 \cap A_4) \\ & - P(A_1 \cap A_2 \cap A_3 \cap A_4). \end{aligned} \tag{3}$$

Since the probabilities $P(A_i) = P(A_j)$, $P(A_i \cap A_j) = P(A_k \cap A_\ell)$, etc., the equation above reduces to

$$P(A) = 4P(A_1) - 6P(A_1 \cap A_2) + 4P(A_1 \cap A_2 \cap A_3) - P(A_1 \cap A_2 \cap A_3 \cap A_4).$$

All we need to do now is fill in the blanks. Intuitively, the probability that the 1st man gets back his own wallet is $\frac{1}{4}$ since only one wallet is his own out of the 4. Now, the probability that the 1st and the 2nd man get their own wallet back is $\frac{1}{4} \cdot \frac{1}{3} = \frac{1}{12}$. Proceeding in this fashion, we have

$$P(A) = 4 \cdot \frac{1}{4} - 6 \cdot \frac{1}{12} + 4 \cdot \frac{1}{24} - \frac{1}{24} = 0.625.$$

■

PROBLEM 1.2.2 (Handout 2, # 7). Calculate the probability that in Bridge, the hand of at least one player is void in a particular suit.

SOLUTION. Label the players “player i ” for $1 \leq i \leq 4$. Let A_i denote the event that player i is void in a particular suit. Then the probability of the event A that at least one player is void in a particular suit is $P(A) = P(\bigcup_{i=1}^4 A_i)$. Again, we can decompose this into

$$P(A) = 4P(A_1) - 6P(A_1 \cap A_2) + 4P(A_1 \cap A_2 \cap A_3) - P(A_1 \cap A_2 \cap A_3 \cap A_4).$$

Thus, it suffices to fill in the blanks in the equation above, i.e., find the probabilities $P(A_1)$, $P(A_1 \cap A_2)$, etc. To this end, the probability that player 1 is void in a particular suit, say red hearts, is

$$P(A_1) = \frac{\binom{52-13}{13}}{\binom{52}{13}}.$$

Similarly, the probability that player 1 and player 2 is void in red hearts is

$$P(A_1) = \frac{\binom{52-13}{13}}{\binom{52}{13}} \cdot \frac{\binom{52-13-13}{13}}{\binom{52-13}{13}} = \frac{\binom{52-13-13}{13}}{\binom{52}{13}};$$

and so on.

Thus,

$$P(A) = 4 \cdot \frac{\binom{52-13}{13}}{\binom{52}{13}} - 6 \cdot \frac{\binom{52-13-13}{13}}{\binom{52}{13}} + 4 \cdot \frac{\binom{52-13-13-13}{13}}{\binom{52}{13}} - 1 \cdot 0 \approx 0.051.$$

■

PROBLEM 1.2.3 (Handout 2, # 12). If n balls are placed at random into n cells, find the probability that exactly one cell remains empty.

SOLUTION. There are n ways to choose a cell to be left empty. Of the remaining $n - 1$ cells, one must contain 2 balls. That cell can be chosen in $n - 1$ ways and the two balls to be placed in it can be chosen in $\binom{2}{n}$ ways. Of the remaining cells, each must contain 1 ball and this pairing can be done in $(n - 2)!$ ways. Thus, the probability of the event A that exactly one cell remains empty is

$$P(A) = \frac{n(n-1)\binom{2}{n}(n-2)!}{n^n} = \frac{\binom{2}{n}n!}{n^n}.$$

■

PROBLEM 1.2.4 (Handout 2, # 13). (*Spread of rumors*). In a town of $n + 1$ inhabitants, a person tells a rumor to a second person, who in turn repeats it to a third person, etc. At each step the recipient of the rumor is chosen at random from the n people available. Find the probability that the rumor told r times without:

- (a) returning to the originator,
- (b) being repeated to any person.

Do the same problem when at each step the rumor told by one person to a gathering of N randomly chosen people. (The first question is the special case $N = 1$).

SOLUTION. For part (a), the originator can tell any one of the other n inhabitants the rumor. In turn, the non-originator can tell the other $n - 1$ inhabitants (not including the originator). Thus, the probability of the event A that the rumor told r does not return to the originator is

$$P(A) = \frac{n(n-1)^{r-1}}{n^r} = \left(\frac{n-1}{n}\right)^{r-1}.$$

For part (b),

■

PROBLEM 1.2.5 (Handout 2, # 14). What is the probability that

- (a) the birthdays of twelve people will fall in twelve different calendar months (assume equal probabilities for the twelve months),
- (b) the birthdays of six people will fall in exactly two calendar months?

SOLUTION.

■

PROBLEM 1.2.6 (Handout 2, # 15). A car is parked among N cars in a row, not at either end. On his return the owner finds exactly r of the N places still occupied. What is the probability that both neighboring places are empty?

SOLUTION.

■

PROBLEM 1.2.7 (Handout 2, # 16). Find the probability that in a random arrangement of 52 bridge card no two aces are adjacent.

SOLUTION.

■

PROBLEM 1.2.8 (Handout 2, # 17). Suppose $P(A) = \frac{3}{4}$, and $P(B) = \frac{1}{3}$. Prove that $P(A \cap B) \geq \frac{1}{12}$. Can it be equal to $\frac{1}{12}$?

SOLUTION.

■

PROBLEM 1.2.9 (Handout 2, # 18). Suppose you have infinitely many events A_1, A_2, \dots , and each one is sure to occur, i.e., $P(A_i) = 1$ for any i . Prove that $P(\bigcap_{i=1}^n A_i) = 1$.

SOLUTION.

■

PROBLEM 1.2.10 (Handout 2, # 19). There are n blue, n green, n red, and n white balls in an urn. Four balls are drawn from the urn with replacement. Find the probability that there are balls of at least three different colors among the four drawn.

SOLUTION.

■

1.3 Homework 3

PROBLEM 1.3.1 (Handout 3, # 3). n sticks are broken into one short and one long part. The $2n$ parts are then randomly paired up to form n new sticks. Find the probability that

- (a) the parts are joined in their original order, i.e., the new sticks are the same as the old sticks;
- (b) each long part is paired up with a short part.

SOLUTION. ■

PROBLEM 1.3.2 (Handout 3, # 5). In a town, there are 3 plumbers. On a certain day, 4 residents need a plumber and they each call one plumber at random.

- (a) What is the probability that all the calls go to one plumber (not necessarily a specific one)?
- (b) What is the expected value of the number of plumbers who get a call?

SOLUTION. ■

PROBLEM 1.3.3 (Handout 4, # 7). (*Polygraphs*). Polygraphs are routinely administered to job applicants for sensitive government positions. Suppose someone actually lying fails the polygraph 90% of the time. But someone telling the truth also fails the polygraph 15% of the time. If a polygraph indicates that an applicant is lying, what is the probability that he is in fact telling the truth? Assume a general prior probability p that the person is telling the truth.

SOLUTION. ■

PROBLEM 1.3.4 (Handout 4, # 8). In a bolt factory machines A , B , C manufacture, respectively, 25, 35, and 40 percent of the total. Of their output 5, 4, and 2 per cent are defective bolts. A bolt is drawn at random from the produce and is found defective. What are the probabilities that it was manufactured by machines A , B , C ?

SOLUTION. ■

PROBLEM 1.3.5 (Handout 4, # 9). Suppose that 5 men out of 100 and 25 women out of 10 000 are colorblind. A colorblind person is chosen at random. What is the probability of his being male? (Assume males and females to be in equal numbers.)

SOLUTION. ■

PROBLEM 1.3.6 (Handout 4, # 10). (*Bridge*). In a Bridge party West has no ace. What probability should he attribute to the event of his partner having

- (a) no ace,

- (b) two or more aces?

Verify the result by a direct argument.

SOLUTION. ■

PROBLEM 1.3.7 (Handout 4, # 12). A true-false question will be posed to a couple on a game show. The husband and the wife each has a probability p of picking the correct answer. Should they decide to let one of them answer the question, or decide that they will give the common answer if they agree and toss a coin to pick the answer if they disagree?

SOLUTION. ■

PROBLEM 1.3.8 (Handout 4, # 13). An urn containing 5 balls has been filled up by taking 5 balls at random from a second urn which originally had 5 black and 5 white balls. A ball is chosen at random from the first urn and is found to be black. What is the probability of drawing a white ball if a second ball is chosen from among the remaining 4 balls in the first urn?

SOLUTION. ■

PROBLEM 1.3.9 (Handout 4, # 15). Events A , B , C have probabilities p_1 , p_2 , p_3 . Given that exactly two of the three events occurred, the probability that C occurred is greater than $\frac{1}{2}$ if and only if ... (write down the necessary and sufficient condition).

SOLUTION. ■

PROBLEM 1.3.10 (Handout 5, # 1). There are five coins on a desk: 2 are double-headed, 2 are double-tailed, and 1 is a normal coin.

One of the coins is selected at random and tossed. It shows heads.

What is the probability that the other side of this coin is a tail?

SOLUTION. ■

PROBLEM 1.3.11 (Handout 5, # 2). (*Genetic testing*). There is a 50-50 chance that the Queen carries the gene for hemophilia. If she does, then each Prince has a 50-50 chance of carrying it. Three Princesses were recently tested and found to be non-carriers. Find the following probabilities:

- (a) that the Queen is a carrier;
- (b) that the fourth Princess is a carrier.

SOLUTION. ■

PROBLEM 1.3.12 (Handout 5, # 4). (*Is Johnny in Jail*). Johnny and you are roommates. You are a terrific student and spend Friday evenings drowned in books. Johnny always goes out on Friday evenings. 40% of the times, he goes out with his girlfriend, and 60% of the times he goes to a bar. If he goes out with his girlfriend, 30% of the times he is just too lazy to come back and spends the night at hers. If he goes to a bar, 40% of the times he gets mad at the person sitting on his right, beats him up, and goes to jail. On one Saturday morning, you wake up to see Johnny is missing. Where is Johnny?

SOLUTION.

■

1.4 Homework 4

PROBLEM 1.4.1 (Handout 5, # 2). In an urn, there are 12 balls. 4 of these are white. Three players: A , B , and C , take turns drawing a ball from the urn, in the alphabetical order. The first player to draw a white ball is the winner. Find the respective winning probabilities: assume that at each trial, the ball drawn in the trial before is put back into the urn (i.e., selection *with replacement*).

SOLUTION. ■

PROBLEM 1.4.2 (Handout 5, # 8). Consider n families with 4 children each. How large must n be to have a 90% probability that at least 3 of the n families are all girl families?

SOLUTION. ■

PROBLEM 1.4.3 (Handout 5, # 10). (*Yahtzee*). In Yahtzee, five fair dice are rolled. Find the probability of getting a Full House, which is three rolls of one number and two rolls of another, in Yahtzee.

SOLUTION. ■

PROBLEM 1.4.4 (Handout 5, # 12). The probability that a coin will show all heads or all tails when tossed four times is 0.25. What is the probability that it will show two heads and two tails?

SOLUTION. ■

PROBLEM 1.4.5 (Handout 5, # 13). Let the events A_1, A_2, \dots, A_n be independent and $P(A_k) = p_k$. Find the probability p that none of the events occurs.

SOLUTION. ■

PROBLEM 1.4.6 (Handout 6, # 5). Suppose a fair die is rolled twice and suppose X is the absolute value of the difference of the two rolls. Find the PMF and the CDF of X and plot the CDF. Find a median of X ; is the median unique?

SOLUTION. ■

PROBLEM 1.4.7 (Handout 6, # 7). Find a discrete random variable X such that $E(X) = E(X^3) = 0$; $E(X^2) = E(X^4) = 1$.

SOLUTION. ■

PROBLEM 1.4.8 (Handout 6, # 9). (*Runs*). Suppose a fair die is rolled n times. By using the indicator variable method, find the expected number of times that a six is followed by at least two other sixes. Now compute the value when $n = 100$.

SOLUTION. ■

PROBLEM 1.4.9 (Handout 6, # 10). (*Birthdays*). For a group of n people find the expected number of days of the year which are birthdays of exactly k people. (Assume 365 days and that all arrangements are equally probable.)

SOLUTION. ■

PROBLEM 1.4.10 (Handout 6, # 11). (*Continuation*). Find the expected number of multiple birthdays. How large should n be to make this expectation exceed 1?

SOLUTION. ■

PROBLEM 1.4.11 (Handout 6, # 12). (*The blood-testing problem*). A large number, N , of people are subject to a blood test. This can be administered in two ways, (i) Each person can be tested separately. In this case N tests are required, (ii) The blood samples of k people can be pooled and analyzed together. If the test is negative, this one test suffices for the k people. If the test is positive, each of the k persons must be tested separately, and in all $k + 1$ tests are required for the k people. Assume the probability p that the test is positive is the same for all people and that people are stochastically independent.

- (b) What is the expected value of the number, X , of tests necessary under plan (ii)?
- (c) Find an equation for the value of k which will minimize the expected number of tests under the second plan. (Do not try numerical solutions.)

SOLUTION. ■

PROBLEM 1.4.12 (Handout 6, # 13). (*Sample structure*). A population consists of r (classes whose sizes are in the proportion $p_1 : p_2 : \cdots : p_r$). A random sample of size n is taken with replacement. Find the expected number of classes not represented in the sample.

SOLUTION. ■

1.5 Homework 5

PROBLEM 1.5.1 (Handout 7, # 6(d, f)). Find the variance of the following random variables

- (d) $X = \#$ of tosses of a fair coin necessary to obtain a head for the first time.
- (f) $X = \#$ matches observed in random sitting of 4 husbands and their wives in opposite sides of a linear table.

This is an example of the *matching problem*.

SOLUTION. ■

PROBLEM 1.5.2 (Handout 7, # 8). (*Nonexistence of variance*).

- (a) Show that for a suitable positive constant c , the function $p(x) = \frac{c}{x^3}$, $x = 1, \dots$, is a valid probability mass function (PMF).
- (b) Show that in this case, the expectation of the underlying random variable exists, but the variance does not!

SOLUTION. ■

PROBLEM 1.5.3 (Handout 7, # 9). In a box, there are 2 black and 4 white balls. These are drawn out one by one at random (without replacement).

- (a) Let X be the draw at which the first black ball comes out. Find the mean the variance of X .
- (b) Let X be the draw at which the second black ball comes out. Find the meman the variance of X .

SOLUTION. ■

PROBLEM 1.5.4 (Handout 7, # 10). Suppose X has a *discrete uniform distribution* on the set $\{1, \dots, N\}$.

Find formulas for the mean and the variance of X .

SOLUTION. ■

PROBLEM 1.5.5 (Handout 7, # 11). (*Be Original*). Give an example of a random variable with mean 1 and variance 100.

SOLUTION. ■

PROBLEM 1.5.6 (Handout 7, # 13). (*Be Original*). Suppose a random variable X has the property that its second and fourth moment are both 1.

What can you say about the nature of X ?

SOLUTION. ■

PROBLEM 1.5.7 (Handout 7, # 14). (*Be Original*). One of the following inequalities is true in general for all nonnegative random variables. Identify which one!

$$E(X)E(X^4) \geq E(X^2)E(X^3);$$

$$E(X)E(X^4) \leq E(X^2)E(X^2).$$

SOLUTION. ■

PROBLEM 1.5.8 (Handout 7, # 15). Suppose X is the number of heads obtained in 4 tosses of a fair coin.

Find the expected value of the weird function

$$\ln\left(2 + \sin\left(\frac{\pi}{4}x\right)\right).$$

SOLUTION. ■

PROBLEM 1.5.9 (Handout 7, # 16). In a sequence of Bernoulli trials let X be the length of the run (of either successes or failures) started by the first trial.

- (a) Find the distribution of X , $E(X)$, $\text{Var}(X)$.

SOLUTION. ■

PROBLEM 1.5.10 (Handout 7, # 17). A man with n keys wants to open his door and tries the keys independently and at random. Find the mean and variance of the number of trials

- (a) if unsuccessful keys are not eliminated from further selections;
 (b) if they are.

(Assume that only one key fits the door. The exact distributions are given in II, 7, but are not required for the present problem.)

SOLUTION. ■

1.6 Homework 6

PROBLEM 1.6.1 (Handout 8, # 2). Identify the parameters n and p for each of the following binomial distributions:

- (a) # boys in a family with 5 children;
- (b) # correct answers in a multiple choice test if each question has a 5 alternatives, there are 25 questions, and the student is making guesses at random.

SOLUTION. ■

PROBLEM 1.6.2 (Handout 8, # 10). A newsboy purchases papers at 20 cents and sells them for 35 cents. He cannot return unsold papers. If the daily demand for papers is modeled as a $\text{Binom}(50, 0.5)$ random variable, what is the optimum number of papers the newsboy should purchase?

SOLUTION. ■

PROBLEM 1.6.3 (Handout 8, # 12). How many independent bridge dealings are required in order for the probability of a preassigned player having four aces at least once to be $\frac{1}{2}$ or better? Solve again for some player instead of a given one.

SOLUTION. ■

PROBLEM 1.6.4 (Handout 8, # 13). A book of 500 pages contains 500 misprints. Estimate the chances that a given page contains at least three misprints.

SOLUTION. ■

PROBLEM 1.6.5 (Handout 8, # 14). Colorblindness appears in 1 per cent of the people in a certain population. How large must a random sample (with replacements) be if the probability of its containing a colorblind person is to be 0.95 or more?

SOLUTION. ■

PROBLEM 1.6.6 (Handout 8, # 15). Two people toss a true coin n times each. Find the probability that they will score the same number of heads.

SOLUTION. ■

PROBLEM 1.6.7 (Handout 8, # 16). (*Binomial approximation to the hypergeometric distribution*). A population of N TV elements is divided into red and black elements in the proportion $p : q$ (where $p + q = 1$). A sample of size n is taken without replacement. The probability that it contains exactly k red elements is given by the hypergeometric distribution of II, 6. Show that as $n \rightarrow \infty$ this probability approaches $\text{Binom}(n, p)$.

SOLUTION. ■

PROBLEM 1.6.8 (Handout 9, # 3). Suppose X, Y, Z are mutually independent random variables, and $E(X) = 0$, $E(Y) = -1$, $E(Z) = 1$, $E(X^2) = 4$, $E(Y^2) = 3$, $E(Z^2) = 10$. Find the variance and the second moment of $2Z - \frac{Y}{2} + eX$, where e is the number such that $\ln e = 1$.

SOLUTION. ■

PROBLEM 1.6.9 (Handout 9, # 14). (*Variance of Product*). Suppose X, Y are independent random variables. Can it ever be true that $\text{Var}(XY) = \text{Var}(X) \text{Var}(Y)$? If it can, when?

SOLUTION. ■

1.7 Homework 7

PROBLEM 1.7.1 (Handout 10, # 4). (*Poisson Approximation*). One hundred people will each toss a fair coin 200 times. Approximate the probability that at least 10 of the 100 people would each have obtained exactly 100 heads and 100 tails.

SOLUTION. ■

PROBLEM 1.7.2 (Handout 10, # 5). (*A Pretty Question*). Suppose X is a Poisson distributed random variable. Can three different values of X have an equal probability?

SOLUTION. ■

PROBLEM 1.7.3 (Handout 10, # 6). (*Poisson Approximation*). There are 20 couples seated at a rectangular table, husbands on one side and the wives on the other, in a random order. Using a Poisson approximation, find the probability that exactly two husbands are seated directly across from their wives; at least three are; at most three are.

SOLUTION. ■

PROBLEM 1.7.4 (Handout 10, # 7). (*Poisson Approximation*). There are 5 coins on a desk, with probabilities 0.05, 0.1, 0.05, 0.01, and 0.04 for heads. By using a Poisson approximation, find the probability of obtaining at least one head when the five coins are each tossed once. Is the number of heads obtained binomially distributed in this problem?

SOLUTION. ■

PROBLEM 1.7.5 (Handout 10, # 8). A book of 500 pages contains 500 misprints. Estimate the chances that a given page contains at least three misprints.

SOLUTION. ■

PROBLEM 1.7.6 (Handout 10, # 9). Estimate the number of raisins which a cookie should contain on the average if it is desired that not more than one cookie out of a hundred should be without raisin.

SOLUTION. ■

PROBLEM 1.7.7 (Handout 10, # 10). The terms $p(k; \lambda)$ of the Poisson distribution reach their maximum when k is the largest integer not exceeding λ .

SOLUTION. ■

PROBLEM 1.7.8 (Handout 10, # 11). Prove

$$p(0; \lambda) + \cdots + p(n; \lambda) = \frac{1}{n!} \int_{\lambda}^{\infty} e^{-x} x^n dx.$$

SOLUTION. ■

PROBLEM 1.7.9 (Handout 10, # 12). There is a random number N of coins in your pocket, where N has a Poisson distribution with mean μ . Each one is tossed once.

Let X be the number of times a head shows.

Find the distribution of X .

SOLUTION. ■

PROBLEM 1.7.10 (Handout 10, # 14). Find the MGF of a general Poisson distribution, and hence prove that the mean and the variance of an arbitrary Poisson distribution are equal.

SOLUTION. ■

PROBLEM 1.7.11 (Handout 10, # 17 (a)). (*Poisson approximations*). 20 couples are seated in a rectangular table, husbands on one side and the wives on the other. First, find the expected number of husbands that sit directly across from their wives. Then, using a Poisson approximation, find the probability that two do; three do; at most five do.

SOLUTION. ■

1.8 Homework 8

PROBLEM 1.8.1 (Handout 12, # 2). Let X be Uniform $[a, b]$. Find the PDF, CDF, mean, and variance of X .

SOLUTION. ■

PROBLEM 1.8.2 (Handout 12, # 8). The diameter of a circular disk cut out by a machine has the following PDF

$$f(x) = \begin{cases} \frac{4x-x^2}{9} & \text{for } 1 \leq x \leq 4, \\ 0 & \text{otherwise.} \end{cases}$$

Find the average diameter of disks coming from this machine (in inches).

SOLUTION. ■

PROBLEM 1.8.3 (Handout 12, # 9). Suppose X is Uniform $[0, 2\pi]$. Find $P(-0.5 \leq \sin X \leq 0.5)$.

SOLUTION. ■

PROBLEM 1.8.4 (Handout 12, # 13). X has a piecewise uniform distribution on $[0, 1]$, $[1, 3]$, $[3, 6]$, and $[6, 10]$. Write its density function.

SOLUTION. ■

PROBLEM 1.8.5 (Handout 12, # 16). Show that for every p , $0 \leq p \leq 1$, the function $f(x) = p \sin x + (1-p) \cos x$, $0 \leq x \leq \frac{\pi}{2}$ (and $f(x) = 0$ otherwise), is a density function. Find its CDF and use it to find all the medians.

SOLUTION. ■

PROBLEM 1.8.6 (Handout 12, # 17). Give an example of a density function on $[0, 1]$ by giving a formula such that the density is finite at zero, unbounded at one, has a unique minimum in the open interval $(0, 1)$ and such that the median is 0.5.

SOLUTION. ■

PROBLEM 1.8.7 (Handout 12, # 18). (*A Mixed Distribution*). Suppose the damage claims on a particular type of insurance policy are uniformly distributed on $[0, 5]$ (in thousands of dollars), but the maximum by the insurance company is 2500 dollars. Find the CDF and the expected value of the payout, and plot the CDF. What is unusual about this CDF?

SOLUTION.

■

PROBLEM 1.8.8 (Handout 12, # 19). (*Random Distribution*). Jen's dog broke her six-inch long pencil off at a random point on the pencil. Find the density function and the expected value of the ratio of the lengths of the shorter piece and the longer piece of the pencil.

SOLUTION.

■

PROBLEM 1.8.9 (Handout 12, # 20). (*Square of a PDF Need Not Be a PDF*). Give an example of a density function $f(x)$ on $[0, 1]$ such that $cf^2(x)$ cannot be a density function for any c .

SOLUTION.

■

PROBLEM 1.8.10 (Handout 12, # 21). (*Percentiles of the Standard Cauchy*). Find the p^{th} percentile of the standard Cauchy density for a general p , and compute it for $p = 0.75$.

SOLUTION.

■

PROBLEM 1.8.11 (Handout 12, # 22). (*Integer Part*). Suppose X has a uniform distribution on $[0, 10.5]$. Find the expected value of the integer part of X .

SOLUTION.

■

PROBLEM 1.8.12 (Handout 12, # 23). X is uniformly distributed on some interval $[a, b]$. If its mean is 2, and variance is 3, what are the values of a , b ?

SOLUTION.

■

2 Finals and Midterms

2.1 Midterm 1

3 Midterms, Exams, and Qualifying Exams

3.1 Qualifying Exams, August '99

PROBLEM 3.1.1. The number of fish that Anirban catches on any given day has a Poisson distribution with mean 20. Due to the legendary softness of his heart, he sets free, on average, 3 out of the 4 fish he catches. Find the mean and the variance of the number of fish Anirban takes home on a given day.

SOLUTION.

■

PROBLEM 3.1.2. A fair die is rolled and at the same time a fair coin is tossed. This is done repeatedly. Find the probability that head occurs (strictly) before six occurs.

SOLUTION.

■

PROBLEM 3.1.3. X, Y are independent random variables with a common density $f(x) = e^{-|x|}/2$, $x \in (-\infty, \infty)$. Find the density function of $X + Y$.

SOLUTION.

■

PROBLEM 3.1.4. Let X_n denote the distance between two points chosen independently at random from the unit cube in \mathbb{R}^n . Evaluate

$$\lim_{n \rightarrow \infty} \frac{E(X_n)}{\sqrt{n}}.$$

SOLUTION.

■

PROBLEM 3.1.5. Let X be distributed as Uniform $[0, 1]$. What is the probability that the digit 5 does not occur in the decimal expansion of X ?

SOLUTION.

■

3.2 Qualifying Exam, January '06

PROBLEM 3.2.1. The birthdays of 5 people are known to fall in exactly 3 calendar months. What is the probability that exactly two of the 5 were born in January?

SOLUTION. ■

PROBLEM 3.2.2. Coupons are drawn, independently, with replacement, one at a time, from a set of 10 coupons. Find, explicitly, the expected number of draws

- (a) until the first draw coupon is drawn again;
- (b) until a duplicate occurs.

SOLUTION. ■

PROBLEM 3.2.3. Let N be a positive integer. Choose an integer at random from $\{1, \dots, N\}$. Let E be the event that your chosen random number is divisible by 3, and divisible by at least one of 4 and 6, but not divisible by 5. Find, explicitly, $\lim_{N \rightarrow \infty} P(E)$.

SOLUTION. ■

PROBLEM 3.2.4. Anirban is driving his Dodge on a highway with 4 lanes each way. He is wired to change lanes every minute on the minute. He changes with equal probability to either adjacent lane if there are two adjacent lanes, and the successive changes are mutually independent. Find, explicitly, the probability that after 4 minutes, Anirban is back to the lane he started from

- (a) if he started at an outside lane;
- (b) if he started at an inside lane.

SOLUTION. ■

PROBLEM 3.2.5. Burgess is going to Moose Pass, Alaska. He is driving his Dodge. He puts his car on cruise control at 70 mph. Gas stations are located every 30 miles, starting from his home. His car runs out of gas at a time distributed as an exponential with mean 4 hours. When that happens, he gets out, takes his bike out of his trunk, and bikes to the next gas station say M , at 10 mph. Let the time elapsed between when Burgess starts his trip and when he arrives at the gas station M be T . Find $E(T)$.

SOLUTION. ■

PROBLEM 3.2.6. A fair coin is tossed n times. Suppose X heads are obtained. Given $X = x$, let Y be generated according to the Poisson distribution with mean x . Find the unconditional variance of Y , and then find the limit of the probability $P(|Y - n/2| > n^{3/4})$, as $n \rightarrow \infty$.

SOLUTION.

■

PROBLEM 3.2.7. Anirban plays a game repeatedly. On each play he wins an amount uniformly distributed in $(0, 1)$ dollars, and then he tips the lady in charge of the game the square of the amount he has won. Then he plays again, tips again, and so on. Approximately calculate the probability that if he plays and tips six hundred times, his total winnings minus his total tips will exceed \$105.

SOLUTION.

■

PROBLEM 3.2.8. Anirban's dog got mad at him and broke his walking cane, first uniformly into two peices, and then the long piece again uniformly into two pieces. Find the probability that Anirban can make a triangle out of the three pieces of his cane.

SOLUTION.

■

PROBLEM 3.2.9. Suppose X, Y, Z are identically independently distributed $\text{Exp}(1)$ random variables. Find the joint density of (X, XY, XYZ) .

SOLUTION.

■

PROBLEM 3.2.10. Let X be the number of Kings and Y the number of Hearts in a Bridge hand. Find the correlation between X and Y .

SOLUTION.

■

3.3 Qualifying Exam, August '14

PROBLEM 3.3.1.

- (a) 3 balls are distributed one by one and at random in 3 boxes. What is the probability that exactly one box remains empty?
- (b) n balls are distributed one by one and at random in n boxes. Find the probability that exactly one box remains empty.
- (c) n balls are distributed one by one and at random in n boxes. Find the probability that exactly two boxes remain empty.

SOLUTION.

■

PROBLEM 3.3.2. n players each roll a fair die. For any pair of players i, j , $i < j$, who roll the same number, the group is awarded one point.

- (a) Find the mean of the total points of the group.
- (b) Find the variance of the total points of the group.

SOLUTION.

■

PROBLEM 3.3.3. Suppose X_1, X_2, \dots , is an infinite sequence of independently identically distributed Uniform $[0, 1]$ random variables. Find the limit

$$\lim_{n \rightarrow \infty} P \left[\frac{(\prod_{i=1}^n X_i)^{1/n}}{(\sum_{i=1}^n X_i)/n} > \frac{3}{4} \right].$$

SOLUTION.

■

PROBLEM 3.3.4. Suppose X is an exponential random variable with density $e^{-x/\sigma_1}/\sigma_1$ and Y is another exponential random variable with density $e^{-y/\sigma_2}/\sigma_2$, and that X, Y are independent.

- (a) Find the CDF of $X/(X + Y)$.
- (b) In the case $\sigma_1 = 2$, $\sigma_2 = 1$, find the mean of $X/(X + Y)$.

SOLUTION.

■

PROBLEM 3.3.5. Ten independently picked Uniform $[0, 100]$ numbers are each rounded to the nearest integer. Use the central limit theorem to approximate the probability that the sum of the ten rounded numbers equals the rounded value of the sum of the ten original numbers.

SOLUTION.

■

PROBLEM 3.3.6. Suppose for some given $m \geq 2$, we choose m independently identically distributed $\text{Uniform}[0, 1]$ random variables X_1, \dots, X_m . Let X_{\min} denote their minimum and X_{\max} denote their maximum. Now continue sampling X_{m+1}, \dots , from the $\text{Uniform}[0, 1]$ density. Let N be the first index k such that X_{m+k} falls outside the interval $[X_{\min}, X_{\max}]$.

- (a) Find a formula for $P(N > n)$ for a general n .
- (b) Hence, explicitly find $E(N)$.

SOLUTION.

■

PROBLEM 3.3.7. A $G_{n,p}$ graph on n vertices is obtained by adding each of the $\binom{n}{2}$ possible edges into the graph mutually independently with probability p . If vertex subsets A, B both have k vertices, and each vertex A shares an edge with each vertex in B , but there are no edges among the vertices within A or within B , then A, B generate a complete bipartate subgraph of order k denoted as $K_{k,k}$.

- (a) For a given n and p , find an expression for the expected number of complete bipartate subgraphs $K_{3,3}$ of order $k = 3$ in a $G_{5n,p}$ graph.
- (b) Let p_n denote the value of p for which the expected value in part (a) equals one. Identify constants α, β such that $\lim_{n \rightarrow \infty} n^\alpha p_n = \beta$.

SOLUTION.

■