

MA 523: Homework 9

Carlos Salinas

November 30, 2016

PROBLEM 9.1

- (a) Show that for $n = 3$ the general solution to the wave equation $u_{tt} - \Delta u = 0$ with spherical symmetry about the origin has the form

$$u = \frac{1}{r}F(r+t) + \frac{1}{r}G(r-t), \quad r = |x|,$$

with suitable F and G .

- (b) Show that the solution with initial data of the form

$$u(r, 0) = 0, \quad u_t(r, 0) = h(r)$$

(h is an even function of r) is given by

$$u = \frac{1}{2r} \int_{r-t}^{r+t} \rho h(\rho) d\rho.$$

SOLUTION. For part (a): Consider the initial-value problem

$$\begin{cases} u_{tt} - \Delta u = 0 & \text{in } \mathbb{R}^3 \times (0, \infty), \\ u = g, \quad u_t = h & \text{on } \mathbb{R}^3 \times \{t = 0\}. \end{cases} \quad (9.1)$$

Suppose that u is a solution to the initial-value problem (9.1) that is symmetric about the origin; i.e., $u(x) = u(x')$ whenever $|x| = |x'|$. Now, make the following substitution ■

PROBLEM 9.2

Show that the solution $w(x_1, t)$ of the initial-value problem for the *Klein-Gordon equation*

$$\begin{cases} w_{tt} = w_{x_1 x_1} - \lambda^2 w, \\ w(x_1, 0) = 0, \end{cases} \quad w_t(x_1, 0) = h(x_1) \quad (9.2)$$

is given by

$$w(x_1, t) = \frac{1}{2} \int_{x_1-t}^{x_1+t} J_0(\lambda s) h(y_1) dy_1.$$

Here $s^2 = t^2 - (x_1 - y_1)^2$, while J_0 denotes the Bessel function defined by

$$J_0(z) := \frac{2}{\pi} \int_0^{\frac{\pi}{2}} \cos(z \sin \theta) d\theta.$$

(*Hint:* Descend to (9.2) from the two-dimensional wave equation satisfied by

$$u(x_1, x_2, t) = \cos(\lambda x_2) w(x_1, t).)$$

SOLUTION. ■

PROBLEM 9.3

Let u solve

$$\begin{cases} u_{tt} - \Delta u = 0 & \text{in } \mathbb{R}^3 \times (0, \infty), \\ u = g, \quad u_t = h & \text{on } \mathbb{R}^3 \times \{t = 0\} \end{cases}$$

where g and h are smooth and have compact support. Show there exists a constant C such that

$$|u(x, t)| \leq Ct^{-1} \quad (x \in \mathbb{R}^3, t > 0).$$

SOLUTION.

■