

# Fall 2016 Notes

Carlos Salinas

September 19, 2016

## Contents

|   |           |
|---|-----------|
| <b>Contents</b>   | <b>1</b>  |
| <b>1 Probability</b>                                    | <b>2</b>  |
| 1.1 Discrete Probability . . . . .                      | 2         |
| <b>2 Introduction to Partial Differential Equations</b> | <b>4</b>  |
| 2.1 Introduction . . . . .                              | 4         |
| 2.2 The transport equation . . . . .                    | 6         |
| 2.3 Characteristics . . . . .                           | 7         |
| 2.4 Boundary conditions . . . . .                       | 11        |
| <b>3 Algebraic Geometry</b>                             | <b>15</b> |
| 3.1 The statement of de Rham's theorem . . . . .        | 15        |
| <b>4 Algebraic Topology</b>                             | <b>16</b> |
| 4.1 Cohomology . . . . .                                | 16        |
| <b>5 Group Theory and Differential Equations</b>        | <b>17</b> |
| <b>Bibliography</b>                                     | <b>18</b> |

# Chapter 1

## Probability

We will devote this chapter to the material that is covered in MA 51900 (discrete probability) as it was covered in DasGupta's class. We will, for the most part, reference Feller's *An introduction to probability theory and its applications, Volume 1* [4] (especially for the discrete noncalculus portion of the class) and DasGupta's own book *Fundamentals of Probability: A First Course* [2].

### 1.1 Discrete Probability

The material in this chapter is mostly pulled from Sheldon Ross's *A First Course in Probability Theory* [8] with some examples from [2] and [4]. I find Ross's book to be better structured than the latter two.

#### Combinatorial Analysis

These are the main results from this section.

**Theorem 1.1** (The basic principle of counting). *Suppose that two experiments are to be performed. Then if experiment 1 can result in any one of  $m$  possible outcomes and if, for each outcome of experiment 1, there are  $n$  possible outcomes of experiment 2, then together there are  $mn$  possible outcomes of the two experiments.*

**Theorem 1.2** (The generalized principle of counting). *If  $r$  experiments that are to be performed are such that the first one may result in any of  $n_1$  possible outcomes; and if, for each of these  $n_1$  possible outcomes, there are  $n_2$  possible outcomes for the second experiment; and if, for each of the possible outcomes of the first two experiments, there are  $n_3$  possible outcomes for the third experiment; etc. ..., then there is a total of  $n_1 n_2 \cdots n_r$  possible outcomes of the  $r$  experiments.*

Using notation as in [4], the number

$$(n)_r = n(n-1) \cdots (n-r+1)$$

represents the number of different ways that a group of  $r$  items could be selected from  $n$  items when the order of selection is relevant, and as each group of  $r$  items will be counted  $r!$  times in this count,

it follows that the number of different groups of  $r$  items that could be formed from a set of  $n$  items is

$$\frac{(n)_r}{r!} = \frac{n!}{(n-r)!r!}$$

for which we reserve the notation

$$\binom{n}{r}$$

read  $n$  choose  $r$ . (This is called a binomial coefficient since it appears in the binomial expansion  $(a+b)^n$ .)

A useful combinatorial identity on binomial coefficients is the following

$$\binom{n}{r} = \binom{n-1}{r-1} + \binom{n-1}{r}$$

for  $1 \leq r \leq n$ .

**Theorem 1.3** (The binomial theorem).

$$(a+b)^n = \sum_{i=0}^n \binom{n}{i} x^i y^{n-i}.$$

*PROOF.* We provide a combinatorial proof of the theorem. Consider the product

$$(a_1 + b_1) \cdots (a_n + b_n).$$

Its expansion consists of the sum of  $2^n$  terms, each term being the product of  $n$  factors. Furthermore, each of the  $2^n$  terms in the sum will contain as a factor either  $a_i$  or  $b_i$  for each  $1 \leq i \leq n$ . Now, how many of the  $2^n$  terms in the sum will have  $k$  of the  $a_i$  and  $n-k$  of the  $b_i$  as factors? As each term consisting of  $k$  of the  $a_i$  and  $n-k$  of the  $b_i$  correspond to a choice of a group of  $k$  from the values  $a_1, \dots, a_n$ , there are  $\binom{n}{k}$  such terms. Thus, letting  $a_i = a$ ,  $b_i = b$ ,  $1 \leq i \leq n$ , we see that

$$(a+b)^n = \sum_{i=0}^n \binom{n}{i} x^i y^{n-i}.$$

■

## Chapter 2

# Introduction to Partial Differential Equations

Here we summarize some important points about PDEs. The material is mostly taken from Evans's *Partial Differential Equations* [3] with occasional detours to Strauss's *Partial Differential Equations: An Introduction* [9]. We will be following Dr. Petrosyan's **Course Log** which can be found here <https://www.math.purdue.edu/~arshak/F16/MA523/courselog/>, i.e., summarizing the appropriate chapters from [3].

### 2.1 Introduction

#### Partial differential equations

**Definition 2.1.** An expression of the form

$$F(D^k u(x), D^{k-1} u(x), \dots, Du(x), u(x), x) = 0, \quad x \in U, \quad (2.1)$$

is called a *kth-order partial differential equation (PDE)*, where

$$F: \mathbf{R}^{n^k} \times \mathbf{R}^{n^{k-1}} \times \dots \times \mathbf{R}^n \times U \longrightarrow \mathbf{R}$$

is given, and

$$u: U \longrightarrow \mathbf{R}$$

is the unknown.

Here are some more definitions,

**Definition 2.2.**

- (i) The partial differential equation (2.1) is called *linear* if it has the form

$$\sum_{|\alpha| \leq k} a_\alpha(x) D^\alpha u = f(x)$$

for given functions  $a_\alpha (|\alpha| \leq k)$ ,  $f$ . This linear PDE is *homogeneous* if  $f = 0$ .

(ii) The PDE (2.1) is *semilinear* if it has the form

$$\sum_{|\alpha|=k} a_\alpha D^\alpha u + a_0(D^{k-1}u, \dots, Du, u, x) = 0.$$

(iii) The PDE (2.1) is *quasilinear* if it has the form

$$\sum_{|\alpha|=k} a_\alpha(D^{k-1}u, \dots, Du, u, x) D^\alpha u + a_0(D^{k-1}u, \dots, Du, u, x) = 0.$$

(iv) The PDE (2.1) is *fully nonlinear* if it depends upon the highest order derivatives.

A *system* of partial differential equations is, informally speaking, a collection of several PDEs for several unknown functions.

**Definition 2.3.** An expression of the form

$$\mathbf{F}(D^k \mathbf{u}(x), D^{k-1} \mathbf{u}(x), \dots, D\mathbf{u}(x), \mathbf{u}(x), x) = 0, \quad x \in U, \quad (2.2)$$

is called a *kth-order system of PDEs*, where

$$\mathbf{F}: \mathbf{R}^{mn^k} \times \mathbf{R}^{mn^{k-1}} \times \dots \times \mathbf{R}^{mn} \times \mathbf{R}^m \times U \longrightarrow \mathbf{R}^m$$

is given and

$$\mathbf{u}: U \longrightarrow \mathbf{R}^m, \quad \mathbf{u} = (u^1, \dots, u^m)$$

is the unknown.

*Remark 2.4.* We haven't talked much about systems of PDEs and I suspect we will not do so very much in this course.

## Examples

This is only a fraction of the PDEs listed in Evan's chapter.

### Linear equations

1. Laplace's equation

$$\Delta u = \sum_{i=1}^n u_{x_i x_i} = 0.$$

2. Helmholtz's (or eigenvalue) equation

$$-\Delta u = \lambda u.$$

3. Linear transport equation

$$u_t + \sum_{i=1}^n b^i u_{x_i} = 0.$$

4. Liouville's equation

$$u_t - \sum_{i=1}^n (b^i u)_{x_i} = 0.$$

5. Heat (or diffusion) equation

$$u_t - \Delta u = 0.$$

6. Wave equation

$$u_{tt} - \Delta u = 0.$$

7. Telegraph equation

$$u_{tt} + du_t - u_{xx} = 0.$$

### Nonlinear equations

1. Eikonal equation

$$|Du| = 1.$$

2. Nonlinear Poisson equation

$$-\Delta u = f(u).$$

3. Inviscid Burgers' equation

$$u_t + uu_x = 0.$$

and so on.

## 2.2 The transport equation

We begin our study with one of the simplest PDEs, the *transport equation* with constant coefficients. This is the PDE

$$u_t + b \cdot Du = 0, \quad \text{in } \mathbf{R}^n \times (0, \infty), \quad (2.3)$$

where  $b$  is a fixed vector in  $\mathbf{R}^n$ ,  $b = (b_1, \dots, b_n)$ ,  $x = (x_1, \dots, x_n) \in \mathbf{R}^n$  is a typical point in space,  $t \geq 0$  denotes a typical time and  $u: \mathbf{R} \times [0, \infty) \rightarrow \mathbf{R}$  is the unknown,  $u = u(x, t)$ . We write  $Du = D_x u = (u_{x_1}, \dots, u_{x_n})$  for the gradient of  $u$  with respect to the spatial variable  $x$ .

So, which functions solve (2.3)? Well, let us suppose for a moment that  $u$  is a smooth solution to the PDE and let us try to compute it. To do so, we first recognize that (2.3) asserts that a particular directional derivative of  $u$  vanishes, namely,  $D_b u = 0$ . We exploit this by fixing a point  $(x, t) \in \mathbf{R}^n \times (0, \infty)$  and defining

$$z(s) := u(x + sb, t + s), \quad s \in \mathbf{R}.$$

Then we calculate

$$\begin{aligned} \dot{z}(s) &= Du(x + sb, t + s) \cdot b + u_t(x + sb, t + s) \\ &= 0, \end{aligned}$$

the second equality holding by (2.3). Thus,  $z$  is a constant function of  $s$ , and consequently for each  $(x, t)$ ,  $u$  is constant on the line through  $(x, t)$  with direction  $(b, 1) \in \mathbf{R}^{n+1}$ . Hence, if we know the value of  $u$  at any point on each such line, we know its value everywhere in  $\mathbf{R}^n \times (0, \infty)$ .

### 2.3 Characteristics

#### Derivation of characteristic ODEs

Consider the nonlinear first-order PDE

$$F(Du, u, x) = 0 \quad \text{in } U, \quad (2.4)$$

subject now to the boundary condition

$$u = g \quad \text{on } \Gamma, \quad (2.5)$$

where  $\Gamma \subseteq \partial U$  and  $g: \Gamma \rightarrow \mathbf{R}$  are given. We hereafter suppose that  $F, g$  are smooth functions.

We now develop the method of *characteristics*, which solves (2.4) and (2.5) by converting the PDE into an appropriate system of ODEs. Suppose  $u$  solves the (2.4), (2.5) and fix any point  $x \in U$ . We would like to calculate  $u(x)$  by finding some curve lying within  $U$ , connecting  $x$  with a point  $x^0 \in \Gamma$  and along which we can compute  $u$ . Since (2.5) says  $u = g$  on  $\Gamma$ , we know the value of  $u$  at the one end  $x^0$ . We hope then to be able to calculate  $u$  all along the curve, and so in particular at  $x$ .

#### Finding the characteristic ODEs

How can we choose the curve so all this will work? Let us suppose it is described parametrically by the function  $\mathbf{x}(s) = (x^1(s), \dots, x^n(s))$ , the parameter  $s$  lying in some subinterval of  $\mathbf{R}$ . Assuming  $u$  is a  $C^2$  solution of (2.4), we define also

$$z(s) := u(\mathbf{x}(s)).$$

In addition, set

$$\mathbf{p}(s) := Du(\mathbf{x}(s));$$

that is,  $\mathbf{p}(s) = (p^1(s), \dots, p^n(s))$ , where

$$p^i(s) = u_{x_i}(\mathbf{x}(s)), \quad (2.6)$$

$1 \leq i \leq n$ . So  $z$  gives the values of  $u$  along the curve and  $\mathbf{p}$  records the values of the gradient  $Du$ . We must choose a function  $\mathbf{x}$  in such a way that we can compute  $z$  and  $\mathbf{p}$ .

For this, first differentiate (2.6)

$$\dot{p}^i(s) = \sum_{j=1}^n u_{x_i x_j}(\mathbf{x}(s)) \dot{x}^j(s)$$

This expression is not too promising, since it involves the second derivatives of  $u$ . On the other hand, we can also differentiate the PDE (2.4) with respect to  $x_i$  to get

$$\sum_{j=1}^n \frac{\partial}{\partial p_j} F(Du, u, x) u_{x_j x_i} + \frac{\partial}{\partial z} F(Du, u, x) u_{x_i} + \frac{\partial}{\partial x_i} F(Du, u, x) = 0.$$

We are able to employ this identity to get rid of the *dangerous* second derivative terms provided we first set

$$\dot{x}^j(s) = \frac{\partial}{\partial p_j} F(\mathbf{p}(s), z(s), \mathbf{x}(s)).$$

Assuming now that the above equation holds, we can evaluate the partials

$$\begin{aligned} \sum_{j=1}^n \frac{\partial}{\partial p_j} F(\mathbf{p}(s), z(s), \mathbf{x}(s)) \\ + \frac{\partial}{\partial z} F(\mathbf{p}(s), z(s), \mathbf{x}(s)) p^i(s) + \frac{\partial}{\partial x_i} F(\mathbf{p}(s), z(s), \mathbf{x}(s)) = 0. \end{aligned}$$

Substitute this expression and the previous one into the derivative for  $p^i$  and we get

$$\dot{p}^i(s) = \frac{\partial}{\partial x_i} F(\mathbf{p}(s), z(s), \mathbf{x}(s))$$

Finally, we differentiate  $z$  to get

$$\dot{z}(s) = \sum_{j=1}^n \frac{\partial}{\partial x_j} u(\mathbf{x}(s)) \dot{x}^j(s) = \sum_{j=1}^n p^j(s) \frac{\partial}{\partial p_j} F(\mathbf{p}(s), z(s), \mathbf{x}(s)),$$

the second equality holding by –fuck this guy for numbering every expression–(5) and (8)–whatever they are.

We summarize by rewriting equations (8)–(10) in vector notation:

$$\begin{cases} \text{(a) } \dot{\mathbf{p}}(s) = -D_x F(\mathbf{p}(s), z(s), \mathbf{x}(s)) - D_z F(\mathbf{p}(s), z(s), \mathbf{x}(s)) \mathbf{p}(s), \\ \text{(b) } \dot{z}(s) = D_p F(\mathbf{p}(s), z(s), \mathbf{x}(s)) \cdot \mathbf{p}(s), \\ \text{(c) } \dot{\mathbf{x}}(s) = D_p F(\mathbf{p}(s), z(s), \mathbf{x}(s)). \end{cases} \quad (2.7)$$

This important system of  $2n + 1$  first-order ODEs comprises the *characteristic equations* of the nonlinear first-order PDE (2.4). The functions  $\mathbf{p} = (p^1, \dots, p^n)$ ,  $z$ ,  $\mathbf{x} = (x^1, \dots, x^n)$  are called the *characteristics*. We will sometimes refer to  $\mathbf{x}$  as the *projected characteristics*: it is the projection of the full characteristics  $(\mathbf{p}, z, \mathbf{x}) \subseteq \mathbf{R}^{2n+1}$  onto the physical region  $U \subseteq \mathbf{R}^n$ .

**Theorem 2.5** (Structure of characteristic ODEs). *Let  $u \in C^2(U)$  solve the nonlinear, first-order partial differential equation (2.4) in  $U$ . Assume  $\mathbf{x}$  solves the ODEs (2.7)(c), where  $\mathbf{p} = Du$ ,  $z = u$ . Then  $\mathbf{p}$  solves the ODE (2.7)(a) and  $z$  solves the ODE (2.7)(b), for those  $s$  such that  $\mathbf{x} \in U$ .*

## Examples

### $F$ linear

Consider first the situation that (2.4) is linear and homogeneous, and thus has the form

$$F(Du, u, x) = \mathbf{b}(x) \cdot Du(x) + c(x)u(x) = 0, \quad x \in U.$$

Then  $F(p, z, x) = \mathbf{b}(x) \cdot p + c(x)z$ , and so

$$D_p F = \mathbf{b}(x).$$

In this circumstance (2.7)(c) becomes

$$\dot{\mathbf{x}}(s) = \mathbf{b}(\mathbf{x}(s)),$$



an ODE involving only the function  $\mathbf{x}$ . Furthermore (2.7)(b) becomes

$$\dot{z}(s) = \mathbf{b}(\mathbf{x}(s)) \cdot \mathbf{p}(s). \quad (2.8)$$

Since  $\mathbf{p}(\cdot) = Du(\mathbf{x}(\cdot))$ , the PDE simplifies the above to

$$\dot{z}(s) = -c(\mathbf{x}(s))z(s).$$

This ODE is linear in  $z$ , once we know the function  $\mathbf{x}$  by solving its ODE.

In summary,

$$\begin{cases} \text{(a)} & \dot{\mathbf{x}}(s) = \mathbf{b}(\mathbf{x}(s)) \\ \text{(b)} & \dot{z}(s) = -c(\mathbf{x}(s))z(s) \end{cases} \quad (2.9)$$

comprise the characteristic equations for the linear, first-order PDE (2.8).

**Example 2.6.** We demonstrate the utility of equations (2.9) by explicitly solving the problem

$$\begin{cases} x_1 u_{x_2} - x_2 u_{x_1} = u & \text{in } U \\ u = g & \text{on } \Gamma, \end{cases} \quad (2.10)$$

where  $U$  is the quadrant  $\{x_1 > 0, x_2 > 0\}$  and  $\Gamma = \{x_1 > 0, x_2 = 0\} \subseteq \partial U$ . The PDE in (2.10) is of the form (2.8), for  $\mathbf{b} = (-x_2, x_1)$  and  $c = -1$ . Thus the equations (2.9) read

$$\begin{cases} (x^1, x^2)(s) = (x^0 \cos s, x^0 \sin s) \\ z(s) = z^0 e^s = g(x^0) e^s, \end{cases} \quad (2.11)$$

where  $x^0 \geq 0$ ,  $0 \leq s \leq \pi/2$ . Fix a point  $(x_1, x_2) \in U$ . We select  $s > 0$ ,  $x^0 > 0$  so that  $(x_1, x_2) = (x^1(s), x^2(s)) = (x^0 \cos s, x^0 \sin s)$ . That is,  $x^0 = (x_1^2 + x_2^2)^{1/2}$ ,  $s = \arctan(x_2/x_1)$ . Therefore,

$$\begin{aligned} u(x_1, x_2) &= u(x^1(s), x^2(s)) \\ &= z(s) \\ &= g(x^0) e^s \\ &= g((x_1^2 + x_2^2)^{1/2}) e^{\arctan(x_2/x_1)}. \end{aligned}$$

### **$F$ quasilinear**

The PDE (2.4) is quasilinear if it has the form

$$F(Du, u, x) = \mathbf{b}(x, u(x)) \cdot Du(x) + c(x, u(x)) = 0. \quad (2.12)$$

In this circumstance  $F(p, z, x) = \mathbf{b}(x, z) \cdot p + c(x, z)$ ; whence

$$D_p F = \mathbf{b}(x, z).$$

Hence equation (2.9)(c) reads

$$\dot{\mathbf{x}} = \mathbf{b}(\mathbf{x}(s), z(s)),$$

and (2.9)(b) becomes

$$\begin{aligned}\dot{z}(s) &= \mathbf{b}(\mathbf{x}(s), z(s)) \cdot \mathbf{p}(s) \\ &= -c(\mathbf{x}(s), z(s))\end{aligned}$$

by (2.12). Consequently

$$\begin{cases} \text{(a)} \quad \dot{\mathbf{x}}(s) = \mathbf{b}(\mathbf{x}(s), z(s)), \\ \text{(b)} \quad \dot{z}(s) = -c(\mathbf{x}(s), z(s)) \end{cases} \quad (2.13)$$

are the characteristic equations for the quasilinear first-order PDE.

**Example 2.7.** The characteristic ODEs (2.13) are in general difficult to solve, and so we work out in this example the simpler case of a boundary-value problem for a semilinear PDE:

$$\begin{cases} u_{x_1} + u_{x_2} = u^2 & \text{in } U \\ u = g & \text{on } \Gamma. \end{cases} \quad (2.14)$$

Now  $U$  is the half-space  $\{x_2 > 0\}$  and  $\Gamma = \{x_2 = 0\} = \partial U$ . Here  $\mathbf{b} = (1, 1)$  and  $c = -z^2$ . Then (2.13) becomes

$$\begin{cases} (\dot{x}^1, \dot{x}^2) = 1 \\ \dot{z} = z^2. \end{cases}$$

Consequently

$$\begin{cases} (x^1, x^2)(s) = (x^0 + s, s) \\ z(s) = \frac{z^0}{1 - sz^0} \\ \quad = \frac{g(x^0)}{1 - sg(x^0)}, \end{cases}$$

where  $x^0 \in \mathbf{R}$ ,  $s \geq 0$ , provided the denominator is not zero.

Fix a point  $x_1, x_2 \in U$ . We select  $s > 0$  and  $x^0 \in \mathbf{R}$  so that  $(x_1, x_2) = (x^1(s), x^2(s)) = (x^0 + s, s)$ ; that is,  $x^0 = x_1 - x_2$ ,  $s = x_2$ . Then

$$\begin{aligned} u(x_1, x_2) &= u(x^1(s), x^2(s)) \\ &= z(s) \\ &= \frac{g(x^0)}{1 - sg(x^0)} \\ &= \frac{g(x_1 - x_2)}{1 - x_2 g(x_1 - x_2)}. \end{aligned}$$

This solution of course make sense only if  $1 - x_2 g(x_1 - x_2) \neq 0$ .

### **$F$ fully nonlinear**

In the general case, the full characteristic equations (2.7) must be integrated, if possible.

**Example 2.8.** Consider the fully nonlinear problem

$$\begin{cases} u_{x_1} u_{x_2} = u & \text{on } U \\ u = x_2^2 & \text{on } \Gamma \end{cases} \quad (2.15)$$

where  $U = \{x_1 > 0\}$ ,  $\Gamma = \{x_1 = 0\} = \partial U$ . Here  $F(p, z, x) = p_1 p_2 - z$ , and hence the characteristic ODEs (2.7) become

$$\begin{cases} (\dot{p}^1, \dot{p}^2) = (p^1, p^2) \\ \dot{z} = 2p^1 p^2 \\ (\dot{x}^1, \dot{x}^2) = (p^2, p^1). \end{cases}$$

We integrate these equations to find

$$\begin{cases} (x^1, x^2)(s) = (p_2^0(e^s - 1), p_1^0(e^s - 1)) \\ z(s) = z^0 + p_1^0 p_2^0 (e^{2s} - 1) \\ (p^1, p^2)(s) = (p_1^0 e^s, p_2^0 e^s), \end{cases}$$

where  $x^0 \in \mathbf{R}$ ,  $s \in \mathbf{R}$ , and  $z^0 = (x^0)^2$ .

We must determine  $p^0 = (p_1^0, p_2^0)$ . Since  $u = x_2^2$  on  $\Gamma$ ,  $p_2^0 = u_{x_2}(0, x^0) = 2x^0$ . Furthermore the PDE  $u_{x_1} u_{x_2} = u$  itself implies  $p_1^0 p_2^0 = z^0 = (x^0)^2$ , and so  $p_1^0 = x^2/2$ . Consequently the formulas above become

$$\begin{cases} (x^1, x^2)(s) = (2x^0(e^s - 1), x^0(e^s + 1)/2) \\ z(s) = (x^0)^2 e^{2s} \\ (p^1, p^2)(s) = (x^0 e^s/2, 2x^0 e^s). \end{cases}$$

Fix a point  $(x_1, x_2) \in U$ . Select  $s$  and  $x^0$  so that

$$(x_1, x_2) = (x^1(s), x^2(s)) = (2x^0(e^s - 1), x^0(e^s + 1)/2).$$

This equality implies  $x^0 = (4x_2 - x_1)/4$ ,  $e^s = (x_1 + 4x_2)/(4x_2 - x_1)$ ; and so

$$\begin{aligned} u(x_1, x_2) &= u(x^1(s), x^2(s)) \\ &= \frac{(x_1 + 4x_2)^2}{16}. \end{aligned}$$

## 2.4 Boundary conditions

### Straightening the boundary

We intend in the following section to invoke the characteristic ODE (2.7) to actually solve the boundary-value problem (2.4), (2.5), at least in a small region near an appropriate portion  $\Gamma$  of  $\partial U$ . In order to simplify the relevant calculations, it is convenient first fix any point  $x^0 \in \partial U$ . Then utilizing the notation from the appendix §C.1 of [3], we find smooth mappings  $\Phi, \Psi: \mathbf{R}^n \rightarrow \mathbf{R}^n$  such that  $\Psi = \Phi^{-1}$  and  $\Phi$  straightens  $\partial U$  near  $x^0$ .

Given a function  $u: U \rightarrow \mathbf{R}$ , let us write  $V := \Phi(U)$  and set

$$v(y) := u(\Psi(y)) \quad y \in V. \quad (2.16)$$

Then

$$u(x) = v(\Phi(x)) \quad x \in U. \quad (2.17)$$

Now suppose that  $u$  is a  $C^1$  solution of our boundary-value problem (2.4), (2.5) in  $U$ . What PDE does  $v$  then satisfy in  $V$ ?

According to (2.17), we have

$$u_{x_i}(x) = \sum_{k=1}^n v_{y_k}(\Phi(x)) \Phi_{x_i}^k(x)$$

i.e.,

$$Du(x) = Dv(y)D\Phi(x).$$

Thus,

$$\begin{aligned} 0 &= F(Du(x), u(x), x) \\ &= F(Dv(y), D\Phi(\Psi(y)), v(y), \Psi(y)). \end{aligned}$$

In addition  $v = h$  on  $\Delta$ , where  $\Delta := \Phi(\Gamma)$  and  $h(y) := g(\Psi(y))$ .

In summary, our problem (2.4), (2.5) converts into a problem having the same form.

### Compatibility conditions on boundary

In view of the foregoing computations, if we are given a point  $x^0 \in \Gamma$  we may as well assume that the outset that  $\Gamma$  is flat near  $x^0$ , lying in the plane  $\{x_n = 0\}$ .

We intend now to utilize the characteristic ODE to construct a solution to (2.4), at least near  $x^0$ , and for this we must discover appropriate initial conditions

$$\mathbf{p}(0) = p^0, \quad z(0) = z^0, \quad \mathbf{x}(0) = x^0. \quad (2.18)$$

Now clearly if the curve  $\mathbf{x}$  passes through  $x^0$ , we should insist that

$$z^0 = g(x^0). \quad (2.19)$$

What should we require concerning  $\mathbf{p}(0) = p^0$ ? Since (2.5) implies  $u(x_1, \dots, x_{n-1}, 0) = g(x_1, \dots, x_{n-1})$  near  $x^0$ , we may differentiate to find

$$u_{x_i}(x^0) = g_{x_i}(x^0) \quad i = 1, \dots, n-1.$$

As we also want the PDE (2.4) to hold, we should therefore insist  $p^0 = (p_1^0, \dots, p_n^0)$  satisfies these relations

$$\begin{cases} p_i^0 = g_{x_i}(x^0) & i = 1, \dots, n-1 \\ F(p^0, z^0, x^0) = 0. \end{cases} \quad (2.20)$$

These identities provide  $n$  equations for the  $n$  quantities  $p^0 = (p_1^0, \dots, p_n^0)$ .

We call (2.19) and (2.20) the *compatibility conditions*. A triple  $(p^0, z^0, x^0) \in \mathbf{R}^{2n+1}$  verifying (2.19), (2.20) is *admissible*. Note that  $z^0$  is uniquely determined by the boundary condition and our choice of the point  $x^0$ , but a vector  $p^0$  satisfying (2.20) may not exist and it may not be unique.

### Noncharacteristic boundary data

So now assume as above that  $x^0 \in \Gamma$ , that  $\Gamma$  near  $x^0$  lies in the plane  $\{x_n = 0\}$ , and that the triple  $(p^0, z^0, x^0)$  is admissible. We are planning to construct a solution  $u$  of (2.4), (2.5) in  $U$  near  $x^0$  by integrating by parts the characteristic ODE (2.7). So far we have ascertained  $\mathbf{x}(0) = x^0$ ,  $z(0) = z^0$ ,  $\mathbf{p}(0) = p^0$  are appropriate boundary conditions for the characteristic ODE, with  $\mathbf{x}$  intersecting  $\Gamma$  at  $x^0$ . But we will need in fact to solve these ODEs for *nearby* initial points as well, and must consequently now ask if we can somehow appropriately perturb  $(p^0, z^0, x^0)$ , keeping the compatibility conditions.

In other words, given a point  $y = (y_1, \dots, y_{n-1}, 0) \in \Gamma$ , with  $y$  close to  $x^0$ , we intend to solve the characteristic ODE

$$\begin{cases} \text{(a) } \dot{\mathbf{p}}(s) = -D_x F(\mathbf{p}(s), z(s), \mathbf{x}(s)) - D_z F(\mathbf{p}(s), z(s), \mathbf{x}(s)) \\ \text{(b) } \dot{z}(s) = D_p F(\mathbf{p}(s), z(s), \mathbf{x}(s)) \cdot \mathbf{p}(s) \\ \text{(c) } \dot{\mathbf{x}}(s) = D_p F(\mathbf{p}(s), z(s), \mathbf{x}(s)), \end{cases} \quad (2.21)$$

with the initial conditions

$$\mathbf{p}(0) = \mathbf{q}(y), \quad z(0) = g(y), \quad \mathbf{x}(0) = y. \quad (2.22)$$

Our task then is to find a function  $\mathbf{q} = (q^1, \dots, q^n)$ , so that

$$\mathbf{q}(x^0) = p^0 \quad (2.23)$$

and  $(\mathbf{q}(y), g(y), y)$  is admissible; that is, the compatibility conditions

$$\begin{cases} q^i(y) = g_{x_i}(y) & 1 \leq i \leq n-1 \\ F(\mathbf{q}(y), g(y), y) = 0 \end{cases} \quad (2.24)$$

hold for all  $y \in \Gamma$  close to  $x^0$ .

**Lemma 2.9** (Noncharacteristic boundary conditions). *There exists a unique solution  $\mathbf{q}$  of (2.23), (2.24) for all  $y \in \Gamma$  sufficiently close to  $x^0$ , provided*

$$F_{p_n}(p^0, z^0, x^0) \neq 0. \quad (2.25)$$

We say the admissible triple  $(p^0, z^0, x^0)$  is *noncharacteristic* if (2.25) holds. We henceforth assume this condition.

*PROOF.* To simplify notation, let us now temporarily write  $y = (y_1, \dots, y_n) \in \mathbf{R}^n$ . We apply the implicit function theorem to the mapping

$$\mathbf{G}: \mathbf{R}^n \times \mathbf{R}^n \longrightarrow \mathbf{R}^n, \quad \mathbf{G}(p, y) = (G^1(p, y), \dots, G^n(p, y)),$$

where

$$\begin{cases} G^i(p, y) = p_i - g_{x_i}(y) & 1 \leq i \leq n-1, \\ G^n(p, y) = F(p, g(y), y). \end{cases}$$

Now  $\mathbf{G}(p^0, x^0) = 0$ , according to (2.19), (2.18). Also

$$D_p \mathbf{G}(p^0, x^0) = \begin{bmatrix} 1 & \cdots & 0 & 0 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \cdots & 1 & 0 \\ F_{p_1}(p^0, z^0, x^0) & \cdots & F_{p_{n-1}}(p^0, z^0, x^0) & F_{p_n}(p^0, z^0, x^0) \end{bmatrix}$$

and thus

$$\det D_p \mathbf{G}(p^0, x^0) = F_{p_n}(p^0, z^0, x^0) \neq 0,$$

in view of the noncharacteristic condition (2.25). ■

## Chapter 3

# Algebraic Geometry

A summary to a course on an introduction to sheaf cohomology. We will mostly reference Donu's notes available here <https://www.math.purdue.edu/~dvb/classroom.html>, but also cite Ravi Vakil's *Fundamentals of Algebraic Geometry* [10] available here <https://math216.wordpress.com/>.

### 3.1 The statement of de Rham's theorem

These are almost verbatim Arapura's notes on the de Rham Complex and cohomology.

Before doing anything fancy, let's start at the beginning. Let  $U \subseteq \mathbf{R}^3$  be an open set. In calculus class, we learn about operations

$$\{ \text{functions} \} \xrightarrow{\nabla} \{ \text{vector fields} \} \xrightarrow{\nabla \times} \{ \text{vector fields} \} \xrightarrow{\nabla \cdot} \{ \text{functions} \}$$

such that  $(\nabla \times)(\nabla) = 0$  and  $(\nabla \cdot)(\nabla \times) = 0$ . This is a prototype for a *complex*. An obvious question: does  $\nabla \times v = 0$  imply that  $v$  is a gradient? Answer: sometimes yes (e.g. if  $U = \mathbf{R}^3$ ) and sometimes no (e.g. if  $U = \mathbf{R}^3$  minus a line).

## Chapter 4

# Algebraic Topology

From my meetings with Mark. We reference Hatcher's *Algebraic Topology* [6] freely available here <https://www.math.cornell.edu/~hatcher/#ATI>.

### 4.1 Cohomology



## Chapter 5

# Group Theory and Differential Equations

This is a summary of Kuga's *Galois' Dream: Group Theory and Differential Equations* book.

# Bibliography

- [1] R. Bott and L.W. Tu. *Differential Forms in Algebraic Topology*. Graduate Texts in Mathematics. Springer New York, 2013.
- [2] A. DasGupta. *Fundamentals of Probability: A First Course*. Springer Texts in Statistics. Springer New York, 2010.
- [3] L.C. Evans. *Partial Differential Equations*. Graduate studies in mathematics. American Mathematical Society, 2010.
- [4] W. Feller. *An introduction to probability theory and its applications*. Number v. 1 in Wiley mathematical statistics series. Wiley, 1950.
- [5] J. Harris. *Algebraic Geometry: A First Course*. Graduate Texts in Mathematics. Springer New York, 2013.
- [6] A. Hatcher. *Algebraic Topology*. Cambridge University Press, 2002.
- [7] D. Mumford. *The Red Book of Varieties and Schemes*. Blurb, Incorporated, 2015.
- [8] S.M. Ross. *A First Course in Probability*. Pearson Prentice Hall, 2010.
- [9] W.A. Strauss. *Partial Differential Equations: An Introduction*. Wiley, 1992.
- [10] R. Vakil. Math 216: Foundations of algebraic geometry, 2016.