

MA571 Homework 9

Carlos Salinas

October 31, 2015

PROBLEM 9.1 (MUNKRES §52, EX. 2)

Let α be a path in X from x_0 to x_1 ; let β be a path in X from x_1 to x_2 . Show that if $\gamma = \alpha * \beta$, then $\hat{\gamma} = \hat{\beta} \circ \hat{\alpha}$.

Proof.

■

PROBLEM 9.2 (MUNKRES §52, EX. 3)

Let x_0 and x_1 be points of the path-connected space X . Show that $\pi_1(X, x_0)$ is Abelian if and only if for every pair α and β of paths from x_0 to x_1 , we have $\hat{\alpha} = \hat{\beta}$.

Proof.

■

PROBLEM 9.3 (MUNKRES §52, EX. 4)

Let $A \subset X$; suppose $r: X \rightarrow A$ is continuous map such that $r(a) = a$ for each $a \in A$. (The map r is called a *retraction* of X onto A .) If $a_0 \in A$, show that

$$r_*: \pi_1(X, x_0) \longrightarrow \pi_1(A, a_0)$$

is surjective.

Proof.

■

PROBLEM 9.4 (MUNKRES §55, EX. 1)

Show that if A is a retract of B^2 , then every continuous map $f: A \rightarrow A$ has a fixed point.

Proof.

■

PROBLEM 9.5 (MUNKRES §55, EX. 2)

Show that if $h: S^1 \rightarrow S^1$ is nullhomotopic, then h has a fixed point and h maps some point x to its antipode $-x$.

Proof.

■

PROBLEM 9.6

Proof.



PROBLEM 9.7 ((A))

Prove that every m -manifold is locally path-connected.

Proof.

■

PROBLEM 9.8 ((B))

Prove that every m -manifold is regular.

Proof.

■

PROBLEM 9.9 ((C))

Prove that there is no 1-1 continuous function $\iota: S^1 \rightarrow \mathbb{R}$. You may assume any fact about trigonometric functions. (Note: this shows in particular that there is no $\iota: S^1 \rightarrow \mathbb{R}$ with $p \circ \iota$ equal to the identity map, where p is the map in the note on the Fundamental Group of the Circle.)

Proof.

