

# MA 523: Homework 4

Carlos Salinas

September 28, 2016



## PROBLEM 4.1 (LEGENDRE TRANSFORM)

Let  $u(x_1, x_2)$  be a solution of the quasilinear equation

$$a^{11}(Du)u_{x_1x_1} + 2a^{12}(Du)u_{x_1x_2} + a^{22}(Du)u_{x_2x_2} = 0$$

in some region of  $\mathbb{R}^2$ , where we can invert the relations

$$p^1 = u_{x_1}(x_1, x_2), \quad p^2 = u_{x_2}(x_1, x_2)$$

to solve for

$$x^1 = x^1(p_1, p_2), \quad x^2 = x^2(p_1, p_2).$$

Define then

$$v(p) := \mathbf{x}(p) \cdot p - u(\mathbf{x}(p)),$$

where  $\mathbf{x} = (x^1, x^2)$ ,  $p = (p_1, p_2)$ . Show that  $v$  satisfies the *linear* equation

$$a^{22}(p)v_{p_1p_2} - 2a^{12}(p)v_{p_1p_2} + a^{11}(p)v_{p_1p_2} = 0.$$

(*Hint:* See [Evans, 4.4.3b], prove the identities (29)).

SOLUTION. ■

## PROBLEM 4.2

Find the solution  $u(x, t)$  of the one-dimensional wave equation

$$u_{tt} - u_{xx} = 0$$

in the quadrant  $x > 0, t > 0$  for which

$$\begin{cases} u(x, 0) = f(x), & u_t(x, 0) = g(x), & \text{for } x > 0 \\ u_t(0, t) = \alpha u_x(0, t), & & \text{for } t > 0, \end{cases}$$

where  $\alpha \neq -1$  is a given constant. Show that generally no solution exists when  $\alpha = -1$ . (*Hint:* Use a representation  $u(x, t) = F(x - t) + G(x + t)$  for the solution.)

*SOLUTION.*

■

## PROBLEM 4.3

- (a) Let  $u$  be a solution of the wave equation  $u_{tt} - c^2 u_{xx} = 0$  for  $0 < x < \pi$ ,  $t > 0$  such that  $u(0, t) = u(\pi, t) = 0$ . Show that the *energy*

$$E(t) = \frac{1}{2} \int_0^\pi (u_t^2 + c^2 u_x^2) dx, \quad t > 0$$

is independent of  $t$ ; i.e.,  $\frac{d}{dt} E = 0$  for  $t > 0$ . Assume that  $u$  is  $C^2$  up to the boundary.

- (b) Express the energy  $E$  of the Fourier series solution

$$u(x, t) = \sum_{n=1}^{\infty} (a_n \cos(nct) + b_n \sin(nct)) \sin(nx)$$

in terms of coefficients  $a_n$ ,  $b_n$ .

*SOLUTION.* For part (a), suppose that  $u$  is, as above, a solution to the wave equation which is  $C^2$  up to the boundary. We show that its energy is independent of  $t$ , i.e., that  $\frac{d}{dt} E = 0$ . Assuming the energy is bounded, the dominated convergence theorem allows us to permute the order of integration and differentiation like so

$$\begin{aligned} \frac{d}{dt} E(t) &= \frac{d}{dt} \left( \frac{1}{2} \int_0^\pi (u_t^2 + c^2 u_x^2) dx \right) \\ &= \frac{1}{2} \int_0^\pi \frac{\partial}{\partial t} (u_t^2 + c^2 u_x^2) dx \\ &= \frac{1}{2} \int_0^\pi 2u_t u_{tt} + 2c^2 u_x u_{xt} dx \end{aligned}$$

which, after using the relation  $u_{tt} = c^2 u_{xx}$ , becomes

$$\begin{aligned} &= c^2 \int_0^\pi u_t u_{xx} + u_x u_{xt} dx \\ &= c^2 \int_0^\pi \frac{\partial}{\partial x} (u_x u_t) dx \\ &= c^2 (u_x(\pi, t) u_t(\pi, t) - u_x(0, t) u_t(0, t)) \\ &= 0 \end{aligned}$$

since the boundary conditions, i.e.,  $u = 0$ , implies  $u_x = u_t = 0$  at the boundary.

For part (b), suppose  $u$  is a Fourier series solution to the wave equation, i.e.,

$$u(x, t) = \sum_{n=1}^{\infty} (a_n \cos(nct) + b_n \sin(nct)) \sin(nx).$$

First we compute  $u_t$  and  $u_x$ . They are

$$\begin{aligned} u_t(x, t) &= \frac{\partial}{\partial t} u(x, t) \\ &= \sum_{n=1}^{\infty} cn(b_n \cos(nct) - a_n \sin(nct)) \sin(nx) \end{aligned}$$

and

$$\begin{aligned} u_x(x, t) &= \frac{\partial}{\partial x} u(x, t) \\ &= \sum_{n=1}^{\infty} n(a_n \cos(nct) + b_n \sin(nct)) \cos(nx). \end{aligned}$$

Thus,

$$\begin{aligned} E(t) &= \frac{1}{2} \int_0^\pi \left[ \left( \sum_{n=1}^{\infty} cn(b_n \cos(nct) - a_n \sin(nct)) \sin(nx) \right)^2 \right. \\ &\quad \left. + c^2 \left( \sum_{n=1}^{\infty} n(a_n \cos(nct) + b_n \sin(nct)) \cos(nx) \right)^2 \right] \end{aligned}$$

which, after expanding and using the fact that  $\cos(nct)$ ,  $\sin(nct)$ ,  $\cos(nx)$ , and  $\sin(nx)$  are orthogonal, becomes

$$\begin{aligned} &= \frac{1}{2} \int_0^\pi \left[ \sum_{n,m=1}^{\infty} c^2 nm(b_n b_m \cos(nct) \cos(mct) + a_n a_m \sin(nct) \sin(mct) \right. \\ &\quad \left. - a_n b_m \cos(mct) \sin(nct) - a_m b_n \cos(nct) \sin(mct)) \sin(nx) \sin(mx) \right. \\ &\quad \left. - c^2 \sum_{n,m=1}^{\infty} n^2 (a_n a_m \cos(nct) \cos(mct) + b_n b_m \sin(nct) \sin(mct) \right. \\ &\quad \left. + a_n b_m \cos(nct) \sin(mct) + a_m b_n \cos(mct) \sin(nct)) \cos(nx) \cos(mx) \right] \\ &= \frac{1}{2} \int_0^\pi \sum_{n=1}^{\infty} \left( cn^2 (b_n^2 \cos^2(nct) + a_n^2 \sin^2(nct)) \sin^2(nx) - cn^2 (a_n^2 \cos^2(nct) + b_n^2 \sin^2(nct)) \cos^2(nx) \right) \end{aligned}$$

Consider the map  $XY$  ■