

MA 523: Homework 1

Carlos Salinas

August 25, 2016

PROBLEM 1.1 (TAYLOR'S FORMULA)

Let $f: \mathbb{R}^n \rightarrow \mathbb{R}$ be smooth, $n \geq 2$. Prove that

$$f(x) = \sum_{|\alpha| \leq k} \frac{1}{\alpha!} D^\alpha f(0) x^\alpha + O(|x|^{k+1})$$

as $x \rightarrow 0$ for each $k = 1, 2, \dots$, assuming that you know this formula for $n = 1$.

Hint: Fix $x \in \mathbb{R}^n$ and consider the function of one variable $g(t) := f(tx)$. Prove that

$$\frac{d^m}{dt^m} g(t) = \sum_{|\alpha|=m} \frac{m!}{\alpha!} D^\alpha f(tx) x^\alpha,$$

by induction on m .

Solution. ► Taking the hint, fix $x \in \mathbb{R}^n$ and consider the function of one variable $g(t) := f(tx)$. We claim that

$$\frac{d^m}{dt^m} g(t) = \sum_{|\alpha|=m} \frac{m!}{\alpha!} D^\alpha f(tx) x^\alpha.$$

Proof of claim. We shall prove this by induction on m . The case $m = 1$ follows from Taylor's formula in 1 dimension as

$$\begin{aligned} \frac{d}{dt} g(t) &= \frac{d}{dt} f(tx) \\ &= \frac{d}{dt} f(tx_1, \dots, tx_n) \\ &= \end{aligned}$$

□

◀

PROBLEM 1.2

Write down the characteristic equation for the p.d.e.

$$u_t + b \cdot Du = f \tag{*}$$

on $\mathbb{R}^n \times (0, \infty)$, where $b \in \mathbb{R}^n$. Using the characteristic equation, solve (*) subject to the initial condition

$$u = g$$

on $\mathbb{R}^n \times \{t = 0\}$. Make sure the answer agrees with formula (5) in §2.1.2 of [E].

Solution. ►

◀

PROBLEM 1.3

Solve using the characteristics:

(a) $x_1^2 u_{x_1} + x_2^2 u_{x_2} = u^2$, $u = 11$ on the line $x_2 = 2x_1$.

(b) $uu_{x_1} + u_{x_2} = 1$, $u(x_1, x_2) = x_1/2$.

(c) $x_1 u_{x_1} + 2x_2 u_{x_2} + u_{x_3} = 3u$, $u(x_1, x_2, 0) = g(x_1, x_2)$.

Solution. ►

◀

PROBLEM 1.4

For the equation

$$u = x_1 u_{x_1} + x_2 u_{x_2} + \frac{1}{2}(u_{x_1}^2 + u_{x_2}^2)$$

find a solution with $u(x_1, 0) = (1 - x_1^2)/2$.

Solution. ►

◀