

MA557 Homework 10

Carlos Salinas

November 16, 2015

PROBLEM 10.1

Let $\varphi: R \rightarrow S$ be a homomorphism of rings, ${}_a\varphi: \operatorname{Spec} S \rightarrow \operatorname{Spec} R$ the induced map of the spectra, and $\mathfrak{p} \in \operatorname{Spec} R$. Show that the fiber $({}_a\varphi)^{-1}(\mathfrak{p})$ is homeomorphic to $\operatorname{Spec}(S \otimes_R k(\mathfrak{p}))$.

Proof.

■

PROBLEM 10.2

Let $R \subset S$ be an integral extension of rings with S a Noetherian ring, and let $\mathfrak{p} \in \operatorname{Spec} R$. Show that there are only finitely many primes in S lying over \mathfrak{p} .

Proof.

■

PROBLEM 10.3

Let $\varphi: R \rightarrow S$ be a homeomorphism of rings with S a Noetherian ring. Show that the following are equivalent:

- (i) φ satisfies going up.
- (ii) ${}_a\varphi: \operatorname{Spec} S \rightarrow \operatorname{Spec} R$ is a closed map.
- (iii) for every $\mathfrak{q} \in \operatorname{Spec} S$, the induced map $\operatorname{Spec}(S/\mathfrak{q}) \rightarrow \operatorname{Spec}(R/\mathfrak{q} \cap S)$ is surjective.

Proof.

■

PROBLEM 10.4

Let $R \subset S$ be an integral extension of domains with R normal, $K = \text{Quot } R$, $\alpha \in S$, $X^n + a_1X^{n-1} + \cdots + a_n$ the minimal polynomial of α over K (recall $a_i \in R$). Show that for any R -ideal I , $\alpha \in \sqrt{IS}$ if and only if $a_i \in \sqrt{I}$ for $1 \leq i \leq n$.

Proof.

■

PROBLEM 10.5

Let k be a field and $R = k[X_1, \dots, X_n]$ a k -algebra. Show that the following are equivalent:

- (i) R is a domain with $\dim R = n - 1$
- (ii) $R \cong k[X_1, \dots, X_n]/(f)$, where $k[X_1, \dots, X_n]$ is a polynomial ring and f is an irreducible polynomial.

Proof.

■