

MA 523: Homework 7

Carlos Salinas

October 31, 2016

PROBLEM 7.1

Solve the Dirichlet problem for the Laplace equation in \mathbb{R}^2

$$\begin{cases} \Delta u = 0 & \text{in } 1 < |x| < 2, \\ u = x_1 & \text{on } |x| = 1, \\ u = 1 + x_1 x_2 & \text{on } |x| = 2. \end{cases}$$

(*Hint:* Use Laurent series.)

SOLUTION. Suppose u is a solution to the Dirichlet problem above with the form

$$u(x_1, x_2) = a \log |x| + \operatorname{Re}(L(x_1, x_2))$$

where

$$L(x_1, x_2) = \sum_{n \in \mathbb{Z}} a_n (x_1 + x_2 i)^n$$

is an honest Laurent series. ■

PROBLEM 7.2

Let Ω be a bounded domain with a C^1 boundary, $g \in C^2(\partial\Omega)$ and $f \in C(\bar{\Omega})$. Consider the so called *Neumann problem*

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ \frac{\partial u}{\partial \nu} = g & \text{on } \partial\Omega, \end{cases} \quad (*)$$

where ν is the outer normal on $\partial\Omega$. Show that the solution of $(*)$ in $C^2(\Omega) \cap C^1(\bar{\Omega})$ is unique up to a constant; i.e., if u_1 and u_2 are both solutions of $(*)$, then $u_2 = u_1 + \text{const.}$ in Ω .

(*Hint*: Look at the proof of the uniqueness for the Dirichlet problem by energy methods [E, 2.2.5a].)

SOLUTION. Suppose u_1 and u_2 are solutions to the Neumann problem $(*)$. Define $v := u_1 - u_2$. Then v is harmonic in Ω and $\frac{\partial v}{\partial \nu} = 0$ on $\partial\Omega$. Consider the energy functional

$$E[v] = \frac{1}{2} \int_{\Omega} |Dv|^2 dx.$$

By Green's formula version (ii),

$$\begin{aligned} E[v] &= \frac{1}{2} \int_{\Omega} |Dv|^2 dx \\ &= -\frac{1}{2} \int_{\Omega} v \Delta v dx + \int_{\partial\Omega} \frac{\partial v}{\partial \nu} v dS(x) \\ &= 0. \end{aligned}$$

This implies that $|Dv|^2 = Dv \cdot Dv = 0$ which, since the standard inner product on \mathbb{R}^n is positive-definite, implies that $Dv \equiv 0$. It follows that $u_1 = u_2 + \text{const.}$, i.e., the solution u to $(*)$ is unique up to a constant. ■

PROBLEM 7.3

Write down an explicit formula for a solution of

$$\begin{cases} u_t - \Delta u + cu = f & \text{in } \mathbb{R}^n \times (0, \infty), \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}, \end{cases}$$

where $c \in \mathbb{R}$.

(*Hint:* Rewrite the problem in terms of $v(x, t) := e^{ct}u(x, t)$.)

SOLUTION.

■