

MA 519: Homework 11

Max Jeter, Carlos Salinas

November 10, 2016

PROBLEM 11.1 (DASGUPTA 7.2 (A), (B), (C), (D), (E))

- (a) Suppose $E|X_n - c|^\alpha \rightarrow 0$, where $0 < \alpha < 1$. Does X_n necessarily converge in probability to c ?
- (b) Suppose $a_n(X_n - \theta) \xrightarrow{\mathcal{L}} \mathcal{N}(0, 1)$. Under what condition on a_n can we conclude that $X_n \xrightarrow{\mathcal{P}} \theta$?
- (c) $\text{o}_p(1) + \text{O}_p(1) = ?$
- (d) $\text{o}_p(1)\text{O}_p(1) = ?$
- (e) $\text{o}_p(1) + \text{o}_p(1)\text{O}_p(1) = ?$

SOLUTION. For part (a), we provide the following counterexample. Let X have the distribution ■

PROBLEM 11.2 (DASGUPTA 7.3 [MONTE CARLO])

Consider the purely mathematical problem of finding a definite integral $\int f(x) dx$ for some (possibly complicated) function $f(x)$. Show that the SLLN provides a method for approximately finding the value of the integral by using appropriate averages $\frac{1}{n} \sum_{k=1}^n f(X_k)$.

Numerical analysts call this Monte Carlo integration.

SOLUTION.

■

PROBLEM 11.3 (DASGUPTA 7.4 (A), (B))

Suppose X_1, \dots , are i.i.d. and that $E(X_1) = \mu \neq 0$, $\text{Var}(X_1) = \sigma^2 < \infty$. Let $S_{m,p} = \sum_{k=1}^m X_k^p$, $m \geq 1$, $p = 1, 2$.

- (a) Identify with proof the almost sure limit of $S_{m,1}/S_{n,1}$ for fixed m , and $n \rightarrow \infty$.
- (b) Identify with proof the almost sure limit of $S_{n-m,1}/S_{n,1}$ for fixed m , and $n \rightarrow \infty$.

SOLUTION. ■

PROBLEM 11.4 (DASGUPTA 7.5 (A))

Let A_n , $n \geq 1$, A be events with respect to a common sample space Ω .

- (a) Prove that $I_{A_n} \xrightarrow{\mathcal{L}} I_A$ if and only if $P(A_n) \rightarrow P(A)$.

SOLUTION.

■

PROBLEM 11.5 (DASGUPTA 7.11 [SAMPLE MAXIMUM])

Let X_k , $k \geq 1$, be an i.i.d. sequence, and $X_{(n)}$ the maximum of X_1, \dots, X_n . Let $\xi(F) = \sup\{x : F(x) < 1\}$, where F is the common CDF of the X_k . Prove that $X_{(n)} \xrightarrow{\text{a.s.}} \xi(F)$.

SOLUTION. ■

PROBLEM 11.6 (DASGUPTA 7.14 (A))

Suppose X_k are i.i.d. standard Cauchy. Show that

- (a) $P(|X_n| > n \text{ infinitely often}) = 1$.

SOLUTION.

■

PROBLEM 11.7 (DASGUPTA 7.16 [COUPON COLLECTION])

Cereal boxes contain independently and with equal probability exactly one of n different celebrity pictures. Someone having the entire set of n pictures can cash them in for money. Let W_n be the minimum number of cereal boxes one would need to purchase to own a complete set of the pictures.

Find a sequence a_n such that $W_n/a_n \xrightarrow{\mathcal{P}} 1$.

(*Hint:* Approximate the mean of W_n .)

SOLUTION. ■

PROBLEM 11.8 (DASGUPTA 7.17)

Let $X \sim \text{Bin}(n, p)$. Show that $(X_n/n)^2$ and $X_n(X_n - 1)/(n(n - 1))$ both converging in probability to p^2 . Do they converge almost surely?

SOLUTION. ■

PROBLEM 11.9 (DASGUPTA 7.21)

Let X_1, X_2, \dots , be i.i.d. $U[0, 1]$. Let

$$G_n = (X_1 \cdots X_n)^{1/n}.$$

Find c such that $G_n \xrightarrow{\mathcal{P}} c$.

SOLUTION.

■

PROBLEM 11.10 (DASGUPTA 7.30 [CONCEPTUAL])

Suppose $X_n \xrightarrow{\mathcal{L}} X$, and also $Y_n \xrightarrow{\mathcal{L}} X$. Does this mean that $X_n - Y_n$ converge in distribution to (the point mass at) zero?

SOLUTION. ■

PROBLEM 11.11 (DASGUPTA 7.31 (A))

- (a) Suppose $a_n(X_n - \theta) \rightarrow \mathcal{N}(0, \tau^2)$; what can be said about the limiting distribution of $|X_n|$, when $\theta \neq 0$, $\theta = 0$?

SOLUTION. ■