

MA 544: Homework 3

Carlos Salinas

January 26, 2016

PROBLEM 3.1 (WHEEDEN & ZYGMUND §3, EX. 5)

Construct a subset of $[0, 1]$ in the same manner as the Cantor set, except that at the k th stage each interval removed has length $\delta 3^{-k}$, $0 < \delta < 1$. Show that the resulting set is perfect, has measure $1 - \delta$, and contains no interval.

Proof.

■

PROBLEM 3.2 (WHEEDEN & ZYGMUND §3, EX. 7)

Prove (3.15).

Proof.

■

PROBLEM 3.3 (WHEEDEN & ZYGMUND §3, EX. 9)

If $\{E_k\}_{k=1}^{\infty}$ is a sequence of sets with $\sum |E_k|_e < +\infty$, show that $\limsup E_k$ (and also $\liminf E_k$) has measure zero.

Proof.

■

PROBLEM 3.4 (WHEEDEN & ZYGMUND §3, EX. 12)

If E_1 and E_2 are measurable subsets of \mathbb{R}^1 , show that $E_1 \times E_2$ is measurable subset of \mathbb{R}^2 and $|E_1 \times E_2|_e = |E_1||E_2|_e$. (Interpret $0 \cdot \infty$ as 0) [HINT: Use a characterization of measurability.]

Proof.

■

PROBLEM 3.5 (WHEEDEN & ZYGMUND §3, EX. 13)

Motivated by (3.7), define the *inner measure of E* to be $|E|_i := \sup|F|$, where the supremum is taken over all closed subsets F of E . Show that

- (i) $|E|_i \leq |E|_e$, and
- (ii) if $|E|_e < +\infty$, then E is measurable if and only if $|E|_e = |E|_i$. [Use (3.22).]

Proof.

■

PROBLEM 3.6 (WHEEDEN & ZYGMUND §3, EX. 14)

Show that the conclusion of part (ii) of Exercise 13 (Problem) is false if $|E|_e = +\infty$.

Proof.

■

PROBLEM 3.7 (WHEEDEN & ZYGMUND §3, EX. 8)

Show that the Borel σ -algebra \mathcal{B} in \mathbb{R}^n is the smallest σ -algebra containing the closed sets in \mathbb{R}^n .

Proof.

■

PROBLEM 3.8 (WHEEDEN & ZYGMUND §3, EX. 10)

Proof.



PROBLEM 3.9 (WHEEDEN & ZYGMUND §3, EX. 15)

Proof.



PROBLEM 3.10 (WHEEDEN & ZYGMUND §3, EX. 16)

Proof.



PROBLEM 3.11 (WHEEDEN & ZYGMUND §3, EX. 18)

Proof.

