

Rank 1 Character Varieties-Part II Generators

Sean Lawton

MRC - Snowbird

June 2016

$SL(2, \mathbb{C})$ and free groups.

- 1 Let $F_r = \langle \gamma_1, \dots, \gamma_r \rangle$ denote the free group of rank r

$SL(2, \mathbb{C})$ and free groups.

- 1 Let $F_r = \langle \gamma_1, \dots, \gamma_r \rangle$ denote the free group of rank r , so it is group of all strings in the symbols $\{\gamma_1^{\pm 1}, \dots, \gamma_r^{\pm 1}\}$.

$SL(2, \mathbb{C})$ and free groups.

- 1 Let $F_r = \langle \gamma_1, \dots, \gamma_r \rangle$ denote the free group of rank r , so it is group of all strings in the symbols $\{\gamma_1^{\pm 1}, \dots, \gamma_r^{\pm 1}\}$.
- 2 $\text{Hom}(F_r, G) \cong G^{\times r}$ by the evaluation map.

SL(2, \mathbb{C}) and free groups.

- 1 Let $F_r = \langle \gamma_1, \dots, \gamma_r \rangle$ denote the free group of rank r , so it is group of all strings in the symbols $\{\gamma_1^{\pm 1}, \dots, \gamma_r^{\pm 1}\}$.
- 2 $\text{Hom}(F_r, G) \cong G^{\times r}$ by the evaluation map.
- 3 Therefore $\text{Hom}(F_r, G)$ is an affine variety:

$$\{(v_{11}^1, v_{12}^1, v_{21}^1, v_{22}^1, \dots, v_{11}^r, v_{12}^r, v_{21}^r, v_{22}^r) \in \mathbb{C}^{4r} \mid v_{11}^k v_{22}^k - v_{12}^k v_{21}^k = 1, 1 \leq k \leq r\}.$$

SL(2, \mathbb{C}) and free groups.

- 1 Let $F_r = \langle \gamma_1, \dots, \gamma_r \rangle$ denote the free group of rank r , so it is group of all strings in the symbols $\{\gamma_1^{\pm 1}, \dots, \gamma_r^{\pm 1}\}$.
- 2 $\text{Hom}(F_r, G) \cong G^{\times r}$ by the evaluation map.
- 3 Therefore $\text{Hom}(F_r, G)$ is an affine variety:

$$\{(v_{11}^1, v_{12}^1, v_{21}^1, v_{22}^1, \dots, v_{11}^r, v_{12}^r, v_{21}^r, v_{22}^r) \in \mathbb{C}^{4r} \mid v_{11}^k v_{22}^k - v_{12}^k v_{21}^k = 1, 1 \leq k \leq r\}.$$

- 4 Therefore it has a coordinate ring $\mathbb{C}[\text{Hom}(F_r, \text{SL}(2, \mathbb{C}))] \cong \mathbb{C}[x_{ij}^k \mid 1 \leq i, j \leq 2, 1 \leq k \leq r] / \langle x_{11}^k x_{22}^k - x_{12}^k x_{21}^k - 1 \mid 1 \leq k \leq r \rangle.$

- 5 G acts on $\text{Hom}(F_r, G)$ by $g \cdot \rho = g\rho g^{-1}$; or equivalently on $G^{\times r}$ by $g \cdot (g_1, \dots, g_r) = (gg_1g^{-1}, \dots, gg_rg^{-1})$.

- 5 G acts on $\text{Hom}(F_r, G)$ by $g \cdot \rho = g\rho g^{-1}$; or equivalently on $G^{\times r}$ by $g \cdot (g_1, \dots, g_r) = (gg_1g^{-1}, \dots, gg_rg^{-1})$.
- 6 So this determines a co-action on the ring of polynomials $\mathbb{C}[\text{Hom}(F_r, G)]$ by $g \cdot f(\rho) = f(g^{-1}\rho g)$.

- 5 G acts on $\text{Hom}(F_r, G)$ by $g \cdot \rho = g\rho g^{-1}$; or equivalently on $G^{\times r}$ by $g \cdot (g_1, \dots, g_r) = (gg_1g^{-1}, \dots, gg_rg^{-1})$.
- 6 So this determines a co-action on the ring of polynomials $\mathbb{C}[\text{Hom}(F_r, G)]$ by $g \cdot f(\rho) = f(g^{-1}\rho g)$.
- 7 Since G is reductive, the ring of invariants $\mathbb{C}[\text{Hom}(F_r, G)]^G := \{f \in \mathbb{C}[\text{Hom}(F_r, G)] \mid g \cdot f = f\}$ is finitely generated.

- 5 G acts on $\text{Hom}(F_r, G)$ by $g \cdot \rho = g\rho g^{-1}$; or equivalently on $G^{\times r}$ by $g \cdot (g_1, \dots, g_r) = (gg_1g^{-1}, \dots, gg_rg^{-1})$.
- 6 So this determines a co-action on the ring of polynomials $\mathbb{C}[\text{Hom}(F_r, G)]$ by $g \cdot f(\rho) = f(g^{-1}\rho g)$.
- 7 Since G is reductive, the ring of invariants $\mathbb{C}[\text{Hom}(F_r, G)]^G := \{f \in \mathbb{C}[\text{Hom}(F_r, G)] \mid g \cdot f = f\}$ is finitely generated. (we will prove this in our special case)

- 5 G acts on $\mathrm{Hom}(F_r, G)$ by $g \cdot \rho = g\rho g^{-1}$; or equivalently on $G^{\times r}$ by $g \cdot (g_1, \dots, g_r) = (gg_1g^{-1}, \dots, gg_rg^{-1})$.
- 6 So this determines a co-action on the ring of polynomials $\mathbb{C}[\mathrm{Hom}(F_r, G)]$ by $g \cdot f(\rho) = f(g^{-1}\rho g)$.
- 7 Since G is reductive, the ring of invariants $\mathbb{C}[\mathrm{Hom}(F_r, G)]^G := \{f \in \mathbb{C}[\mathrm{Hom}(F_r, G)] \mid g \cdot f = f\}$ is finitely generated. (we will prove this in our special case)
- 8 Therefore, $\mathbb{C}[\mathrm{Hom}(F_r, G)]^G \cong \mathbb{C}[t_1, \dots, t_N]/\mathfrak{I}$ for some ideal \mathfrak{I} .

- 9 Hilbert's Basis Theorem says that \mathfrak{J} is finitely generated.

- 9 Hilbert's Basis Theorem says that \mathfrak{I} is finitely generated. (we will describe $\{t_1, \dots, t_N\}$ and \mathfrak{I} explicitly in our special case)

- 9 Hilbert's Basis Theorem says that \mathfrak{J} is finitely generated. (we will describe $\{t_1, \dots, t_N\}$ and \mathfrak{J} explicitly in our special case)
- 10 Hilbert's Nullstellensatz says that the maximal ideals in $\mathbb{C}[\text{Hom}(F_r, G)]^G$ are exactly of the form $\langle t_1 - a_1, \dots, t_N - a_N \rangle + \mathfrak{J}$ where $(a_1, \dots, a_N) \in \mathbb{C}^N$ is in the simultaneous zero set of the generators of \mathfrak{J} .

- 9 Hilbert's Basis Theorem says that \mathfrak{J} is finitely generated. (we will describe $\{t_1, \dots, t_N\}$ and \mathfrak{J} explicitly in our special case)
- 10 Hilbert's Nullstellensatz says that the maximal ideals in $\mathbb{C}[\text{Hom}(F_r, G)]^G$ are exactly of the form $\langle t_1 - a_1, \dots, t_N - a_N \rangle + \mathfrak{J}$ where $(a_1, \dots, a_N) \in \mathbb{C}^N$ is in the simultaneous zero set of the generators of \mathfrak{J} .
- 11 This correspondence, $\langle t_1 - a_1, \dots, t_N - a_N \rangle + \mathfrak{J} \mapsto (a_1, \dots, a_N)$, determines an algebraic set.

- 9 Hilbert's Basis Theorem says that \mathfrak{J} is finitely generated. (we will describe $\{t_1, \dots, t_N\}$ and \mathfrak{J} explicitly in our special case)
- 10 Hilbert's Nullstellensatz says that the maximal ideals in $\mathbb{C}[\text{Hom}(F_r, G)]^G$ are exactly of the form $\langle t_1 - a_1, \dots, t_N - a_N \rangle + \mathfrak{J}$ where $(a_1, \dots, a_N) \in \mathbb{C}^N$ is in the simultaneous zero set of the generators of \mathfrak{J} .
- 11 This correspondence, $\langle t_1 - a_1, \dots, t_N - a_N \rangle + \mathfrak{J} \mapsto (a_1, \dots, a_N)$, determines an algebraic set.
- 12 This space is the character variety

$$\mathfrak{X}_r(\text{SL}(2, \mathbb{C})).$$

Generators and the Non-Commutative Picture

- Recall that $\mathbb{C}[\mathrm{Hom}(F_r, \mathrm{SL}(2, \mathbb{C}))] = \mathbb{C}[x_{ij}^k] / \Delta$ where Δ is the ideal generated by the r irreducible polynomials

$$x_{11}^k x_{22}^k - x_{12}^k x_{21}^k - 1.$$

Generators and the Non-Commutative Picture

- Recall that $\mathbb{C}[\mathrm{Hom}(F_r, \mathrm{SL}(2, \mathbb{C}))] = \mathbb{C}[x_{ij}^k]/\Delta$ where Δ is the ideal generated by the r irreducible polynomials

$$x_{11}^k x_{22}^k - x_{12}^k x_{21}^k - 1.$$

- Let \bar{x}_{ij}^k be the image of x_{ij}^k under $\mathbb{C}[x_{ij}^k] \rightarrow \mathbb{C}[x_{ij}^k]/\Delta$.

Generators and the Non-Commutative Picture

- Recall that $\mathbb{C}[\mathrm{Hom}(F_r, \mathrm{SL}(2, \mathbb{C}))] = \mathbb{C}[x_{ij}^k] / \Delta$ where Δ is the ideal generated by the r irreducible polynomials

$$x_{11}^k x_{22}^k - x_{12}^k x_{21}^k - 1.$$

- Let \bar{x}_{ij}^k be the image of x_{ij}^k under $\mathbb{C}[x_{ij}^k] \rightarrow \mathbb{C}[x_{ij}^k] / \Delta$.
- Define

$$\mathbf{X}_k = \begin{pmatrix} \bar{x}_{11}^k & \bar{x}_{12}^k \\ \bar{x}_{21}^k & \bar{x}_{22}^k \end{pmatrix}.$$

Generators and the Non-Commutative Picture

- Recall that $\mathbb{C}[\text{Hom}(F_r, \text{SL}(2, \mathbb{C}))] = \mathbb{C}[x_{ij}^k]/\Delta$ where Δ is the ideal generated by the r irreducible polynomials

$$x_{11}^k x_{22}^k - x_{12}^k x_{21}^k - 1.$$

- Let \bar{x}_{ij}^k be the image of x_{ij}^k under $\mathbb{C}[x_{ij}^k] \rightarrow \mathbb{C}[x_{ij}^k]/\Delta$.
- Define

$$\mathbf{X}_k = \begin{pmatrix} \bar{x}_{11}^k & \bar{x}_{12}^k \\ \bar{x}_{21}^k & \bar{x}_{22}^k \end{pmatrix}.$$

They are called *generic unimodular matrices*.

Generators and the Non-Commutative Picture

- Recall that $\mathbb{C}[\text{Hom}(F_r, \text{SL}(2, \mathbb{C}))] = \mathbb{C}[x_{ij}^k] / \Delta$ where Δ is the ideal generated by the r irreducible polynomials

$$x_{11}^k x_{22}^k - x_{12}^k x_{21}^k - 1.$$

- Let \bar{x}_{ij}^k be the image of x_{ij}^k under $\mathbb{C}[x_{ij}^k] \rightarrow \mathbb{C}[x_{ij}^k] / \Delta$.
- Define

$$\mathbf{X}_k = \begin{pmatrix} \bar{x}_{11}^k & \bar{x}_{12}^k \\ \bar{x}_{21}^k & \bar{x}_{22}^k \end{pmatrix}.$$

They are called *generic unimodular matrices*.

- We note that

$$\begin{pmatrix} x_{11}^k & x_{12}^k \\ x_{21}^k & x_{22}^k \end{pmatrix}$$

are simply called *generic matrices*.

- Let's shorten $\mathfrak{X}_{F_r}(\mathrm{SL}(2, \mathbb{C}))$ to simply \mathfrak{X}_r .

- Let's shorten $\mathfrak{X}_{F_r}(\mathrm{SL}(2, \mathbb{C}))$ to simply \mathfrak{X}_r .
- Closely related to $\mathbb{C}[\mathfrak{X}_r] := \mathbb{C}[\mathrm{Hom}(F_r, \mathrm{SL}(2, \mathbb{C}))]^{\mathrm{SL}(2, \mathbb{C})}$ is the ring of invariants

$$\mathbb{C}[\mathfrak{Y}_r] := \mathbb{C}[\mathfrak{gl}(2, \mathbb{C})^{\times r}]^{\mathrm{SL}(2, \mathbb{C})} = \mathbb{C}[x_{ij}^k]^{\mathrm{SL}(2, \mathbb{C})}.$$

- Let's shorten $\mathfrak{X}_{F_r}(\mathrm{SL}(2, \mathbb{C}))$ to simply \mathfrak{X}_r .
- Closely related to $\mathbb{C}[\mathfrak{X}_r] := \mathbb{C}[\mathrm{Hom}(F_r, \mathrm{SL}(2, \mathbb{C}))]^{\mathrm{SL}(2, \mathbb{C})}$ is the ring of invariants

$$\mathbb{C}[\mathfrak{Y}_r] := \mathbb{C}[\mathfrak{gl}(2, \mathbb{C})^{\times r}]^{\mathrm{SL}(2, \mathbb{C})} = \mathbb{C}[x_{ij}^k]^{\mathrm{SL}(2, \mathbb{C})}.$$

- In fact $\mathbb{C}[\mathfrak{Y}_r]/\Delta \approx \mathbb{C}[\mathfrak{X}_r]$.

- Let's shorten $\mathfrak{X}_{F_r}(\mathrm{SL}(2, \mathbb{C}))$ to simply \mathfrak{X}_r .
- Closely related to $\mathbb{C}[\mathfrak{X}_r] := \mathbb{C}[\mathrm{Hom}(F_r, \mathrm{SL}(2, \mathbb{C}))]^{\mathrm{SL}(2, \mathbb{C})}$ is the ring of invariants

$$\mathbb{C}[\mathfrak{Y}_r] := \mathbb{C}[\mathfrak{gl}(2, \mathbb{C})^{\times r}]^{\mathrm{SL}(2, \mathbb{C})} = \mathbb{C}[x_{ij}^k]^{\mathrm{SL}(2, \mathbb{C})}.$$

- In fact $\mathbb{C}[\mathfrak{Y}_r]/\Delta \approx \mathbb{C}[\mathfrak{X}_r]$.
- Otherwise stated,

$$\mathbb{C}[x_{ij}^k]^{\mathrm{SL}(2, \mathbb{C})}/\Delta \approx \left(\mathbb{C}[x_{ij}^k]/\Delta \right)^{\mathrm{SL}(2, \mathbb{C})};$$

which is true because $\mathrm{SL}(2, \mathbb{C})$ is *linearly* reductive and the generators of Δ are invariants.

First Fundamental Theorem of Matrix Invariants

In 1976 Procesi proved

Theorem (Procesi)

$\mathbb{C}[\mathfrak{M}_r]$ is generated by the invariants $\text{tr}(\mathbf{X}_{i_1} \mathbf{X}_{i_2} \cdots \mathbf{X}_{i_k})$, where \mathbf{X}_j are generic matrices.

First Fundamental Theorem of Matrix Invariants

In 1976 Procesi proved

Theorem (Procesi)

$\mathbb{C}[\mathfrak{M}_r]$ is generated by the invariants $\text{tr}(\mathbf{X}_{i_1} \mathbf{X}_{i_2} \cdots \mathbf{X}_{i_k})$, where \mathbf{X}_j are generic matrices.

- As an immediate corollary, we have that $\mathbb{C}[\mathfrak{X}_r]$ is generated by the invariants $\text{tr}(\mathbf{X}_{i_1} \mathbf{X}_{i_2} \cdots \mathbf{X}_{i_k})$, where \mathbf{X}_j are generic *unimodular* matrices.

First Fundamental Theorem of Matrix Invariants

In 1976 Procesi proved

Theorem (Procesi)

$\mathbb{C}[\mathfrak{M}_r]$ is generated by the invariants $\text{tr}(\mathbf{X}_{i_1} \mathbf{X}_{i_2} \cdots \mathbf{X}_{i_k})$, where \mathbf{X}_j are generic matrices.

- As an immediate corollary, we have that $\mathbb{C}[\mathfrak{X}_r]$ is generated by the invariants $\text{tr}(\mathbf{X}_{i_1} \mathbf{X}_{i_2} \cdots \mathbf{X}_{i_k})$, where \mathbf{X}_j are generic *unimodular* matrices.
- **Executive Comment:** As the determinant measures volume, one should think of the trace as measuring length. So Procesi's coordinates are “length coordinates.”

Introduction to “word play”

- The Cayley-Hamilton equation gives

$$\mathbf{X}^2 - \operatorname{tr}(\mathbf{X})\mathbf{X} + \det(\mathbf{X})\mathbf{I} = \mathbf{0}.$$

And if we assume $\det(\mathbf{X}) = 1$, as is the case in $\mathbb{C}[\mathfrak{X}_r]$, we easily derive $\operatorname{tr}(\mathbf{X}^{-1}) = \operatorname{tr}(\mathbf{X})$ and $\operatorname{tr}(\mathbf{X}^2) = \operatorname{tr}(\mathbf{X})^2 - 2$.

Introduction to “word play”

- The Cayley-Hamilton equation gives

$$\mathbf{X}^2 - \operatorname{tr}(\mathbf{X})\mathbf{X} + \det(\mathbf{X})\mathbf{I} = \mathbf{0}.$$

And if we assume $\det(\mathbf{X}) = 1$, as is the case in $\mathbb{C}[\mathfrak{X}_r]$, we easily derive $\operatorname{tr}(\mathbf{X}^{-1}) = \operatorname{tr}(\mathbf{X})$ and $\operatorname{tr}(\mathbf{X}^2) = \operatorname{tr}(\mathbf{X})^2 - 2$.

- Hence the generators $\operatorname{tr}(\mathbf{X}^2)$ in $\mathbb{C}[\mathfrak{Y}_r]$ project to $\operatorname{tr}(\mathbf{X})^2 - 2$ in $\mathbb{C}[\mathfrak{X}_r]$ and so are freely eliminated.

Introduction to “word play”

- The Cayley-Hamilton equation gives

$$\mathbf{X}^2 - \operatorname{tr}(\mathbf{X})\mathbf{X} + \det(\mathbf{X})\mathbf{I} = \mathbf{0}.$$

And if we assume $\det(\mathbf{X}) = 1$, as is the case in $\mathbb{C}[\mathfrak{X}_r]$, we easily derive $\operatorname{tr}(\mathbf{X}^{-1}) = \operatorname{tr}(\mathbf{X})$ and $\operatorname{tr}(\mathbf{X}^2) = \operatorname{tr}(\mathbf{X})^2 - 2$.

- Hence the generators $\operatorname{tr}(\mathbf{X}^2)$ in $\mathbb{C}[\mathfrak{Y}_r]$ project to $\operatorname{tr}(\mathbf{X})^2 - 2$ in $\mathbb{C}[\mathfrak{X}_r]$ and so are freely eliminated.

Example: $r = 1$

- We also get from the characteristic equation (multiplying by \mathbf{X}^{n-2}): $\mathbf{X}^n - \text{tr}(\mathbf{X})\mathbf{X}^{n-1} + \mathbf{X}^{n-2} = \mathbf{0}$, which in turn gives $\text{tr}(\mathbf{X}^n) = \text{tr}(\mathbf{X})\text{tr}(\mathbf{X}^{n-1}) - \text{tr}(\mathbf{X}^{n-2})$.

Example: $r = 1$

- We also get from the characteristic equation (multiplying by \mathbf{X}^{n-2}): $\mathbf{X}^n - \text{tr}(\mathbf{X})\mathbf{X}^{n-1} + \mathbf{X}^{n-2} = \mathbf{0}$, which in turn gives $\text{tr}(\mathbf{X}^n) = \text{tr}(\mathbf{X})\text{tr}(\mathbf{X}^{n-1}) - \text{tr}(\mathbf{X}^{n-2})$.
- Consequently, there are exactly r generators of type $\text{tr}(\mathbf{X})$ in $\mathbb{C}[\mathcal{X}_r]$ and none of type $\text{tr}(\mathbf{X}^n)$, $n \neq 1$; and this is minimal among these generators.

Example: $r = 1$

- We also get from the characteristic equation (multiplying by \mathbf{X}^{n-2}): $\mathbf{X}^n - \text{tr}(\mathbf{X})\mathbf{X}^{n-1} + \mathbf{X}^{n-2} = \mathbf{0}$, which in turn gives $\text{tr}(\mathbf{X}^n) = \text{tr}(\mathbf{X})\text{tr}(\mathbf{X}^{n-1}) - \text{tr}(\mathbf{X}^{n-2})$.
- Consequently, there are exactly r generators of type $\text{tr}(\mathbf{X})$ in $\mathbb{C}[\mathcal{X}_r]$ and none of type $\text{tr}(\mathbf{X}^n)$, $n \neq 1$; and this is minimal among these generators.
- Precisely, the words are $\{\mathbf{X}_1, \dots, \mathbf{X}_r\}$.

Example: $r = 1$

- We also get from the characteristic equation (multiplying by \mathbf{X}^{n-2}): $\mathbf{X}^n - \text{tr}(\mathbf{X})\mathbf{X}^{n-1} + \mathbf{X}^{n-2} = \mathbf{0}$, which in turn gives $\text{tr}(\mathbf{X}^n) = \text{tr}(\mathbf{X})\text{tr}(\mathbf{X}^{n-1}) - \text{tr}(\mathbf{X}^{n-2})$.
- Consequently, there are exactly r generators of type $\text{tr}(\mathbf{X})$ in $\mathbb{C}[\mathfrak{X}_r]$ and none of type $\text{tr}(\mathbf{X}^n)$, $n \neq 1$; and this is minimal among these generators.
- Precisely, the words are $\{\mathbf{X}_1, \dots, \mathbf{X}_r\}$.
- Since the dimension of \mathfrak{X}_1 is 1, we have proved: $\mathbb{C}[\mathfrak{X}_1] \cong \mathbb{C}[t]$ where t corresponds to the invariant function $\text{tr}(\mathbf{X})$.

Example: $r = 2$

- More generally, note that the dimension of \mathfrak{X}_r is equal to $3r - 3$ for $r \geq 2$.
- Multiplying the Cayley-Hamilton equation on both sides by words \mathbf{U} and \mathbf{V} allows us to freely eliminate the generators of type: $\text{tr}(\mathbf{U}\mathbf{X}^n\mathbf{V})$ as long as $n \geq 2$ and at least one of \mathbf{U} or \mathbf{V} is not the identity.

Example: $r = 2$

- More generally, note that the dimension of \mathfrak{X}_r is equal to $3r - 3$ for $r \geq 2$.
- Multiplying the Cayley-Hamilton equation on both sides by words \mathbf{U} and \mathbf{V} allows us to freely eliminate the generators of type: $\text{tr}(\mathbf{U}\mathbf{X}^n\mathbf{V})$ as long as $n \geq 2$ and at least one of \mathbf{U} or \mathbf{V} is not the identity.
- So for the case, $\mathbb{C}[\mathfrak{X}_2]$ we are left with the generators $\text{tr}(\mathbf{X}_1), \text{tr}(\mathbf{X}_2), \text{tr}(\mathbf{X}_1\mathbf{X}_2)$ since any other expression in two letters would result in a sub-expression with an exponent greater than one, which we just showed was impossible.

- Since in this case the dimension of the variety is 3, there can be no further relations and thus these generators are minimal.

- Since in this case the dimension of the variety is 3, there can be no further relations and thus these generators are minimal.
- Consequently, there are exactly $\binom{r}{2}$ generators of type $\text{tr}(\mathbf{XY})$ in $\mathbb{C}[\mathfrak{X}_r]$.

- Since in this case the dimension of the variety is 3, there can be no further relations and thus these generators are minimal.
- Consequently, there are exactly $\binom{r}{2}$ generators of type $\text{tr}(\mathbf{XY})$ in $\mathbb{C}[\mathfrak{X}_r]$.
- This also gives a direct (and short) proof of the Fricke-Vogt theorem: $\mathfrak{X}_2 \cong \mathbb{C}^3$

- Since in this case the dimension of the variety is 3, there can be no further relations and thus these generators are minimal.
- Consequently, there are exactly $\binom{r}{2}$ generators of type $\text{tr}(\mathbf{XY})$ in $\mathbb{C}[\mathfrak{X}_r]$.
- This also gives a direct (and short) proof of the Fricke-Vogt theorem: $\mathfrak{X}_2 \cong \mathbb{C}^3$ (equivalently $\mathbb{C}[\mathfrak{X}_2] \cong \mathbb{C}[x, y, z]$).

A general remark

The only connected rank 1 complex Lie groups are $SL(2, \mathbb{C})$ and $PSL(2, \mathbb{C})$. They are related by the quotient map $SL(2, \mathbb{C}) \rightarrow PSL(2, \mathbb{C})$ with fibre $\{\pm I\}$. Since the fibre is central (i.e. commutes with the conjugation action) there is a natural map $\mathfrak{X}_\Gamma(SL(2, \mathbb{C})) \rightarrow \mathfrak{X}_\Gamma(PSL(2, \mathbb{C}))$ with fibre $\mathfrak{X}_\Gamma(\mathbb{Z}/2\mathbb{Z})$. Using this map, one can determine the relations and generators of $\mathfrak{X}_\Gamma(PSL(2, \mathbb{C}))$ from thos of $\mathfrak{X}_\Gamma(SL(2, \mathbb{C}))$.
See *G-Character varieties for $G = SO(n, \mathbb{C})$ and other not simply connected groups* by Adam S. Sikora.

First step to fundamental relation: Polarization

- Replacing \mathbf{X} with $\mathbf{X} + \mathbf{Y}$ in the Cayley-Hamilton equation gives

$$(\mathbf{X} + \mathbf{Y})^2 - \text{tr}(\mathbf{X} + \mathbf{Y})(\mathbf{X} + \mathbf{Y}) + \det(\mathbf{X} + \mathbf{Y})\mathbf{I} = \mathbf{0}.$$

First step to fundamental relation: Polarization

- Replacing \mathbf{X} with $\mathbf{X} + \mathbf{Y}$ in the Cayley-Hamilton equation gives

$$(\mathbf{X} + \mathbf{Y})^2 - \text{tr}(\mathbf{X} + \mathbf{Y})(\mathbf{X} + \mathbf{Y}) + \det(\mathbf{X} + \mathbf{Y})\mathbf{I} = \mathbf{0}.$$

- Simplifying this expression yields

$$\mathbf{XY} + \mathbf{YX} = \text{tr}(\mathbf{X})\mathbf{Y} + \text{tr}(\mathbf{Y})\mathbf{X} - \text{tr}(\mathbf{X})\text{tr}(\mathbf{Y})\mathbf{I} + \text{tr}(\mathbf{XY})\mathbf{I}.$$

Second step, but an important step...

Multiplying on the right by \mathbf{Z} we get the expression

$$\begin{aligned}\mathrm{tr}(\mathbf{XYZ}) + \mathrm{tr}(\mathbf{YXZ}) &= \mathrm{tr}(\mathbf{X})\mathrm{tr}(\mathbf{YZ}) + \mathrm{tr}(\mathbf{Y})\mathrm{tr}(\mathbf{XZ}) \\ &\quad - \mathrm{tr}(\mathbf{X})\mathrm{tr}(\mathbf{Y})\mathrm{tr}(\mathbf{Z}) + \mathrm{tr}(\mathbf{XY})\mathrm{tr}(\mathbf{Z}).\end{aligned}$$

Second step, but an important step...

Multiplying on the right by \mathbf{Z} we get the expression

$$\begin{aligned}\mathrm{tr}(\mathbf{XYZ}) + \mathrm{tr}(\mathbf{YXZ}) = & \mathrm{tr}(\mathbf{X})\mathrm{tr}(\mathbf{YZ}) + \mathrm{tr}(\mathbf{Y})\mathrm{tr}(\mathbf{XZ}) \\ & - \mathrm{tr}(\mathbf{X})\mathrm{tr}(\mathbf{Y})\mathrm{tr}(\mathbf{Z}) + \mathrm{tr}(\mathbf{XY})\mathrm{tr}(\mathbf{Z}).\end{aligned}$$

At this point, we see that we only need $\binom{r}{3}$ generators of the form $\mathrm{tr}(\mathbf{XYZ})$, and no others of length 3 or more in three letters

Second step, but an important step...

Multiplying on the right by \mathbf{Z} we get the expression

$$\begin{aligned}\mathrm{tr}(\mathbf{XYZ}) + \mathrm{tr}(\mathbf{YXZ}) = & \mathrm{tr}(\mathbf{X})\mathrm{tr}(\mathbf{YZ}) + \mathrm{tr}(\mathbf{Y})\mathrm{tr}(\mathbf{XZ}) \\ & - \mathrm{tr}(\mathbf{X})\mathrm{tr}(\mathbf{Y})\mathrm{tr}(\mathbf{Z}) + \mathrm{tr}(\mathbf{XY})\mathrm{tr}(\mathbf{Z}).\end{aligned}$$

At this point, we see that we only need $\binom{r}{3}$ generators of the form $\mathrm{tr}(\mathbf{XYZ})$, and no others of length 3 or more in three letters. Remember we already have shown we never need exponents beyond 1 in any letter.

Last step...

- Now taking this relation and substituting $\mathbf{Z} \mapsto \mathbf{ZW}$ gives a relation for $\text{tr}(\mathbf{XYZW}) + \text{tr}(\mathbf{YXZW})$.

Last step...

- Now taking this relation and substituting $\mathbf{Z} \mapsto \mathbf{ZW}$ gives a relation for $\text{tr}(\mathbf{XYZW}) + \text{tr}(\mathbf{YXZW})$.
- And substituting $\mathbf{Y} \mapsto \mathbf{WY}$ gives $\text{tr}(\mathbf{XWYZ}) + \text{tr}(\mathbf{WYXZ})$.

Last step...

- Now taking this relation and substituting $\mathbf{Z} \mapsto \mathbf{ZW}$ gives a relation for $\text{tr}(\mathbf{XYZW}) + \text{tr}(\mathbf{YXZW})$.
- And substituting $\mathbf{Y} \mapsto \mathbf{WY}$ gives $\text{tr}(\mathbf{XWYZ}) + \text{tr}(\mathbf{WYXZ})$.
- Sending $\mathbf{X} \mapsto \mathbf{XW}$ gives $\text{tr}(\mathbf{XWYZ}) + \text{tr}(\mathbf{YXWZ})$; and $\mathbf{Z} \mapsto \mathbf{WZ}$ gives $\text{tr}(\mathbf{XYWZ}) + \text{tr}(\mathbf{YXWZ})$.

Last step...

- Now taking this relation and substituting $\mathbf{Z} \mapsto \mathbf{ZW}$ gives a relation for $\text{tr}(\mathbf{XYZW}) + \text{tr}(\mathbf{YXZW})$.
- And substituting $\mathbf{Y} \mapsto \mathbf{WY}$ gives $\text{tr}(\mathbf{XWYZ}) + \text{tr}(\mathbf{WYXZ})$.
- Sending $\mathbf{X} \mapsto \mathbf{XW}$ gives $\text{tr}(\mathbf{XWYZ}) + \text{tr}(\mathbf{YXWZ})$; and $\mathbf{Z} \mapsto \mathbf{WZ}$ gives $\text{tr}(\mathbf{XYWZ}) + \text{tr}(\mathbf{YXWZ})$.
- Subtracting, adding, and subtracting these four relations gives an expression for $\text{tr}(\mathbf{XYZW}) - \text{tr}(\mathbf{XYWZ})$.

Fundamental Relation

- However, sending $\mathbf{X} \mapsto \mathbf{W} \mapsto \mathbf{Y} \mapsto \mathbf{Z} \mapsto \mathbf{X}$ in the first expression gives $\text{tr}(\mathbf{XYZW}) + \text{tr}(\mathbf{XYWZ})$.

Fundamental Relation

- However, sending $\mathbf{X} \mapsto \mathbf{W} \mapsto \mathbf{Y} \mapsto \mathbf{Z} \mapsto \mathbf{X}$ in the first expression gives $\text{tr}(\mathbf{XYZW}) + \text{tr}(\mathbf{XYWZ})$.
- This adds to our sum to give:

$$\begin{aligned}
 2\text{tr}(\mathbf{XYZW}) &= \text{tr}(\mathbf{X})\text{tr}(\mathbf{Y})\text{tr}(\mathbf{Z})\text{tr}(\mathbf{W}) + \text{tr}(\mathbf{X})\text{tr}(\mathbf{YZW}) \\
 &\quad + \text{tr}(\mathbf{Y})\text{tr}(\mathbf{XZW}) + \text{tr}(\mathbf{Z})\text{tr}(\mathbf{XYW}) + \text{tr}(\mathbf{W})\text{tr}(\mathbf{XYZ}) \\
 &\quad - \text{tr}(\mathbf{XZ})\text{tr}(\mathbf{YW}) + \text{tr}(\mathbf{XW})\text{tr}(\mathbf{YZ}) + \text{tr}(\mathbf{XY})\text{tr}(\mathbf{ZW}) \\
 &\quad - \text{tr}(\mathbf{X})\text{tr}(\mathbf{Y})\text{tr}(\mathbf{ZW}) - \text{tr}(\mathbf{X})\text{tr}(\mathbf{W})\text{tr}(\mathbf{YZ}) \\
 &\quad - \text{tr}(\mathbf{Y})\text{tr}(\mathbf{Z})\text{tr}(\mathbf{XW}) - \text{tr}(\mathbf{Z})\text{tr}(\mathbf{W})\text{tr}(\mathbf{XY})
 \end{aligned}$$

Fundamental Relation

- However, sending $\mathbf{X} \mapsto \mathbf{W} \mapsto \mathbf{Y} \mapsto \mathbf{Z} \mapsto \mathbf{X}$ in the first expression gives $\text{tr}(\mathbf{XYZW}) + \text{tr}(\mathbf{XYWZ})$.
- This adds to our sum to give:

$$\begin{aligned}
 2\text{tr}(\mathbf{XYZW}) &= \text{tr}(\mathbf{X})\text{tr}(\mathbf{Y})\text{tr}(\mathbf{Z})\text{tr}(\mathbf{W}) + \text{tr}(\mathbf{X})\text{tr}(\mathbf{YZW}) \\
 &\quad + \text{tr}(\mathbf{Y})\text{tr}(\mathbf{XZW}) + \text{tr}(\mathbf{Z})\text{tr}(\mathbf{XYW}) + \text{tr}(\mathbf{W})\text{tr}(\mathbf{XYZ}) \\
 &\quad - \text{tr}(\mathbf{XZ})\text{tr}(\mathbf{YW}) + \text{tr}(\mathbf{XW})\text{tr}(\mathbf{YZ}) + \text{tr}(\mathbf{XY})\text{tr}(\mathbf{ZW}) \\
 &\quad - \text{tr}(\mathbf{X})\text{tr}(\mathbf{Y})\text{tr}(\mathbf{ZW}) - \text{tr}(\mathbf{X})\text{tr}(\mathbf{W})\text{tr}(\mathbf{YZ}) \\
 &\quad - \text{tr}(\mathbf{Y})\text{tr}(\mathbf{Z})\text{tr}(\mathbf{XW}) - \text{tr}(\mathbf{Z})\text{tr}(\mathbf{W})\text{tr}(\mathbf{XY})
 \end{aligned}$$

- So length 4 words are not need to generate the ring.

- Since \mathbf{W} can be any word in the generic matrices, we have proved that $\mathbb{C}[\mathfrak{X}_r]$ is generated by at most $\binom{r}{1} + \binom{r}{2} + \binom{r}{3} = \frac{r(r^2+5)}{6}$ generators (so the ring is finitely generated!)

- Since \mathbf{W} can be any word in the generic matrices, we have proved that $\mathbb{C}[\mathfrak{X}_r]$ is generated by at most

$\binom{r}{1} + \binom{r}{2} + \binom{r}{3} = \frac{r(r^2+5)}{6}$ generators (so the ring is finitely generated!)

- In particular, here are the generators:

$\mathcal{G}_1 = \{\text{tr}(\mathbf{X}_1), \dots, \text{tr}(\mathbf{X}_r)\}$ of order r .

$\mathcal{G}_2 = \{\text{tr}(\mathbf{X}_i \mathbf{X}_j) \mid 1 \leq i, j \leq r \text{ and } i \neq j\}$ of order $\frac{r(r-1)}{2}$.

$\mathcal{G}_3 = \{\text{tr}(\mathbf{X}_i \mathbf{X}_j \mathbf{X}_k) \mid 1 \leq i < j < k \leq r\}$ of order $\frac{r(r-1)(r-2)}{3}$.

- Since \mathbf{W} can be any word in the generic matrices, we have proved that $\mathbb{C}[\mathfrak{X}_r]$ is generated by at most $\binom{r}{1} + \binom{r}{2} + \binom{r}{3} = \frac{r(r^2+5)}{6}$ generators (so the ring is finitely generated!)
- In particular, here are the generators:
 $\mathcal{G}_1 = \{\text{tr}(\mathbf{X}_1), \dots, \text{tr}(\mathbf{X}_r)\}$ of order r .
 $\mathcal{G}_2 = \{\text{tr}(\mathbf{X}_i \mathbf{X}_j) \mid 1 \leq i, j \leq r \text{ and } i \neq j\}$ of order $\frac{r(r-1)}{2}$.
 $\mathcal{G}_3 = \{\text{tr}(\mathbf{X}_i \mathbf{X}_j \mathbf{X}_k) \mid 1 \leq i < j < k \leq r\}$ of order $\frac{r(r-1)(r-2)}{3}$.
- We will see in a minute that this is a minimal generating set (we can't get rid of any either!)

Geometrically, this says that the minimal (trace) embedding of \mathfrak{X}_r into \mathbb{C}^N is when $N = \frac{r(r^2+5)}{6}$ and the mapping is exactly

$$[\rho] \mapsto \left(\operatorname{tr}(\rho(\gamma_1)), \dots, \operatorname{tr}(\rho(\gamma_r)), \operatorname{tr}(\rho(\gamma_1\gamma_2)), \dots, \operatorname{tr}(\rho(\gamma_{r-1}\gamma_r)), \right. \\ \left. \operatorname{tr}(\rho(\gamma_1\gamma_2\gamma_3)), \dots, \operatorname{tr}(\rho(\gamma_{r-2}\gamma_{r-1}\gamma_r)) \right).$$

Example $r = 3$ and the proof of minimality

- In this case the variety has dimension 6.

Example $r = 3$ and the proof of minimality

- In this case the variety has dimension 6.
- From the $r = 1$ and $r = 2$ cases and our previous theorems, we know there are only two generators not accounted for: $\text{tr}(\mathbf{XYZ})$ and $\text{tr}(\mathbf{YXZ})$.

Example $r = 3$ and the proof of minimality

- In this case the variety has dimension 6.
- From the $r = 1$ and $r = 2$ cases and our previous theorems, we know there are only two generators not accounted for: $\text{tr}(\mathbf{XYZ})$ and $\text{tr}(\mathbf{YXZ})$.
- However, we just saw that only one of these two generator types is necessary.

Example $r = 3$ and the proof of minimality

- In this case the variety has dimension 6.
- From the $r = 1$ and $r = 2$ cases and our previous theorems, we know there are only two generators not accounted for: $\text{tr}(\mathbf{XYZ})$ and $\text{tr}(\mathbf{YXZ})$.
- However, we just saw that only one of these two generator types is necessary.
- There are then only 7 generators.

Example $r = 3$ and the proof of minimality

- In this case the variety has dimension 6.
- From the $r = 1$ and $r = 2$ cases and our previous theorems, we know there are only two generators not accounted for: $\text{tr}(\mathbf{XYZ})$ and $\text{tr}(\mathbf{YXZ})$.
- However, we just saw that only one of these two generator types is necessary.
- There are then only 7 generators.
- If $\text{tr}(\mathbf{XYZ})$ was allowed to be eliminated, we would conclude that \mathfrak{X}_3 was affine \mathbb{C}^6 .

Example $r = 3$ and the proof of minimality

- In this case the variety has dimension 6.
- From the $r = 1$ and $r = 2$ cases and our previous theorems, we know there are only two generators not accounted for: $\text{tr}(\mathbf{XYZ})$ and $\text{tr}(\mathbf{YXZ})$.
- However, we just saw that only one of these two generator types is necessary.
- There are then only 7 generators.
- If $\text{tr}(\mathbf{XYZ})$ was allowed to be eliminated, we would conclude that \mathfrak{X}_3 was affine \mathbb{C}^6 .
- However, it is not hard to show there exists two representations which agree on the six generators of word length two or less but differ at $\text{tr}(\mathbf{XYZ})$.

For instance, $\mathbf{X} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, $\mathbf{Y} = \begin{pmatrix} 0 & 2 \\ -1/2 & 0 \end{pmatrix}$, and $\mathbf{Z} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$ or $\mathbf{Z} = \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix}$ gives two such representations.

For instance, $\mathbf{X} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, $\mathbf{Y} = \begin{pmatrix} 0 & 2 \\ -1/2 & 0 \end{pmatrix}$, and $\mathbf{Z} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$ or $\mathbf{Z} = \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix}$ gives two such representations.

Therefore, generators of the form $\text{tr}(\mathbf{XYZ})$ cannot be written in terms of traces of shorter words. This proves the aforementioned generators form a minimal generating set.

For instance, $\mathbf{X} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, $\mathbf{Y} = \begin{pmatrix} 0 & 2 \\ -1/2 & 0 \end{pmatrix}$, and $\mathbf{Z} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$ or $\mathbf{Z} = \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix}$ gives two such representations.

Therefore, generators of the form $\text{tr}(\mathbf{XYZ})$ cannot be written in terms of traces of shorter words. This proves the aforementioned generators form a minimal generating set.

Remark

For the $n = 2$ case, Sikora showed that the embedding above is in fact minimal among all algebraic embeddings (not just trace embeddings).

For instance, $\mathbf{X} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, $\mathbf{Y} = \begin{pmatrix} 0 & 2 \\ -1/2 & 0 \end{pmatrix}$, and $\mathbf{Z} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$ or $\mathbf{Z} = \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix}$ gives two such representations.

Therefore, generators of the form $\text{tr}(\mathbf{XYZ})$ cannot be written in terms of traces of shorter words. This proves the aforementioned generators form a minimal generating set.

Remark

For the $n = 2$ case, Sikora showed that the embedding above is in fact minimal among all algebraic embeddings (not just trace embeddings).

Remark

One can further show there exists a product relation for $\text{tr}(\mathbf{XYZ})\text{tr}(\mathbf{YXZ})$. Together with the sum relation, we conclude that \mathfrak{X}_3 is a hypersurface.

Deriving the Product Relation

$$\begin{aligned}\mathrm{tr}(\mathbf{XYZ})\mathrm{tr}(\mathbf{XZY}) &= \mathrm{tr}(\mathbf{X})^2 + \mathrm{tr}(\mathbf{Y})^2 + \mathrm{tr}(\mathbf{Z})^2 \\ &\quad + \mathrm{tr}(\mathbf{XY})^2 + \mathrm{tr}(\mathbf{YZ})^2 + \mathrm{tr}(\mathbf{XZ})^2 \\ &\quad - \mathrm{tr}(\mathbf{X})\mathrm{tr}(\mathbf{Y})\mathrm{tr}(\mathbf{XY}) - \mathrm{tr}(\mathbf{Y})\mathrm{tr}(\mathbf{Z})\mathrm{tr}(\mathbf{YZ}) \\ &\quad - \mathrm{tr}(\mathbf{X})\mathrm{tr}(\mathbf{Z})\mathrm{tr}(\mathbf{XZ}) \\ &\quad + \mathrm{tr}(\mathbf{XY})\mathrm{tr}(\mathbf{YZ})\mathrm{tr}(\mathbf{XZ}) - 4\end{aligned}$$

Deriving the Product Relation

$$\begin{aligned}
 \operatorname{tr}(\mathbf{XYZ})\operatorname{tr}(\mathbf{XZY}) &= \operatorname{tr}(\mathbf{X})^2 + \operatorname{tr}(\mathbf{Y})^2 + \operatorname{tr}(\mathbf{Z})^2 \\
 &\quad + \operatorname{tr}(\mathbf{XY})^2 + \operatorname{tr}(\mathbf{YZ})^2 + \operatorname{tr}(\mathbf{XZ})^2 \\
 &\quad - \operatorname{tr}(\mathbf{X})\operatorname{tr}(\mathbf{Y})\operatorname{tr}(\mathbf{XY}) - \operatorname{tr}(\mathbf{Y})\operatorname{tr}(\mathbf{Z})\operatorname{tr}(\mathbf{YZ}) \\
 &\quad - \operatorname{tr}(\mathbf{X})\operatorname{tr}(\mathbf{Z})\operatorname{tr}(\mathbf{XZ}) \\
 &\quad + \operatorname{tr}(\mathbf{XY})\operatorname{tr}(\mathbf{YZ})\operatorname{tr}(\mathbf{XZ}) - 4
 \end{aligned}$$

- Using the characteristic equation, we derive that
 (*) $\operatorname{tr}(\mathbf{AB}) = \operatorname{tr}(\mathbf{A})\operatorname{tr}(\mathbf{B}) - \operatorname{tr}(\mathbf{A}^{-1}\mathbf{B})$.

Deriving the Product Relation

$$\begin{aligned}
 \operatorname{tr}(\mathbf{XYZ})\operatorname{tr}(\mathbf{XZY}) &= \operatorname{tr}(\mathbf{X})^2 + \operatorname{tr}(\mathbf{Y})^2 + \operatorname{tr}(\mathbf{Z})^2 \\
 &\quad + \operatorname{tr}(\mathbf{XY})^2 + \operatorname{tr}(\mathbf{YZ})^2 + \operatorname{tr}(\mathbf{XZ})^2 \\
 &\quad - \operatorname{tr}(\mathbf{X})\operatorname{tr}(\mathbf{Y})\operatorname{tr}(\mathbf{XY}) - \operatorname{tr}(\mathbf{Y})\operatorname{tr}(\mathbf{Z})\operatorname{tr}(\mathbf{YZ}) \\
 &\quad - \operatorname{tr}(\mathbf{X})\operatorname{tr}(\mathbf{Z})\operatorname{tr}(\mathbf{XZ}) \\
 &\quad + \operatorname{tr}(\mathbf{XY})\operatorname{tr}(\mathbf{YZ})\operatorname{tr}(\mathbf{XZ}) - 4
 \end{aligned}$$

- Using the characteristic equation, we derive that
 $(*) \operatorname{tr}(\mathbf{AB}) = \operatorname{tr}(\mathbf{A})\operatorname{tr}(\mathbf{B}) - \operatorname{tr}(\mathbf{A}^{-1}\mathbf{B}).$
- Then $\operatorname{tr}(\mathbf{BCA})\operatorname{tr}(\mathbf{BAC}) - \operatorname{tr}(\mathbf{A}^{-1}\mathbf{C}^{-1}\mathbf{AC}) = \operatorname{tr}([\mathbf{BCA}][\mathbf{BAC}])$

Deriving the Product Relation

$$\begin{aligned}
 \operatorname{tr}(\mathbf{XYZ})\operatorname{tr}(\mathbf{XZY}) &= \operatorname{tr}(\mathbf{X})^2 + \operatorname{tr}(\mathbf{Y})^2 + \operatorname{tr}(\mathbf{Z})^2 \\
 &\quad + \operatorname{tr}(\mathbf{XY})^2 + \operatorname{tr}(\mathbf{YZ})^2 + \operatorname{tr}(\mathbf{XZ})^2 \\
 &\quad - \operatorname{tr}(\mathbf{X})\operatorname{tr}(\mathbf{Y})\operatorname{tr}(\mathbf{XY}) - \operatorname{tr}(\mathbf{Y})\operatorname{tr}(\mathbf{Z})\operatorname{tr}(\mathbf{YZ}) \\
 &\quad - \operatorname{tr}(\mathbf{X})\operatorname{tr}(\mathbf{Z})\operatorname{tr}(\mathbf{XZ}) \\
 &\quad + \operatorname{tr}(\mathbf{XY})\operatorname{tr}(\mathbf{YZ})\operatorname{tr}(\mathbf{XZ}) - 4
 \end{aligned}$$

- Using the characteristic equation, we derive that
 $(*) \operatorname{tr}(\mathbf{AB}) = \operatorname{tr}(\mathbf{A})\operatorname{tr}(\mathbf{B}) - \operatorname{tr}(\mathbf{A}^{-1}\mathbf{B}).$
- Then $\operatorname{tr}(\mathbf{BCA})\operatorname{tr}(\mathbf{BAC}) - \operatorname{tr}(\mathbf{A}^{-1}\mathbf{C}^{-1}\mathbf{AC}) = \operatorname{tr}([\mathbf{BCA}][\mathbf{BAC}])$
- And $\operatorname{tr}(\mathbf{BC})\operatorname{tr}(\mathbf{ABAC}) - \operatorname{tr}(\mathbf{B}^{-1}\mathbf{ABA}) = \operatorname{tr}([\mathbf{BC}][\mathbf{ABAC}])$

Deriving the Product Relation

$$\begin{aligned}
 \operatorname{tr}(\mathbf{XYZ})\operatorname{tr}(\mathbf{XZY}) &= \operatorname{tr}(\mathbf{X})^2 + \operatorname{tr}(\mathbf{Y})^2 + \operatorname{tr}(\mathbf{Z})^2 \\
 &\quad + \operatorname{tr}(\mathbf{XY})^2 + \operatorname{tr}(\mathbf{YZ})^2 + \operatorname{tr}(\mathbf{XZ})^2 \\
 &\quad - \operatorname{tr}(\mathbf{X})\operatorname{tr}(\mathbf{Y})\operatorname{tr}(\mathbf{XY}) - \operatorname{tr}(\mathbf{Y})\operatorname{tr}(\mathbf{Z})\operatorname{tr}(\mathbf{YZ}) \\
 &\quad - \operatorname{tr}(\mathbf{X})\operatorname{tr}(\mathbf{Z})\operatorname{tr}(\mathbf{XZ}) \\
 &\quad + \operatorname{tr}(\mathbf{XY})\operatorname{tr}(\mathbf{YZ})\operatorname{tr}(\mathbf{XZ}) - 4
 \end{aligned}$$

- Using the characteristic equation, we derive that
 $(*) \operatorname{tr}(\mathbf{AB}) = \operatorname{tr}(\mathbf{A})\operatorname{tr}(\mathbf{B}) - \operatorname{tr}(\mathbf{A}^{-1}\mathbf{B}).$
- Then $\operatorname{tr}(\mathbf{BCA})\operatorname{tr}(\mathbf{BAC}) - \operatorname{tr}(\mathbf{A}^{-1}\mathbf{C}^{-1}\mathbf{AC}) = \operatorname{tr}([\mathbf{BCA}][\mathbf{BAC}])$
- And $\operatorname{tr}(\mathbf{BC})\operatorname{tr}(\mathbf{ABAC}) - \operatorname{tr}(\mathbf{B}^{-1}\mathbf{ABA}) = \operatorname{tr}([\mathbf{BC}][\mathbf{ABAC}])$
- Again using $(*)$, we can simplify $\operatorname{tr}(\mathbf{B}^{-1}\mathbf{ABA})$,
 $\operatorname{tr}(\mathbf{ACA}^{-1}\mathbf{C}^{-1})$, and $\operatorname{tr}(\mathbf{ABAC})$.

Deriving the Product Relation

$$\begin{aligned}
 \text{tr}(\mathbf{XYZ})\text{tr}(\mathbf{XZY}) &= \text{tr}(\mathbf{X})^2 + \text{tr}(\mathbf{Y})^2 + \text{tr}(\mathbf{Z})^2 \\
 &\quad + \text{tr}(\mathbf{XY})^2 + \text{tr}(\mathbf{YZ})^2 + \text{tr}(\mathbf{XZ})^2 \\
 &\quad - \text{tr}(\mathbf{X})\text{tr}(\mathbf{Y})\text{tr}(\mathbf{XY}) - \text{tr}(\mathbf{Y})\text{tr}(\mathbf{Z})\text{tr}(\mathbf{YZ}) \\
 &\quad - \text{tr}(\mathbf{X})\text{tr}(\mathbf{Z})\text{tr}(\mathbf{XZ}) \\
 &\quad + \text{tr}(\mathbf{XY})\text{tr}(\mathbf{YZ})\text{tr}(\mathbf{XZ}) - 4
 \end{aligned}$$

- Using the characteristic equation, we derive that
 $(*) \text{tr}(\mathbf{AB}) = \text{tr}(\mathbf{A})\text{tr}(\mathbf{B}) - \text{tr}(\mathbf{A}^{-1}\mathbf{B}).$
- Then $\text{tr}(\mathbf{BCA})\text{tr}(\mathbf{BAC}) - \text{tr}(\mathbf{A}^{-1}\mathbf{C}^{-1}\mathbf{AC}) = \text{tr}([\mathbf{BCA}][\mathbf{BAC}])$
- And $\text{tr}(\mathbf{BC})\text{tr}(\mathbf{ABAC}) - \text{tr}(\mathbf{B}^{-1}\mathbf{ABA}) = \text{tr}([\mathbf{BC}][\mathbf{ABAC}])$
- Again using $(*)$, we can simplify $\text{tr}(\mathbf{B}^{-1}\mathbf{ABA})$,
 $\text{tr}(\mathbf{ACA}^{-1}\mathbf{C}^{-1})$, and $\text{tr}(\mathbf{ABAC})$.
- Together, these formulae give the product relation.

Exercises

- 1 By hand fill in the details for the derivations for the product formula $\text{tr}(\mathbf{ABC})\text{tr}(\mathbf{ACB})$.
- 2 By hand fill in the details for the derivation for the $\text{tr}(\mathbf{XYZW})$.
- 3 Verify that the example representations I gave are the same on $\text{tr}(\mathbf{A})$, $\text{tr}(\mathbf{B})$, $\text{tr}(\mathbf{C})$, $\text{tr}(\mathbf{AB})$, $\text{tr}(\mathbf{AC})$, $\text{tr}(\mathbf{BC})$ but differ on $\text{tr}(\mathbf{ABC})$ and $\text{tr}(\mathbf{ACB})$.
- 4 Write algorithms by hand that turn $\text{tr}(\mathbf{W})$ into a trace expression with every letter of every represented word having exponent 1 (non-negative and no-multiplicity) and no word having length greater than 3.
- 5 Together implement the above algorithm in *Mathematica*.