

MA 523: Homework 4

Carlos Salinas

September 28, 2016

PROBLEM 4.1 (LEGENDRE TRANSFORM)

Let $u(x_1, x_2)$ be a solution of the quasilinear equation

$$a^{11}(Du)u_{x_1x_1} + 2a^{12}(Du)u_{x_1x_2} + a^{22}(Du)u_{x_2x_2} = 0$$

in some region of \mathbb{R}^2 , where we can invert the relations

$$p^1 = u_{x_1}(x_1, x_2), \quad p^2 = u_{x_2}(x_1, x_2)$$

to solve for

$$x^1 = x^1(p_1, p_2), \quad x^2 = x^2(p_1, p_2).$$

Define then

$$v(p) := \mathbf{x}(p) \cdot p - u(\mathbf{x}(p)),$$

where $\mathbf{x} = (x^1, x^2)$, $p = (p_1, p_2)$. Show that v satisfies the *linear* equation

$$a^{22}(p)v_{p_1p_2} - 2a^{12}(p)v_{p_1p_2} + a^{11}(p)v_{p_1p_2} = 0.$$

(*Hint:* See [Evans, 4.4.3b], prove the identities (29)).

SOLUTION.

$$\Phi = \Phi \Phi$$

■

PROBLEM 4.2

Find the solution $u(x, t)$ of the one-dimensional wave equation

$$u_{tt} - u_{xx} = 0$$

in the quadrant $x > 0, t > 0$ for which

$$\begin{cases} u(x, 0) = f(x), & u_t(x, 0) = g(x) & \text{for } x > 0 \\ u_t(0, t) = \alpha u_x(0, t), & & \text{for } t > 0, \end{cases}$$

where $\alpha \neq -1$ is a given constant. Show that generally no solution exists when $\alpha = -1$. (*Hint:* Use a representation $u(x, t) = F(x - t) + G(x + t)$ for the solution.)

SOLUTION.

■

PROBLEM 4.3

- (a) Let u be a solution of the wave equation $u_{tt} - c^2 u_{xx} = 0$ for $0 < x < \pi$, $t > 0$ such that $u(0, t) = u(\pi, t) = 0$. Show that the *energy*

$$E(t) = \frac{1}{2} \int_0^\pi (u_t^2 + c^2 u_x^2) dx, \quad t > 0$$

is independent of t ; i.e., $dE/dt = 0$ for $t > 0$. Assume that u is C^2 up to the boundary.

- (b) Express the energy E of the Fourier series solution

$$u(x, t) = \sum_{n=1}^{\infty} (a_n \cos(nct) + b_n \sin(nct)) \sin nx$$

in terms of coefficients a_n , b_n .

SOLUTION. ■