

# MA571 Problem Set 1

Carlos Salinas

August 28, 2015



**Problem 1.1 (Munkres §2, 1(a,b).)**

Let  $f: A \rightarrow B$ . Let  $A_0 \subset A$  and  $B_0 \subset B$ .

- (a) Show that  $A_0 \subset f^{-1}(f(A_0))$  and that equality holds if  $f$  is injective.
- (b) Show that  $f(f^{-1}(B_0)) \subset B_0$  and that equality holds if  $f$  is surjective.

*Proof.* (a). First, we will show  $A_0 \subset f^{-1}(f(A_0))$ . Let  $x \in A_0$ . Then  $f(x) \in f(A_0)$ . By definition,  $f^{-1}(f(A_0))$  is the set of those points  $x_0 \in A$  such that  $f(x_0) \in f(A_0)$  and in particular we see that the containment  $A_0 \subset f^{-1}(f(A_0))$  holds. Thus,  $x \in f^{-1}(f(A_0))$ .

Now, let us suppose the map  $f$  is injective. By our former argument, we have that  $A_0 \subset f^{-1}(f(A_0))$  therefore, we will show the reverse containment. If  $y \in f(A_0)$ , then  $f(x) = y$  for some  $x \in A_0$ . By the injectivity of  $f$ , if  $f(x_0) = y$  for some  $x_0 \in A$ , then we must have that  $x_0 = x$ . In particular,  $x_0 \in A_0$ . Thus  $f^{-1}(f(A_0)) \subset A_0$  and equality  $A_0 = f^{-1}(f(A_0))$  holds.

(b). First, we will show that  $f(f^{-1}(B_0)) \subset B_0$ . Consider the preimage  $f^{-1}(B_0)$  of  $B_0$ . Let  $x \in f^{-1}(B_0)$ . Then  $f(x) = y$  for some  $y \in B_0$ . Since  $f(f^{-1}(B_0))$  is, by definition, the set of all points  $f(x) \in B$  where  $x \in f^{-1}(B_0)$  and  $f(x) = y$  for  $y \in B_0$ , we have that  $f(f^{-1}(B_0)) \subset B_0$ .

Now, let us suppose the map  $f$  is surjective. Let  $y \in B_0$ , then there exists  $x \in A$  such that  $f(x) = y$ . Thus,  $x \in f^{-1}(B_0)$ . Then  $y = f(x) \in f(f^{-1}(B_0))$  (in particular  $B_0 \subset f(f^{-1}(B_0))$ ) and we have equality  $B_0 = f(f^{-1}(B_0))$ . ■

**Problem 1.2 (Munkres, §2, 2(g).)**

Let  $f: A \rightarrow B$  and let  $A_i \subset A$  and  $B_i \subset B$  for  $i = 0$  and  $i = 1$ . Show that  $f^{-1}$  preserves inclusion, unions, intersections, and differences of sets:

(g)  $f(A_0 \cap A_1) \subset f(A_0) \cap f(A_1)$ ; show that equality holds if  $f$  is injective.

*Proof of (g).* The claim is evident if  $A_0$  and  $A_1$  are disjoint subsets. Suppose  $A_0 \cap A_1 \neq \emptyset$ . Let  $y \in f(A_0 \cap A_1)$ . Then  $y = f(x)$  for some  $x \in A_0$ ,  $x \in A_1$ . Then  $f(x) \in f(A_0)$  and  $f(x) \in f(A_1)$  so  $y \in f(A_0) \cap f(A_1)$ . Thus,  $f(A_0 \cap A_1) \subset f(A_0) \cap f(A_1)$ .

Now, suppose  $f$  is injective. Then, if  $f(x) = f(x') = y$  for some  $y \in B$ , then  $x = x'$ . Let  $y \in f(A_0) \cap f(A_1)$ . Then  $y = f(x_0)$ ,  $y = f(x_1)$  for some  $x_0 \in A_0$ ,  $x_1 \in A_1$ . But, by the injectivity of  $f$ ,  $x_0 = x_1$  so  $x_0 \in A_0 \cap A_1$ . Hence,  $y \in f(A_0 \cap A_1)$  and the equality  $f(A_0 \cap A_1) = f(A_0) \cap f(A_1)$  holds. ■

**Problem 1.3 (Munkres, §13, 3.)**

Show that the collection  $\mathcal{T}_c$  given in Example 4 of §12 is a topology on the set  $X$ . Is the collection

$$\mathcal{T}_\infty = \{U \mid X \setminus U \text{ is infinite or empty or all of } X\}$$

a topology on  $X$ ?

*Proof.* Recall that  $\mathcal{T}_c$  is the collection of all subsets  $U$  of  $X$  such that  $X \setminus U$  is either countable or is all of  $X$ . Let us verify that  $\mathcal{T}_c$  defines a topology on  $X$ . First,  $\emptyset \in \mathcal{T}_c$  since  $X \setminus \emptyset = X$  and  $X \in \mathcal{T}_c$  since  $X \setminus X = \emptyset$  is countable. Second, let  $\{U_\alpha\}$ ,  $\alpha \in A$ , be an indexed family of nonempty elements of  $\mathcal{T}_c$ , then  $X \setminus U_\alpha$  is countable for all  $\alpha$ . Thus, by DeMorgan's laws, we have that

$$X \setminus \bigcup U_\alpha = \bigcap X \setminus U_\alpha$$

is countable (this follows from Corollary 7.3, since  $\bigcap_\alpha X \setminus U_\alpha$  is a subset of  $U_\beta$  for all  $\beta \in A$ , hence it is countable). Thus, the union  $\bigcup U_\alpha$  is in  $\mathcal{T}_c$ . Lastly, let  $U_1, \dots, U_n$  be nonempty elements of  $\mathcal{T}_c$ , then by DeMorgan's laws, we have that

$$X \setminus \bigcap_{i=1}^n U_i = \bigcup_{i=1}^n (X \setminus U_i)$$

is countable by Theorem 7.5 since  $\bigcup_{i=1}^n (X \setminus U_i)$  is a countable union of countable sets. So the finite intersection  $\bigcap_{i=1}^n U_i \in \mathcal{T}_c$ . Therefore,  $\mathcal{T}_c$  satisfies all the properties to define a topology on  $X$ .

Now, let us consider the collection of subsets of  $X$ ,  $\mathcal{T}_\infty$ , given above. We will show that arbitrary unions of elements of  $\mathcal{T}_\infty$  are, in general, not in  $\mathcal{T}_\infty$ . Let  $X = \mathbf{Z}_+$  and suppose that  $\mathcal{T}_\infty$  defines a topology on  $X$ . Consider the collection of subsets  $\{\{i\}\}_{i=1}^\infty$ .  $\mathbf{Z}_+ \setminus \{n\} = \{1, \dots, n-1, n+1, \dots\}$  is infinite hence,  $\{i\} \in \mathcal{T}_\infty$  for all  $i \in \{1, \dots\}$ . However,  $\mathbf{Z}_+ \setminus \bigcup_{i=1}^\infty \{i\} = \{0\}$  is finite so  $\bigcup_{i=1}^\infty \{i\} \notin \mathcal{T}_\infty$ , this is a contradiction. Therefore,  $\mathcal{T}_\infty$  does not define a topology on  $X$ . ■

**Problem 1.4 (Munkres, §13, 5.)**

Show that if  $\mathcal{A}$  is a basis for a topology on  $X$ , then the topology generated by  $\mathcal{A}$  equals the intersection of all topologies on  $X$  that contain  $\mathcal{A}$ . Prove the same if  $\mathcal{A}$  is a subbasis.

*Proof.* Let  $\mathcal{T}$  be the topology generated by  $\mathcal{A}$  and let  $\mathcal{S}$  be the collection of all topologies  $\mathcal{T}'$  that contain  $\mathcal{A}$ . By Lemma 13.3, it suffices to check that  $\mathcal{T} = \bigcap \mathcal{T}'$ . First we will show that the intersection  $\bigcap \mathcal{T}'$  indeed defines a topology on  $X$ . To that end we shall prove the following lemma:

**Lemma 1.** *Let  $X$  be a nonempty set and let  $\{\mathcal{T}_\alpha\}$  be an indexed collection of topologies on  $X$ . Then  $\bigcap \mathcal{T}_\alpha$  defines a topology on  $X$ .*

*Proof of Lemma 1.* Let  $\mathcal{T} = \bigcap \mathcal{T}_\alpha$ . First, since  $\emptyset \in \mathcal{T}_\alpha$  and  $X \in \mathcal{T}_\alpha$  for all  $\alpha \in A$ ,  $\emptyset$  and  $X$  are in  $\mathcal{T}$ . Second, let  $\{U_\beta\}$ ,  $\beta \in B$ , be an indexed family of nonempty elements of  $\mathcal{T}$ . Then,  $U_\beta \in \mathcal{T}_\alpha$  for all  $\beta \in B$  for all  $\alpha \in A$  so  $\bigcup U_\beta \in \mathcal{T}_\alpha$  for all  $\alpha \in A$ . Hence,  $\bigcup U_\beta \in \mathcal{T}$ . Lastly, let  $U_1, \dots, U_n$  be nonempty elements of  $\mathcal{T}$ . Then,  $U_1, \dots, U_n \in \mathcal{T}_\alpha$  for all  $\alpha \in A$  so  $\bigcap_{i=1}^n U_i \in \mathcal{T}_\alpha$  for all  $\alpha \in A$  thus,  $\bigcap_{i=1}^n U_i \in \mathcal{T}$ . We see that, indeed,  $\mathcal{T}$  defines a topology on  $X$ . ♦

By the Lemma 1 above, it follows that  $\bigcap \mathcal{T}'$  gives a topology on  $X$ . Now, it is easy to see that  $\bigcap \mathcal{T}' \subset \mathcal{T}$  since  $\mathcal{T} \in \mathcal{S}$  is the coarsest topology containing  $\mathcal{A}$ . Let us prove this fact:

**Lemma 2.** *Let  $X$  be a nonempty set. Let  $\mathcal{A}$  be a basis for the topology  $\mathcal{T}$  on  $X$ . Then  $\mathcal{T}$  is the coarsest topology containing  $\mathcal{A}$ .*

*Proof of Lemma 2.* This can be easily proven by contradiction for suppose  $\mathcal{T}$  is not the coarsest topology containing  $\mathcal{A}$ . Let  $\mathcal{C}$  be a strictly coarser topology that contains  $\mathcal{A}$ . Then there exists some open set  $U \in \mathcal{T}$  not in  $\mathcal{C}$ . Thus,  $\mathcal{C}$  is not generated by  $\mathcal{A}$ . ♦

On the other hand we see by Lemma 13.1 that  $\mathcal{T} \subset \bigcap \mathcal{T}'$  since each  $\mathcal{T}' \in \mathcal{S}$  contains the basis  $\mathcal{A}$  of  $\mathcal{T}$ , hence contains the open sets of  $\mathcal{T}$ .

T

■

**Problem 1.5 (Munkres, §13, 8(b).)**

(b) Show that the collection

$$\mathcal{C} = \{ [a, b) \mid a < b, a \text{ and } b \text{ rational} \}$$

is a basis that generates a topology different from the lower limit topology on  $\mathbf{R}$ .

*Proof of (b).*

■

**Problem 1.6 (Munkres, §16, 1.)**

Show that if  $Y$  is a subspace of  $X$ , and  $A$  is a subset of  $Y$ , then the topology  $A$  inherits as a subspace of  $Y$  is the same as the topology it inherits as a subspace of  $X$ .

*Proof.*

■



**Problem 1.7 (Munkres, §16, 4.)**

A map  $f: X \rightarrow Y$  is said to be an *open map* if for every open set  $U$  of  $X$ , the set  $f(U)$  is open in  $Y$ . Show that  $\pi_1: X \times Y \rightarrow X$  and  $\pi_2: X \times Y \rightarrow Y$  are open maps.

*Proof.*

■

**Problem 1.8 (Munkres, §16, 6.)**

Show that the countable collection

$$\{ (a, b) \times (c, d) \mid a < b \text{ and } c < d, \text{ and } a, b, c, d \text{ are rational} \}$$

is a basis for  $\mathbf{R}^2$ .

*Proof.*

■

**Problem 1.9 (Munkres, §16, 9.)**

Show that the dictionary order topology on the set  $\mathbf{R} \times \mathbf{R}$  is the same as the product topology  $\mathbf{R}_d \times \mathbf{R}$ , where  $\mathbf{R}_d$  denotes  $\mathbf{R}$  in the discrete topology. Compare this topology with the standard topology on  $\mathbf{R}^2$ .

*Proof.*

■