

Part IB Physics A : Lent 2022

QUANTUM PHYSICS EXAMPLES III

Prof. C. Castelnovo

1. Obtain the following commutation relations for the angular momentum operators $\hat{L} = \hat{r} \times \hat{p}$, and comment on the results:

$$\begin{aligned}\left[\hat{L}_x, \hat{x}\right] &= 0 & \left[\hat{L}_x, \hat{y}\right] &= i\hbar\hat{z} \\ \left[\hat{L}_x, \hat{p}_x\right] &= 0 & \left[\hat{L}_x, \hat{p}_y\right] &= i\hbar\hat{p}_z \\ \left[\hat{L}_x, \hat{L}^2\right] &= \left[\hat{L}_x, \hat{r}^2\right] = \left[\hat{L}_x, \hat{p}^2\right] = 0\end{aligned}$$

(All other commutation relations follow by the cyclic permutations $x \rightarrow y \rightarrow z \rightarrow x$.)

2. Use the commutation relations for the angular momentum operators,

$$\left[\hat{L}_x, \hat{L}_y\right] = i\hbar\hat{L}_z \quad \left[\hat{L}_y, \hat{L}_z\right] = i\hbar\hat{L}_x \quad \left[\hat{L}_z, \hat{L}_x\right] = i\hbar\hat{L}_y,$$

and the definitions of angular momentum raising and lowering operators,

$$\hat{L}_{\pm} = \hat{L}_x \pm i\hat{L}_y = \hbar e^{\pm i\phi} \left(\pm \frac{\partial}{\partial \theta} + i \cot \theta \frac{\partial}{\partial \phi} \right),$$

to show that

$$\hat{L}^2 = \hat{L}_+ \hat{L}_- + \hat{L}_z^2 - \hbar \hat{L}_z$$

and that

$$\left[\hat{L}_+, \hat{L}_-\right] = 2\hbar\hat{L}_z.$$

Hence show that

$$\hat{L}_z = -i\hbar \frac{\partial}{\partial \phi}$$

and that

$$\hat{L}^2 = -\hbar^2 \left(\frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \sin \theta \frac{\partial}{\partial \theta} + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \phi^2} \right).$$

Finally, obtain the angular momentum quantum numbers for an electron in the hydrogen atom for the following eigenfunctions:

$$\psi(r, \theta, \phi) = R_1(r) \quad \psi(r, \theta, \phi) = R_2(r) \sin \theta e^{i\phi} \quad \psi(r, \theta, \phi) = R_3(r)(3 \cos^2 \theta - 1).$$

3. The orthogonal wave functions $\psi_x = xf(r)$, $\psi_y = yf(r)$ and $\psi_z = zf(r)$ represent three of the electronic bound state solutions for a hydrogen atom. Prove the relations shown in the first row of the table below:

$$\begin{array}{lll}
\hat{L}_x \psi_x = 0 & \hat{L}_x \psi_y = i\hbar \psi_z & \hat{L}_x \psi_z = -i\hbar \psi_y \\
\hat{L}_y \psi_x = -i\hbar \psi_z & \hat{L}_y \psi_y = 0 & \hat{L}_y \psi_z = i\hbar \psi_x \\
\hat{L}_z \psi_x = i\hbar \psi_y & \hat{L}_z \psi_y = -i\hbar \psi_x & \hat{L}_z \psi_z = 0
\end{array}$$

Use the results in the table to prove that the expectation value of each component of the angular momentum of any one of ψ_x , ψ_y and ψ_z is zero. Show, however, that each is an eigenfunction of the operator $\hat{L}^2 = \hat{L}_x^2 + \hat{L}_y^2 + \hat{L}_z^2$ and determine the eigenvalue.

Show that the linear combinations $\psi_{\pm} = \psi_x \pm i\psi_y$ are eigenfunctions of \hat{L}_z and determine their orbital angular momentum quantum numbers m and ℓ .

For Questions 4 and 5 you can use the following information about a hydrogen-like atom

The normalised wavefunctions $Y_{\ell m_{\ell}}(\theta, \phi)$ for $\ell = 0, 1$ and 2 are:

$$\begin{aligned}
Y_{00} &= \sqrt{\frac{1}{4\pi}} \\
Y_{10} &= \sqrt{\frac{3}{4\pi}} \cos \theta & Y_{1\pm 1} &= \mp \sqrt{\frac{3}{8\pi}} \sin \theta e^{\pm i\phi} \\
Y_{20} &= \sqrt{\frac{5}{16\pi}} (3 \cos^2 \theta - 1) & Y_{2\pm 1} &= \mp \sqrt{\frac{15}{8\pi}} \sin \theta \cos \theta e^{\pm i\phi} & Y_{2\pm 2} &= \sqrt{\frac{15}{32\pi}} \sin^2 \theta e^{\pm 2i\phi}
\end{aligned}$$

and the normalised hydrogen-like radial wavefunctions $R_{n\ell}$ for $n = 1, 2$ are:

$$\begin{aligned}
R_{10} &= (Z/a_0)^{3/2} 2 \exp(-Zr/a_0) \\
R_{20} &= (Z/2a_0)^{3/2} (2 - Zr/a_0) \exp(-Zr/2a_0) \\
R_{21} &= (Z/2a_0)^{3/2} (1/\sqrt{3}) (Zr/a_0) \exp(-Zr/2a_0)
\end{aligned}$$

where a_0 is the Bohr radius and Z is the atomic number.

Note also that $\int_0^{\infty} x^n e^{-x} dx = n!$.

4. Confirm, for the cases $\ell = 1$ and $\ell = 2$, that

$$\sum_{m_{\ell}=-\ell}^{m_{\ell}=\ell} |Y_{\ell m_{\ell}}|^2 = \text{constant}.$$

Discuss the significance of this result for the electron probability distributions in the hydrogen atom. (The theorem for general ℓ follows from an addition formula for Legendre polynomials, see Whittaker and Watson, p. 327.)

5. An electron is in the ground state of a hydrogen-like atom with nuclear charge $+Ze$.

- What is its average distance from the nucleus?
- At what distance from the nucleus is it most likely to be found?
- Show that the expectation value of the potential energy operator of the electron is $-Z^2 e^2 / 4\pi\epsilon_0 a_0$.
- Show that the expectation value of the kinetic energy operator is $Z^2 e^2 / 8\pi\epsilon_0 a_0$.
- Hence verify that the expectation value of the Hamiltonian is the energy of the ground state.

6. The potential energy for a three-dimensional harmonic oscillator of mass m and frequency ω is $V(x, y, z) = \frac{1}{2}m\omega^2(x^2 + y^2 + z^2)$.

What are the energies and degeneracies of the three lowest levels?

Show that the degeneracy of the n^{th} excited level is $\frac{1}{2}(n+1)(n+2)$.

7. In a one-dimensional system two particles each of mass m interact through the potential $\frac{1}{2}m\omega^2(x_1 - x_2)^2$, where x_1 and x_2 are their position coordinates. Find the energy levels of the system when its centre of mass is at rest.

8. The Hamiltonian \hat{H} of two interacting particles a and b is given by

$$\hat{H} = \frac{\hat{p}_a^2}{2m_a} + \frac{\hat{p}_b^2}{2m_b} + \hat{V}(|\mathbf{r}|)$$

where $\mathbf{r} = \mathbf{r}_a - \mathbf{r}_b$ is the relative position of the particles.

Derive the commutation relations of the centre-of-mass and relative position and momentum operators \hat{R} , \hat{r} , \hat{P} and \hat{p} , where:

$$\hat{\mathbf{R}} = \frac{m_a \hat{\mathbf{r}}_a + m_b \hat{\mathbf{r}}_b}{m_a + m_b} \quad \hat{\mathbf{p}} = \frac{m_a m_b}{m_a + m_b} \left(\frac{\hat{\mathbf{p}}_a}{m_a} - \frac{\hat{\mathbf{p}}_b}{m_b} \right)$$

Comment on your results.

ANSWERS:

2. $\ell = 0, m_\ell = 0$; $\ell = 1, m_\ell = 1$; $\ell = 2, m_\ell = 0$.

3. Eigenvalue of \hat{L}^2 is $2\hbar^2$; $\ell = 1, m_\ell = \pm 1$.

4. $\ell = 1$: $3/(4\pi)$; $\ell = 2$: $5/(4\pi)$.

5. (a) $3a_0/2Z$; (b) a_0/Z .

6. $\frac{3}{2}\hbar\omega$, $\frac{5}{2}\hbar\omega$, $\frac{7}{2}\hbar\omega$; 1, 3, 6.

7. $E_n = \sqrt{2}(n + \frac{1}{2})\hbar\omega$.

8. $[\hat{R}_j, \hat{P}_k] = [\hat{r}_j, \hat{p}_k] = i\hbar\delta_{jk}$, where $j, k = x, y, z$. All other commutators are zero.