

The geometry of the Universe

Relativistic Astrophysics and Cosmology: Lecture 16

Sandro Tacchella

Friday 10th November 2023

Pre-lecture question:

Can you measure how fast you are moving?

Last time

- ▶ Gravitational lensing

This lecture

- ▶ What is cosmology
- ▶ The geometry of the universe
- ▶ The Friedmann-Robertson-Walker metric

Next lecture


- ▶ The dynamics of the universe

Cosmology


- ▶ The remainder of the course covers cosmology
 - ▶ Week 6: The relativistic Universe
 - ▶ Lecture 16: The geometry of the Universe
 - ▶ Lecture 17: The dynamics of the Universe
 - ▶ Lecture 18: The evolution of the Universe
 - ▶ Week 7: Observational cosmology
 - ▶ Lecture 19: Measuring the Universe
 - ▶ Lecture 20: The constituents of the Universe
 - ▶ Lecture 21: Cosmological data and the Cosmic Microwave Background
 - ▶ Week 8: Beyond homogeneity and the quantum Universe
 - ▶ Lecture 22: The primordial universe
 - ▶ Lecture 23: The perturbed universe
 - ▶ Lecture 24: The quantum universe & wrap-up
- ▶ Aim here is to consolidate understanding of Part II material, and fill in many of the details.
- ▶ In particular, aim to give a physically- rather than mathematically-based approach, to complement what you did last year.
- ▶ Then in later part of course we can cover new material, such as observations, inflation & perturbation theory.

What is cosmology?



- ▶ Cosmology is the study of the universe as a whole, and on the largest scales.
- ▶ Modern precision cosmology involves using a combination of data, theory and inference to study.
 - ▶ Its geometry (spatial size, shape and age),
 - ▶ Its evolution (origin, dynamics & fate),
 - ▶ Its contents (matter, radiation, neutrinos, dark matter & dark energy),
 - ▶ ... and the interplay between the above.
- ▶ Interestingly, there was never any Newtonian cosmology.
- ▶ This is completely possible – Galilean relativity allows for an infinite space filled with relatively expanding matter.
- ▶ However it took the innovation of having spacetime as a dynamical quantity for physicists to start thinking about cosmology physically.



You are here






You are here





You are here

INTERSTELLAR NEIGHBORHOOD




You are here

MILKY WAY GALAXY



You are here

LOCAL GALACTIC GROUP



You are here


VIRGO SUPERCLUSTER

LOCAL GALACTIC GROUP




You are here

LOCAL SUPERCLUSTERS




VIRGO SUPERCLUSTER

LOCAL GALACTIC GROUP




You are here


OBSERVABLE UNIVERSE




LOCAL SUPERCLUSTERS



VIRGO SUPERCLUSTER



- ▶ Start by looking again at the concepts underlying the **Friedmann-Robertson-Walker** metric, discussed last year in Part II.
- ▶ Given our **Gauss Theorema Egregium** approach of using the curvature of 2d subsurfaces, we can quickly find both the metric and dynamical equations.
- ▶ We will then move on to consider what we can infer from the metric itself — this includes the concept of **redshift**, Hubble's law, and the important concept of the measurement of distances in the universe.
- ▶ Right: Distribution of galaxies in space as observed by the SLOAN digital sky survey (SDSS).



- ▶ Our fundamental cosmological assumption is that on the largest scales, the universe satisfies the **Extended Copernican Principle**:
- ▶ All positions in the universe are equivalent for the purposes of cosmology.
- ▶ This implies *homogeneity*.
- ▶ Approximation to a metric describing the whole universe – spatial part completely isotropic and homogeneous.
- ▶ Work at a constant time – what is the most general 3-space metric with complete isotropy and homogeneity?
- ▶ To get isotropy we use spherical coordinates and require

$$ds_{r=\text{const}}^2 = r^2(d\theta^2 + \sin^2 \theta d\phi^2) \equiv r^2 d\Omega.$$

- ▶ To get homogeneity, but allow for a true curvature of space, we write (**Lecture 3**)

$$ds^2 = f(r)dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2) \equiv f(r)dr^2 + r^2 d\Omega, \quad (1)$$

- ▶ $\sqrt{f(r)}dr$ is the proper distance between neighbouring points (r, θ, ϕ) and $(r + dr, \theta, \phi)$. r is just a label which we use and is defined by how we write the metric.

- ▶ For spatial homogeneity $f(r)$ must give constant curvature $K(r)$.
- ▶ This was discussed in [Lecture 3](#) in the context of trying to achieve a 2-surface with constant curvature everywhere, using Gauss's formula for curvature. We may immediately use the result obtained before:

$$K = \frac{f'}{2f^2r} \quad \Rightarrow \quad f(r) = \frac{1}{1 - Kr^2}. \quad (2)$$

- ▶ K is the Gaussian curvature of the 2-dimensional surface $\theta = \pi/2$.
- ▶ Isotropy implies this should be independent of the specific choice of coordinate system, and equations (1) and (2) provide a metric which is genuinely homogeneous over the whole of 3-d space.

What sort of spaces are these?

$K = 0$ (flat universe)

- ▶ This is just ordinary Euclidean space $ds^2 = dr^2 + r^2 d\Omega$.

$K > 0$ positive (closed universe)

- ▶ The area of the sphere $r = \text{const}$ is just $A = 4\pi r^2$, but what *actual* distance from the origin does the coordinate distance r correspond to?
- ▶ This is given by the *proper* interval $a(r)$


$$a(r) = \int_0^{a(r)} ds = \int_0^r \frac{dr}{\sqrt{1 - Kr^2}} = \frac{1}{\sqrt{K}} \sin^{-1}(r\sqrt{K}) \quad \Rightarrow \quad r = \frac{1}{\sqrt{K}} \sin(a\sqrt{K}). \quad (3)$$

- ▶ The proper distance is what we would physically measure if we could simultaneously lay down metre sticks over the cosmic distances required.

- From (3) the proper area of a sphere with proper radius a is

$$A = (4\pi/K) \sin^2(a\sqrt{K}).$$

- For small spheres, with $a \ll 1/\sqrt{K}$, $A \approx 4\pi a^2$ – the Euclidean value.
- As a increases we get a maximum $A_{\max} = 4\pi/K$.
- A decreases as we continue to increase a .
- Compare: circumference of a circle of latitude on a sphere $C = 2\pi R \sin(a/R)$.
- These universes are spheres with three-dimensional surfaces, and hence termed closed.



$K < 0$ negative (open universe)

- ▶ The same integration here gives

$$r = \frac{1}{\sqrt{-K}} \sinh(a\sqrt{-K}), \quad A = \frac{4\pi}{|K|} \sinh^2(a\sqrt{|K|}).$$

- ▶ The area thus now increases *faster* than in flat space, and tends to ∞ as $a \rightarrow \infty$.
- ▶ This case is called an **open** universe, or *hyperbolic* space.
- ▶ So we know the spatial part of the metric. What do we need to get a **time** part?


Derivation of time part of FRW metric

- ▶ The fundamental observation is that the universe is expanding, and therefore the metric is changing in time.
- ▶ This expansion is captured by *Hubble's Law* (Hubble, 1929 — *predicted* by Weyl, 1923).

- ▶ The law is:

$$v = H_0 d$$



- ▶ i.e. galaxies are systematically moving away from us with speed proportional to distance.
- ▶ Will discuss how one could actually *define* the quantities involved in this later.
- ▶ This is consistent with a uniform expansion



- Infer recession velocity from looking at spectra (recall quasars, Lecture 10)
- Identifiable pattern of emission (or absorption) lines (H, Mg, Ca, Na, etc. all give optical lines).
- Express this quantitatively via the redshift z defined by

$$z = \frac{\lambda_{\text{obs}} - \lambda_{\text{em}}}{\lambda_{\text{em}}}.$$

- Note that other definitions of redshift would be possible (e.g. dividing by λ_{obs}) — the one given is the one used in cosmology.
- If we choose to interpret this as a (non-relativistic) Doppler shift, then $v \approx zc$ (since $\nu_{\text{obs}} = \nu_{\text{em}}(1 - v/c)$).




- ▶ So the Hubble effect, is that for cases where we know the distance d to the object (e.g. as provided by Cepheids) then we find

$$v \approx zc = H_0 d.$$

- ▶ H_0 is **Hubble's Constant**.
- ▶ Exact value has continued to be controversial, and over the last 30 years or so has spanned the range $H_0 = 50 - 80 \text{ kms}^{-1} \text{Mpc}^{-1}$.
- ▶ (Will discuss up-to-date determinations later.)
- ▶ $1 \text{ Mpc} = 3.0857 \times 10^{22} \text{ m}$, so translated into SI units this is $H_0 = 1.6 - 2.6 \times 10^{-18} \text{ s}^{-1}$.
- ▶ This has big implications for the time evolution of the universe.

- Consider a galaxy a distance d from us, e.g. M101 which is $\approx 4 \text{ Mpc}$ away.




- Projecting back in time, when would it have been on top of us?

$$v = 4 \times H_0 \text{ kms}^{-1} \implies t = \frac{d}{v} = \frac{4 \text{ Mpc}}{4 \text{ Mpc} \times H_0} = H_0^{-1} \text{ s}$$

- I.e. t_{collide} is approximately:
 - 15 billion years (byr) for H_0 in the middle of the above range;
 - 12.5 byr if $H_0 = 80 \text{ kms}^{-1} \text{ Mpc}^{-1}$;
 - and 20 byr if $H_0 = 50 \text{ kms}^{-1} \text{ Mpc}^{-1}$.

- ▶ Clearly this time is independent of which galaxy we are considering, and thus the whole of the universe must have been in a very small volume this time ago.
- ▶ But the age of the Earth is known to be $\sim 4.5 \text{ byr}$ and the oldest stars in the Galaxy are $\sim 11 \text{ byr}$ old.
- ▶ This poses great problems for higher values of H_0 , since the universe is then in danger of being younger than some of its constituents!
- ▶ (If include dynamics, find t_{collide} is modified somewhat, and is most likely even shorter than we have just deduced!)
- ▶ In any case, it is clear that the universe must have come from an earlier stage at which it was much smaller and denser.
- ▶ Roughly, conservation of matter tells us that $\rho \propto R^{-3}$, where ρ is the matter density and R some characteristic size — the **scale factor** — associated with the universe — we will now look at how to define this physically.



- ▶ Must preserve isotropy and homogeneity at each instant of time for the 3-space geometry corresponding to spatial sections.
- ▶ Only possible change with time is for the curvature to vary $K = K(t)$.
- ▶ Since $K \sim 1/(\text{radius of space})^2$, we define a *scale factor* $R(t)$ via

$$K(t) = \frac{k}{R^2(t)} \begin{cases} k = +1 & \text{if curvature positive,} \\ k = 0 & \text{if space is flat,} \\ k = -1 & \text{is curvature is negative.} \end{cases}$$

- ▶ Define a dimensionless coordinate $\sigma = r/R(t)$, then

$$\frac{dr^2}{1 - Kr^2} + r^2(d\theta^2 + \sin^2 \theta d\phi^2),$$


- ▶ becomes

$$R^2(t) \left\{ \frac{d\sigma^2}{1 - k\sigma^2} + \sigma^2(d\theta^2 + \sin^2 \theta d\phi^2) \right\}.$$

- ▶ Corresponds in time to a metric which is just a simple scaling of a fixed 3-metric.
- ▶ Objects with a fixed σ strictly obey the overall uniform expansion:
 - ▶ they have no *peculiar motion* of their own,
 - ▶ on the scales we are thinking of would correspond to galaxies with no peculiar motion,
 - ▶ called '**fundamental**', or '**comoving**' observers – f.o.'s for short.
- ▶ How do we distinguish fundamental observers? – And what did we mean by 'time' when we referred to wanting the same curvature over the whole of a given spacelike slice at the same instant?
- ▶ Not obvious how we can define an absolute time covering the whole universe, which observers can agree upon, and at successive instants get the same answers for curvature, density etc.
- ▶ There *is* an overall rest-frame, and therefore choice of time axis.
- ▶ It is provided by the **cosmic microwave background radiation (CMBR)**!
- ▶ We can detect our motion relative to it via the **dipole anisotropy**


Fundamental Observers

Fundamental observers are those with zero velocity with respect to the frame defined by the microwave background radiation




The COBE Satellite, showing the main instruments


- Mollweide projection of the Earth's surface. Astro plots the same principle, but looking up.




- ▶ Image of the (galactic aligned) milky way sky in the infrared




- ▶ Image of the sky in microwaves. Blackbody spectrum $T = 2.725$ K




- Subtract off this constant term $T - 2.725$ to reveal dipole




- Subtract off dipole to reveal anisotropy




- Subtract off galaxy to reveal primordial anisotropy from the beginning of the Universe




- ▶ Replace COBE with WMAP



- ▶ Replace WMAP with Planck





Maps made by COBE. Top: The sky 'monopole'; Middle: what's left after subtraction of the monopole; Bottom: after subtraction of monopole and dipole

Cosmic time

- ▶ Cosmic time corresponds to the *proper time* (in the usual special relativistic sense) as measured by these fundamental observers.
- ▶ Having seen that for comoving observers the interval of cosmic time, dt say, that they will measure, is just the interval of proper time ds , we have that the total metric for the universe, i.e. with space and time parts joined together, is

$$ds^2 = c^2 dt^2 - R^2(t) \left\{ \frac{d\sigma^2}{1 - k\sigma^2} + \sigma^2(d\theta^2 + \sin^2 \theta d\phi^2) \right\}.$$

- ▶ In the exercises you will be asked to show that comoving observers follow *geodesics* of this metric, which is called the **Friedmann-Robertson-Walker metric**.
- ▶ This means that comoving observers are **freely falling**, and can apply all the usual SR laws in their local frame.

Final form of the FRW metric

- ▶ Most of our information in cosmology comes from radial propagation of photons.
- ▶ Therefore define a coordinate transformation which puts the FRW metric into a form which is especially convenient for this.
- ▶ Introduce a new comoving radial coordinate χ :

$$d\chi = \frac{d\sigma}{\sqrt{1 - k\sigma^2}}, \quad \text{i.e. } \chi = \begin{cases} \sin^{-1} \sigma & k = +1, \\ \sigma & k = 0, \\ \sinh^{-1} \sigma & k = -1. \end{cases}$$

- ▶ The overall FRW metric then becomes

$$ds^2 = c^2 dt^2 - R^2(t) \{ d\chi^2 + S^2(\chi)(d\theta^2 + \sin^2 \theta d\phi^2) \},$$

where $S^2(\chi) = \begin{cases} \sin^2 \chi & k = +1, \\ \chi^2 & k = 0, \\ \sinh^2 \chi & k = -1. \end{cases}$

- ▶ Have arrived at this after two radial coordinate transformations (first $r \rightarrow \sigma$, then $\sigma \rightarrow \chi$), and is essentially our final form.

Propagation of photons — the redshift

- ▶ We now apply the FRW metric to photon propagation to deduce the redshift formula.
- ▶ Suppose a photon is travelling to us from an external galaxy. Its motion is radial, so we can ignore the θ and ϕ coordinates.
- ▶ The condition for photon propagation is that the interval should vanish. Thus


$$ds^2 = c^2 dt^2 - R^2(t) d\chi^2 = 0 \quad \Rightarrow \quad \frac{d\chi}{dt} = -\frac{c}{R(t)},$$

for an incoming photon. This is how its χ coordinate changes with time.

- ▶ Suppose the photon was emitted at cosmic time t_1 and is received by us at time t_0 .
- ▶ (Conventionally t_0 denotes the time here now.) The χ coordinate of the emitter is given by

$$\chi_1 = \int_{t_1}^{t_0} \frac{c dt}{R(t)}.$$

- ▶ The crucial point is that if the galaxy can be treated as a fundamental observer (i.e. moving with the cosmic fluid without a peculiar velocity of its own), then its χ coordinate is fixed.
- ▶ This enables us to work out the redshift.



If the interval between emission of successive wavecrests is δt_1 and that between reception is δt_0 , we have

$$\chi_1 = \int_{t_1}^{t_0} \frac{c dt}{R(t)} = \int_{t_1 + \delta t_1}^{t_0 + \delta t_0} \frac{c dt}{R(t)} = \left(\int_{t_1 + \delta t_1}^{t_1} + \int_{t_1}^{t_0} + \int_{t_0}^{t_0 + \delta t_0} \right) \frac{c dt}{R(t)}.$$

Thus

$$\int_{t_1}^{t_1 + \delta t_1} \frac{c dt}{R(t)} = \int_{t_0}^{t_0 + \delta t_0} \frac{c dt}{R(t)}.$$

- We now approximate by assuming that $R(t)$ hardly changes between emission of successive crests, i.e. that

$$\frac{\dot{R}(t)}{R(t)} \gg \delta t, \quad \text{i.e. "age of universe" } \gg \text{"period of oscillation of wave".}$$

(This is one of the best approximations you are likely to make in physics!)

- In this case


$$\frac{\delta t_1}{R(t_1)} = \frac{\delta t_0}{R(t_0)}.$$

- Now frequency $\nu = 1/\delta t = c/\lambda$, and hence $\lambda_0/\lambda_1 = R(t_0)/R(t_1)$.
- With redshift following our usual definition of $(\lambda_{\text{obs}} - \lambda_{\text{em}})/\lambda_{\text{em}}$, we derive

$$z = \frac{(\lambda_0 - \lambda_1)}{\lambda_1} = \frac{R(t_0)}{R(t_1)} - 1,$$

- i.e. $1 + z = \frac{R(t_0)}{R(t_1)} = \frac{\text{scale factor of universe on reception}}{\text{scale factor of universe when emitted}}$

- ▶ This coincides nicely with our conception of **redshift** as being due to '*stretching*' of the photon wavelength:



- ▶ Can also derive it from the fact that radially p_χ is conserved, so identifying $p^\mu = (\hbar\omega, \hbar\omega/R(t_1))$, and conserving $p_\chi = R(t_1)\hbar\omega_1$ at t_1 , we find that at time t_0 that $p^\chi = \frac{R(t_1)}{R(t_0)^2}\hbar\omega_1 = \hbar\omega_0/R(t_0)$, so $1 + z = \frac{\lambda_0}{\lambda_1} = \frac{\omega_1}{\omega_0} = \frac{R(t_0)}{R(t_1)}$.
- ▶ To get much further with applying the FRW metric, we will need to know the time history of $R(t)$, i.e. we need to know something about the 'dynamical history' of the universe.
- ▶ Thus next in cosmology we will aim to get to grips with **the basic dynamical equations**

Summary

- ▶ Flat, closed and open universes.
- ▶ Fundamental observers, the CMB dipole, and cosmic time.
- ▶ The Friedmann-Robertson-Walker metric.

$$ds^2 = c^2 dt^2 - R(t)^2 dx^2,$$

$$dx^2 = d\chi^2 + S^2(\chi) d\Omega,$$

$$S^2(\chi) = \begin{cases} \sin^2 \chi & k = +1, \\ \chi^2 & k = 0, \\ \sinh^2 \chi & k = -1. \end{cases}$$

- ▶ The redshift formula

$$1 + z = \frac{R(t_0)}{R(t_1)}.$$

Next time

The dynamics of the universe from the Einstein Field equations.