

Final Review

Andrew Wang

August 13, 2024

1 Counting

Theorem. (Binomial Theorem) Let $x, y \in \mathbb{R}$ and $n \in \mathbb{N}$. Then

$$(x + y)^n = \sum_{0 \leq i \leq n} \binom{n}{i} x^i y^{n-i}.$$

Theorem. (Multinomial Theorem) Let $a_1, \dots, a_n \in \mathbb{R}$ and $n \in \mathbb{N}$. Then

$$(a_1 + \dots + a_k)^n = \sum_{m_1 + \dots + m_k = n} \binom{n}{m_1, \dots, m_k} a_1^{m_1} \dots a_k^{m_k}.$$

2 Probabilities

Definition. The *conditional probability* of E given F is

$$\mathbb{P}(E|F) = \frac{\mathbb{P}(E \cap F)}{\mathbb{P}(F)}$$

Definition. Two events E and F are said to be *independent* if

$$\mathbb{P}(E \cap F) = \mathbb{P}(E)\mathbb{P}(F).$$

Theorem. (Bayes' Formula)

$$\mathbb{P}(E|F) = \frac{\mathbb{P}(F|E)\mathbb{P}(E)}{\mathbb{P}(F)}.$$

3 Random Variables

Definition. The *variance* of a real valued random variable X is given by

$$\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = \mathbb{E}[(X - \mathbb{E}[X])^2].$$

Definition. Two random variables X, Y are said to be independent if

$$\mathbb{P}(X = a, Y = b) = \mathbb{P}(X = a)\mathbb{P}(Y = b)$$

for all a, b .

Proposition. Let X, Y be two independent random variables. then:

- (a) $g(X), f(Y)$ are independent for all functions g, f on X, Y .
- (b) $\mathbb{E}[g(X)h(Y)] = \mathbb{E}[g(X)]\mathbb{E}[f(Y)]$.
- (c) $\text{Var}(g(X) + f(Y)) = \text{Var}(g(X)) + \text{Var}(f(Y))$

4 Poisson Random Variables

Definition. X is a *binomial random variable* X with parameter $p \in (0, 1)$ and $n \in \mathbb{N}$ is given by

$$\mathbb{P}(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}$$

for some $k \in \{1, 2, \dots, n\}$.

5 Moment Generation Functions

Definition. For a real-valued random variable X , its *moment-generating function* is $M_X : \mathbb{R} \rightarrow [0, \infty]$,

$$M_X(t) = \mathbb{E}[e^{tX}], t \in \mathbb{R}$$

Proposition.

- (a) $M_X(0) = 1$.
- (b) $M_{X+Y} = M_X M_Y$ for X, Y independent,
- (c) M_X is fully determined by the moments of X so long as $\mathbb{E}[X^n] \leq (cn)^n$ for some $c > 0$.

Definition. For random variables X_1, \dots, X_n , they have moment generating function $M_{X_1, \dots, X_n}(t_1, \dots, t_n) = \mathbb{E}[e^{t_1 X_1 + \dots + t_n X_n}]$.

Definition. X_1, X_2, \dots, X_n are *multivariate normal random variable* if for any $X_i \in X_1, \dots, X_n$ there exist independent normal random variables Y_1, Y_2, \dots, Y_m and $a_{ij} \in \mathbb{R}$ such that

$$\sum_{0 \leq j \leq m} a_{ij} Y_j = X_i.$$

6 Conditional Expectations and Inequalities.

Definition. For a set S and $A \subset S$, the *indicator/characteristic function* of A is

$$\mathcal{X}_A(x) = \begin{cases} 1, & x \in A, \\ 0, & x \notin A. \end{cases}$$

Proposition. (Chevyshev's Inequality) For $a > 0$ and a real-valued random variable X ,

$$\mathbb{P}(|X - \mathbb{E}[X]| \geq a) \leq \frac{\text{Var}(X)}{a^2}.$$

Proposition. (Markov's Inequality) For $a > 0$, and a real-valued random variable $X \geq 0$,

$$\mathbb{P}(X \geq a) \leq \frac{\mathbb{E}[X]}{a}.$$

Definition. The *conditional expectation* of X given Y is a random variable T with

$$\mathbb{P}(T = s) = \sum_{y \in Y: \mathbb{E}[X|Y=y]=s} \mathbb{P}(Y = y)$$

and denoted by $\mathbb{E}[X|Y]$. Similarly, when X, Y are jointly continuous,

$$\mathbb{E}[X|Y = y] = \frac{1}{f_Y(y)} \int_{\mathbb{R}} x f(x, y) dx$$

$$\mathbb{P}(T \in I) = \int_{S(I)} f(y) dy$$

where $S(I) = \{y \in \mathbb{R} : \mathbb{E}[X|Y = y] \in I\}$

Proposition.

- (a) $\mathbb{E}[X] = \mathbb{E}[\mathbb{E}[X|Y]]$,
- (b) $\mathbb{E}[Xg(Y)|Y] = g(Y)\mathbb{E}[X|Y]$.

7 Law of Large Numbers

Definition. X_n converges in probability to X if for any $\varepsilon > 0$,

$$\lim_{n \rightarrow \infty} \mathbb{P}(|X_n - X| \geq \varepsilon) = 0$$

denoted by $X_n \xrightarrow{p} X$.

Definition. X_n converges almost surely to X if

$$\mathbb{P}\left(\lim_{n \rightarrow \infty} X_n = X\right) = 1.$$

denoted by $X_n \xrightarrow{a.s.} X$.

Theorem. (Weak and Strong Law of Large Numbers) Suppose $\{X_n\}$ are i.i.d random variables with $X_n \xrightarrow{d} X, \mathbb{E}[X] < \infty$, then

$$\lim_{n \rightarrow \infty} \frac{X_1 + X_2 + \cdots + X_n}{n} \xrightarrow{p \text{ \& a.s.}} \mathbb{E}[X].$$

8 Central Limit Theorem

Definition. $X_n \in \mathbb{R}$ converges in distribution to $X \in \mathbb{R}$ if

$$\lim_{n \rightarrow \infty} \mathbb{P}(X_n \leq a) = \mathbb{P}(X \leq a)$$

for all $a \in \mathbb{R}$ at which the CDF of X is continuous. This is denoted as

$$X_n \xrightarrow{d} X.$$

Theorem. (Central Limit Theorem) Suppose $X_n \in \mathbb{R}$ are i.i.d. with $\mathbb{E}[X^2] < \infty, \text{Var}(X) > 0$. Then

$$\frac{X_1 + \cdots + X_n - n\mathbb{E}[X]}{\sqrt{n \cdot \text{Var}(X)}} \Rightarrow \mathcal{N}(0, 1).$$